

Open Peer Review on Qeios

Stage IVA Thyroid Gland Follicular Carcinoma AJCC v7

National Cancer Institute

Source

National Cancer Institute. <u>Stage IVA Thyroid Gland Follicular Carcinoma AJCC v7</u>. NCI Thesaurus. Code C115035.

Stage IVA includes: (T4a, N0, M0); (T4a, N1a, M0); (T1, N1b, M0); (T2, N1b, M0); (T3, N1b, M0); (T4a, N1b, M0). T4a: Moderately advanced local disease. Tumor of any size extending beyond the thyroid gland capsule to invade subcutaneous soft tissues, larynx, trachea, esophagus, or recurrent laryngeal nerve. T1: Tumor size 2 cm or less in greatest dimension, limited to the thyroid gland. T2: Tumor more than 2 cm but not more than 4 cm in greatest dimension, limited to the thyroid gland. T3: Tumor more than 4 cm in greatest dimension and limited to the thyroid gland, or tumor of any size with minimal extrathyroid extension (e.g., extension to sternothyroid muscle or perithyroid soft tissues). N0: No regional lymph node metastasis. N1a: Metastasis to pretracheal, paratracheal or prelaryngeal/Delphian lymph nodes (Level VI lymph nodes). N1b: Metastasis to unilateral, bilateral or contralateral cervical (Levels I, II, III, IV, or V) or retropharyngeal or superior mediastinal lymph nodes (Level VII). M0: No distant metastasis. (AJCC 7th ed.)

Qeios ID: 0Z99N3 · https://doi.org/10.32388/0Z99N3