

[Open Peer Review on Qeios](#)

Stage IVA Hypopharyngeal Squamous Cell Carcinoma AJCC v7

National Cancer Institute

Source

National Cancer Institute. *Stage IVA Hypopharyngeal Squamous Cell Carcinoma AJCC v7*. NCI Thesaurus. Code C6132.

Stage IVA includes: (T 4a, N0, M0); (T 4a, N1, M0); (T 1, N2, M0); (T 2, N2, M0); (T 3, N2, M0); (T 4a, N2, M0). T 4a: Tumor with moderately advanced local disease. Tumor invades the thyroid/cricoid cartilage, hyoid bone, thyroid gland, or central compartment soft tissue. Central compartment soft tissue includes prelaryngeal strap muscles and subcutaneous fat. T 1: Tumor limited to one subsite of hypopharynx and/or 2 cm or less in greatest dimension. T 2: Tumor invades more than one subsite of the hypopharynx or an adjacent site, or measures more than 2 cm but not more than 4 cm in greatest diameter without fixation of hemilarynx. T 3: Tumor measuring more than 4 cm in greatest dimension or tumor with fixation of hemilarynx or extension to esophagus. N0: No regional lymph node metastasis. N1: Metastasis in a single ipsilateral lymph node, 3 cm or less in greatest dimension. N2: Tumor with metastasis in a single ipsilateral lymph node, more than 3 cm but not more than 6 cm in greatest dimension, or in multiple ipsilateral lymph nodes, none more than 6 cm in greatest dimension, or in bilateral or contralateral lymph nodes, none more than 6 cm in greatest dimension. M0: No distant metastasis. (AJCC 7th ed.)