

[Open Peer Review on Qeios](#)

Stage II Mycosis Fungoides AJCC v7

National Cancer Institute

Source

National Cancer Institute. *Stage II Mycosis Fungoides AJCC v7*. NCI Thesaurus. Code C7798.

Stage II includes: IIA (T 1-2, N1-2, M0, B0-1); IIB (T3, N0-2, M0, B0-1). T1: Limited patches, papules, and/or plaques covering less than 10% of the skin surface. May further stratify into T1a (patch only) vs. T1b (plaque +/- patch). T2: Patches, papules, or plaques covering 10% or more of the skin surface. May further stratify into T2a (patch only) vs. T2b (plaque +/- patch). T3: One or more tumors (equal or greater than 1 cm diameter). N0: No clinically abnormal peripheral lymph nodes; biopsy not required. N1: Clinically abnormal peripheral lymph nodes; histopathology Dutch grade 1 or NCI LN0-2. N2: Clinically abnormal peripheral lymph nodes; histopathology Dutch grade 2 or NCI LN3. M0: No visceral organ involvement. B0: Absence of significant blood involvement: 5% or less of peripheral blood lymphocytes are atypical (Sezary cells). B1: Low blood tumor burden: more than 5% of peripheral blood lymphocytes are atypical (Sezary cells) but does not meet the criteria of B2. (AJCC 7th ed.)