

[Open Peer Review on Qeios](#)

Stage II Prostate Cancer AJCC v7

National Cancer Institute

Source

National Cancer Institute. *Stage II Prostate Cancer AJCC v7*. NCI Thesaurus. Code C89233.

Stage II includes: IIA (T 1a-c, N0, M0, PSA less than 20, Gleason 7); (T 1a-c, N0, M0, PSA equal or more than 10 and less than 20, Gleason equal or less than 6); (T 2a, N0, M0, PSA less than 20, Gleason equal or less than 7); (T 2b, N0, M0, PSA less than 20, Gleason equal or less than 7); (T 2b, N0, M0, PSA X, Gleason X); IIB (T 2c, N0, M0, Any PSA, Any Gleason); (T 1-2, N0, M0, PSA equal or more than 20, Any Gleason); (T 1-2, N0, M0, Any PSA, Gleason equal or more than 8). T 1a: Tumor incidental histologic finding in 5% or less of tissue resected. T 1b: Tumor incidental histologic finding in more than 5% of tissue resected. T 1c: Tumor identified by needle biopsy (e.g., because of elevated PSA). T 2a: Tumor involves one-half of one lobe or less. T 2b: Tumor involves more than one-half of one lobe, but not both lobes. T 2c: Tumor involves both lobes. cN0: No regional lymph node metastasis. pN0: No positive regional nodes. M0: No distant metastasis. Gleason 7: Moderately differentiated (moderate anaplasia). Gleason Equal or Less than 6: Well differentiated (slight anaplasia). Gleason equal or more than 8: Poorly differentiated/undifferentiated (marked anaplasia). (AJCC 7th ed.)