

Estimating the Fish Stocks of Six Commercial Species in the Middle Madeira River, Southwest Amazon

Igor Hister Lourenço¹, Marcos André Braz Vaz², Marcelo Rodrigues dos Anjos¹

¹ Federal University of Amazonas

² Universidade Federal de Santa Catarina

Funding: This research was funded by the 2020 to 2021 Biodiversa and Water JPI joint call for research projects, under the BiodivRestore ERA-NET Cofund (GA no. 101003777), with the EU and the funding organisations ANR, FCT, DFG, FUNDECT, and FAPEAM (through the Chamada Transnacional Conjunta BiodivRestore 2020-2021, “Conservação e restauração de ecossistemas degradados e sua biodiversidade, incluindo o foco nos sistemas aquáticos”—Resolução FAPEAM no. 016/2020), through the project Forest Fisher,” and by the CNPq, through the projects BACIAS (CNPq/MCTI/FNDCT no. 39/2022 call), and Banzeiro da Educação (CNPq/MCTI no. 55/2022 call). The authors would also like to thank Mastering Degree Grant provided by FAPEAM to the first author through the Resolução No. 003/2019—POSGRAD UFAM—Edição 2019, developed in the Programa de Pós-Graduação em Ciências Ambientais (PPGCA), at the Instituto de Educação, Agricultura e Ambiente (IEAA) from the Universidade Federal do Amazonas (UFAM).

Potential competing interests: No potential competing interests to declare.

Abstract

This manuscript evaluates the fish stocks of six commercially significant species in the middle Madeira River, Southwest Amazon, by leveraging refined fisheries data and an adapted catch effort method. Through daily monitoring at the Humaitá station from May 2018 to April 2019, it examines the catchability and stock estimates for *Semaprochilodus insignis*, *Mylossoma aureum*, *Mylossoma duriventre*, *Prochilodus nigricans*, *Brycon amazonicus*, and *Pseudoplatystoma fasciatum*. Despite the observed low model adjustments and the influence of capture site diversity, significant p-values indicate less than 7% exploitation of the estimated fish populations, suggesting under-exploration and potential for sustainable growth in the region’s fishing activities. However, the study emphasizes cautious interpretation due to unaccounted factors like seasonality, highlighting the necessity for longer data series and consideration of Amazonian peculiarities in future research. This effort contributes to understanding the sustainable management and conservation needs of Amazonian fish resources amidst environmental and human impacts.

Keywords: Amazon fish stocks, Catch effort method, Fish population assessment, Madeira River, Sustainable fishing.

1. Introduction

Fishing is a crucial activity in the Amazon region, providing income, food security, and leisure to local populations (MEGGERS, 1977). This activity is also the primary and most easily accessible source of protein for people living in the region (CERDEIRA et al., 1997; BATISTA et al., 2004). Given its importance, recent studies have emphasized the need to discuss its sustainability and responsible management (SANTOS & SANTOS, 2005; COUX et al., 2010; SANTOS,

FERREIRA & VAL, 2010; WELCOMME, 2011; PRYSTHON, 2021).

These studies highlight a critical problem that must be addressed immediately - the lack of refined data about fisheries. This limitation restricts the use of advanced analyses. However, recent studies have introduced new and feasible methodologies to generate more refined fishery monitoring data (Lourenço et al., 2020). These studies have also provided fresh perspectives on the Amazon's fishery resources in light of increased exploration (Lourenço et al., 2024) and large hydroelectric projects' impact on fishery resources (Lourenço et al., 2023).

Among the possibilities opened with a more refined fisheries database, fish stock estimations deserve a prominent position, especially when considering the fish stock depletions of several commercial species reported across the Amazon in the last decades (PETRERE, 1983; ALAN et al., 2005; CASTELLO et al., 2014; ROBINSON et al., 2020; SU et al., 2021).

Several methods are available to estimate fish stocks in inland waters (FRITZGERALD, DELANTY & SHEPHARD, 2018). When choosing, it is crucial to consider several factors that can influence the decision-making process. Hoyle et al. (2024) argue that despite the attractiveness of more powerful models, such as Structural Topic Models (STMs), more straightforward approaches should be used before exploring complex models.

A recent study by Silva Júnior et al. (2017) in Amazon estimated fish stocks using an adapted version of the catch effort method proposed by Mao (2007). While the approach seems feasible, due to the lack of refined data in the region, the authors violated a statistical assumption in Mao's method by considering the total biomass instead of the total number of individuals in the regressions, which led to partially unreliable, or at least overestimated results.

This study uses the adaptation made by Silva Júnior et al. (2016) to Mao's (2007) method using refined fisheries data produced by the Lourenço et al. (2020) method to estimate fish stocks of six commercial species in the middle Madeira region, Southwest Amazon.

2. Material and Methods

2.1. Study area and data collection

We gathered the data used in this study from May 2018 to April 2019 by daily monitoring the landings that arrived at the monitoring station in Humaitá, Amazonas, Brazil, middle Madeira River (GIBBS, 1967; McCLAIN et al., 1995; GOULDING et al., 2003; QUEIROZ et al., 2013), following the method proposed by Lourenço et al., (2020).

We selected six commercial fish species that were landed more than 50 times during the sampled period, namely *Semaprochilodus insignis*, *Mylossoma aureum*, *Mylossoma duriventre*, *Prochilodus nigricans*, *Brycon amazonicus*, and *Pseudoplatystoma fasciatum*.

2.2. Statistical analyses

To estimate the fish stocks of the six selected species, we applied the adaptation of Silva Júnior et al. (2016) of Mao's (2007) method, but using number of captured individuals instead of total captured biomass. This adaptation is based on the hypothesis that a population "N" that undergoes a sampling process "g" with a standardized effort "f" has a catch probability that is related to time "t". The catchability "π" is determined by:

$$g_{\pi}(t) = p_q(f_t) \prod_{k=1}^{t-1} \{1 - p_q(f_k)\}$$

where: "q" is the catchability, " $p_q(f_t)$ " represents the probability of an individual be captured given an amount of effort "f", and the " $\prod_{k=1}^{t-1} \{1 - p_q(f_k)\}$ " represents the probability that it was not captured in previous events.

That way, the remaining population, following the catch probability of each individual is provided by "θ":

$$\theta = \frac{1 - \sum_{k=1}^W g_q(k)}{\sum_{k=1}^W g_q(k)}$$

Considering "n" individuals captured, the likelihood estimator of the original population is given by:

$$B = b + b\theta$$

where: "b" is the number of captured individuals, and "B" is the total population.



Finally, the function that relates the catch probability is linearly related to effort and parameterized by catchability:



$$p_q(f) = qf$$

Basically, the method involves counting and removing individuals from a population while applying a given effort repeatedly. An estimative of catchability, known as "q", is determined based on the trend line obtained from the regression between the CPUE and cumulated effort. By replacing the captured individuals' number, effort, and catchability values in Equation 4, it is possible to estimate the fish stock of the analyzed population. In other words, the population size is estimated by modeling the decrease of catchability after successive tries (MAO, 2007).

3. Results

The regressions obtained are presented next (Figure 1). The tendency lines presented a downward bias for *S. insignis*, *M. aureum*, *M. duriventre*, and *F. fasciatum* and an upward bias for *P. nigricans* and *B. amazonicus*.









Figure 1. Regressions between the CPUE and the cumulated effort of six commercial species landed in Humaitá, Amazonas, from May 2018 to April 2019.

Despite the low adjustment of the models (r^2), the regression's p-values were all significant (Table 1). In addition, by analyzing the total estimated population for each species, we observe that less than 7% is currently being explored, complicating the interpretation of the tendency lines.

Table 1. Number of captured individuals, mean effort, catchability, regression's p-value between CPUE and cumulated effort, estimated population, and percentage of explored population of six commercial fish species landed in Humaitá, Amazonas, from May 2018 and April 2019.

Species	N Individuals	Mean effort	Catchability	p-value	Estimated population	Explored %
<i>Semaprochilodus insignis</i>	6380	4.52	-5.20E-03	1.49E-02*	271734.64	2.35
<i>Mylossoma aureum</i>	9108	6.48	-2.80E-03	2.07E-02*	502195.49	1.81
<i>Mylossoma duriventre</i>	8079	8.60	-4.20E-03	4.22E-04*	223671.10	3.61
<i>Prochilodus nigricans</i>	1914	9.10	7.50E-03	1.14E-03*	28058.64	6.82
<i>Brycon amazonicus</i>	1259	8.30	2.50E-03	9.81E-03*	60707.95	2.07
<i>Pseudoplatystoma fasciatum</i>	583	4.59	-1.07E-02	2.19E-04*	17879.37	3.26

Eforço: homens.dia pescados⁻¹; *: significativo com intervalo de confiança de 95%.



Rendimento Sustentável (MSY). Adaptado de Lorenzen et al. (2016)

4. Discussion

The low adjustment of the models can be explained by the inclusion of different capture sites and, consequently, environments. The ideal would be to generate one regression for each capture site separately; however, given the small historical series, there are not enough entries to do so.

As Lorenzen et al. (2016) pointed out, the tendency lines can hold critical information about the exploration levels of each species. Theoretically, given a certain amount of cumulated effort, the regression's tendency lines (our capturability index) will tend to go up until we reach our maximum sustainable yield (MSY) (Figure 2). After reaching the MSY, the catchability will tend to decrease, even if we continue to increase the effort, marking the point where fishermen should stop the activity and give time for the stocks to recover (Martell and Froese, 2012).

In light of this, our results suggest that the species *P. nigricans* and *P. punctifer* still haven't reached the MSY; in other words, the more effort we apply to capture these species, the more the catch. Parallely, this logic can be applied to the

species that presented a downward tendency line, suggesting an exceeding of the MSY.

However, we must interpret these results carefully since we didn't consider the effect of the seasonality over the catch, which has a close relation with the hydrological seasons (SOUSA et al., 2017; LOPES & FREITAS, 2018), a fact consolidated in the studies developed by Issac, Milstein & Ruffino (1996) and Barthem & Fabré (2003). When we take into consideration, for instance, the capture of the genera *Mylossoma* over the year (LOURENÇO, ANJOS E BARREIROS, 2020), the catchability will naturally tend to decline, since the data used (starting in May 2018) coincides with the beginning of the captures, reaching a peak around September, and tending to decline again after that. The same logic can be applied to the other analyzed species, not necessarily meaning that they reached the MSY.

A more accurate analysis would require a longer data series, preferably several years, so we can plot the Lorenzen et al. (2016) parable and have a better glimpse into the conservation status of the analyzed species. A longer data series would also allow us to isolate the effects of the seasonality and environment over the estimates, reinforcing the importance of the continuity of the monitoring method employed here.

The results raised a few questions: What is the support capacity of fish stocks in Amazon inland waters? How much can we extract without compromising the activity and the ecosystem services? How much time do we need to let the stocks rest? At first, the information generated here suggests that the activity can still grow since the number of captured individuals didn't reach 7% of the total population.

When comparing the catch from our study with the catch data from the closest capital in the Madeira River, Porto Velho (approximately two hundred kilometers upstream from Humaitá), we can see that the production here is almost inexpressive (DORIA & LIMA, 2008; DORIA et al., 2012; DORIA, LIMA & ANGELINI, 2018)—opening space for the discussions around the under-exploration of the region, and what factors are influencing it.

Finally, it is worth mentioning that the method developed by Mao (2007) doesn't consider the seasonality, diversity of environments, migration cycles, specificities of the explored species, and fishing gear used, among other factors not considered in this study. Further studies considering these nuances are required for a better understanding of the exploration of this resource and, much likely, adaptations to Mao's (2007) method which will cover the Amazonian peculiarities. Future studies aiming to incorporate each species size are currently under construction and welcoming collaborators and orientations.

References

- ALAN, J. D.; ZBELL, R.; HOGAN, Z.; REVENGA, C.; TAYLOR, B. W.; WELCOMME, R. L.; DINEMILLER, K. Overfishing of Inland Waters. *BioScience*, v. 55, n. 12, p. 1041-1051, 2005.
- Barthem, R. B.; Fabré, N. N. Biologia e diversidade dos recursos pesqueiros da Amazônia. In: Ruffino, Mauro Luís (Org.). *A pesca e os recursos pesqueiros na Amazônia Brasileira*. Manaus: Ibama/ ProVárzea. p.17-59, 2003.
- BATISTA, V. S.; ISSAC, V. J.; VIANA, J. P. "Exploração e manejo dos recursos pesqueiros da Amazônia". In: Ruffino,

- M. L. (ed.). A pesca e os recursos pesqueiros na Amazônia brasileira. Pro Várzea. p. 63-152, Manaus, 2004;
- CASTELLO, L.; ARANTES, C. C.; MCRATH, D. G.; STEWART, D. J. SOUSA, F. S. Understanding fishing-induced extinctions in the Amazon. *Aquatic Conservation: Marine and Freshwater Ecosystems*, v. 25, n. 5, p. 587-598, 2015. <https://doi.org/10.1002/aqc.2491>.
 - CERDEIRA, R. G. P.; RUFFINO, M. L.; ISAAC, V. J. "Consumo de pescado e outros alimentos pela população ribeirinha do lago grande de Monte Alegre, PA. Brasil". *Acta Amazonica*. v. 27, p. 213-228, 1997.
 - COUX, I. G.; ALMEIDA, O.; BENE, C.; BRUMMETT, R.; BUSH, S.; DARWALL, W.; PITTOCK, J.; VAN BRAKEL, M. A. valoração da pesca em águas continentais. *Novos Cadernos NAEA*, v. 13, n. 1, p. 71-103, 2010.
 - DORIA, C. R. C.; LIMA, M. A. L. A pesca do pacu (Cuvier, 1818) (Characiformes: Characidae). *Revista Biotema*, v. 21, n. 3, 2008.
 - DORIA, C. R. C.; RUFFINO, M. L.; HIJAZI, N. C.; CRUZ, R. L. A pesca comercial na bacia do rio Madeira no estado de Rondônia, Amazônia Brasileira. *Acta Amazonica*, v. 42, n. 1, p. 29-40, 2012.
 - DORIA, C. R. C.; LIMA, M. A. L.; ANGELINI, R. Ecosystem indicators of a small-scale fisheries with limited data in Madeira River (Brazil). *Boletim do Instituto de Pesca*, v. 44, n. 3, e317, 2018. <https://dx.doi.org/10.20950/1678-2305.2018.317>.
 - FRITZGERALD, C. J.; DELANTY, K.; SHEPHARD, S. Inland fish stock assessment: Applying data-poor methods from marine systems. *Fisheries Management and Ecology*, v. 25, n. 4, p. 240-252, 2018. <https://doi.org/10.1111/fme.12284>.
 - GIBBS, R. J. The geochemistry of the Amazon River system. Part I. The factors that control the salinity and the composition and concentration of the suspended solids. *Bull. Geol. Soc. Am.* v. 78, p. 1203-1232, 1967.
 - GOULDING, M.; BARTHEM, R.; FERREIRA, E. The Smithsonian atlas of the Amazon. Smithsonian Books, pp. 253, Washington D. C., 2003.
 - HILBORN, R.; WALTERS, C. J. Quatitative fisheries stock assessment and management: choice, dynamics and uncertainty. Chapman and Hall, pp. 570, New York, 1992.
 - HILBORN, R. The state of the art in stock assessment: where we are and where we are going. *Scientia Marina*, v. 67, p. 15-20, 2003.
 - ISAAC, V. J.; MILSTEIN, A.; RUFFINO, M. L. A pesca artesanal no baixo Amazonas: Análise multivariada da captura por espécie. *Acta Amazonica*, v. 26, n. 3, p. 185-208, 1996.
 - LIMA, F. C. T. & CAIRES, R. A. Peixes da Estação Ecológica Serra Geral do Tocantins, bacias dos rios Tocantins e São Francisco, com observações sobre as implicações biogeográficas das "águas emendadas" dos rios Sapão e Galheiros. *Biota Neotropical*, v. 11, n. 1, p. 231-250, 2011.
 - LOPES, G. C. S.; FREITAS, C. E. C. Avaliação da pesca comercial desembarcada em duas cidades localizadas no rio Solimões – Amazonas. *Biota Amazônia*, v. 8, n. 4, 2018. <http://dx.doi.org/10.18561/2179-5746/biotaamazonia.v8n4p36-41>
 - LORENZEN, K.; COWX, I. G.; ENSUA-MENSAH, R. E. M.; LESTER, N. P.; KOEHN, J. D.; RANDALL, R. G.; SO, N.; BONAR, S. A.; BUNNELL, D. B.; VENTURELLI, P.; BOWER, S. D.; COOKE, S. J. Stock assessment in inland fisheies: a foundation for sustainable use and conservation. *Reviews in Fish Biology and Fisheries*, v. 26, p. 405-440, 2016. <https://dx.doi.org/10.1007/s11160-016-9435-0>.

- LOURENÇO, I. H.; ANJOS, M. R.; BARREIROS, J. P. Low-cost technology for fish monitoring applied to the fishing of two species of pacu in Amazonas, Brazil. *Boletim do Instituto de Pesca*, v. 46, n. 2, 2020. <https://doi.org/10.20950/1678-2305.2020.46.2.585>.
- MARTELL, S.; FROESE, R. A simple method for estimating MSY from catch and resilience. *Fish and Fisheries*, v. 14, n. 4, p. 504-514, 2012. <https://doi.org/10.1111/j.1467-2979.2012.00485.x>.
- MCCLAIN, M. E.; RIKEY, J. E.; VICTORIA, R. I. Andean contributions to the biogeochemistry of the Amazon River system. *Bulletin de l'Institut Français d'Etudes Andines*, v. 24, n. 3, p. 1-13, 1995.
- MAO, C. X. Estimating population sizes by catch-effort methods. *Statistical Methodology*, v. 4, p. 111-119, 2007.
- MEGGERS, B. Amazônia: a ilusão de um paraíso, *Civilização Brasileira*. p. 207, Rio de Janeiro, 1977;
- PETRERE JR, R. Yield per recruit of the Tambaqui, *Colossoma macropomum* Cuvier, in the Amazonas State, Brazil. *Journal of Fish Biology*, v. 22, p. 133-144, 1983.
- QUEIROZ, L. J.; TORRENTE-VILARA, G.; VIEIRA, F. G.; OHARA, W. M.; ZUANON, J.; DORIA, C. R. C. Fishes of Cuniã Lake, Madeira River Basin, Brazil. *Check List-Journal of species and distribution*, v. 9, n. 3, p. 540-548p, 2013.
- PETRERE JUNIOR, M. Pesca e esforço de pesca no Estado do Amazonas. *Acta Amazonica*, v. 8, n. 3, p. 439-454, 1978. <http://dx.doi.org/10.1590/1809-43921978083439>.
- PRYTHON, A. Revisitando a Pesca Artesanal na Embrapa. Um “Oceano azul” de oportunidades ou um “oceano vermelho” de ameaças? *Research, Society and Development*, v. 10, n. 3, p. e45010313572, 2021. DOI: <http://dx.doi.org/10.33448/rsd-v10i3.13572>.
- ROBINSON, J. P. W.; ROBINSON, J.; GERRY, C.; GOVIDEN, R.; FRESHWATER, C.; GRAHAM, N. A. J. Diversification insulates fisher catch and revenue in heavily exploited tropical fisheries. *Science Advances*, v. 6, eaaz0587, 2020.
- SALES, M. K. G.; FONSECA, R.; ZANCHI, F. B.; SANTOS, S. Caracterização do desembarque pesqueiro efetuado na colônia de pescadores Z-31 no município de Humaitá-AM. *Revista Igapó*, v. 5, n. 1, Jun, 2011.
- SANTOS, G. M.; SANTOS, A. C. M. Sustentabilidade da pesca na Amazônia. *Estudos Avançados*, v. 54, p. 165-182, 2005.
- SANTOS, G. M.; FERREIRA, E. J. G.; VAL, A. L. Recursos pesqueiros e sustentabilidade na Amazônica: Fatos e perspectivas. *Hiléia: Revista do Direito Ambiental*, n. 8, p. 43-77, 2010.
- SILVA JÚNIOR, U. L. S.; RASEIRA, M. B.; RUFFINO, M. L.; BATISTA, V. S.; LEITE, R. G. Estimativas do Tamanho do Estoque de algumas Espécies de Peixes Comerciais da Amazônia a partir de Dados de Captura e Esforço. *Biodiversidade Brasileira*, v. 7, n. 1, p. 105-121, 2017.
- SOUSA, T. G. C.; SOUZA, L. A.; FRUTUOSO, M. E.; FREITAS, C. E. C. Seasonal dynamic of Amazonian small-scale fisheries is dictated by the hydrologic pulse. *Boletim do Instituto de Pesca*, v. 43, n. 2, p. 207-221, 2017. <https://www.dx.doi.org/10.20950/1678-2305.2017v43n2p207>.
- SU, G.; LOGEZ, M.; XU, J. TAO, S.; VILLÉGER, S.; BROSSE, S. Human impacts on global freshwater fish biodiversity. *Science*, v. 371, n. 6531, p. 835-838, 2021. <https://www.dx.doi.org/10.1126/science.abd3369>.
- WELCOMME, R. L. A review of a model for qualitative evaluation of exploitation levels in multi-species fisheries. *Fisheries Management, and Ecology*, v. 6, p. 1-19, 1999.

- WELCOMME, R. L.; An overview of global catch statistics for inland fish. ICES Journal of Marine Science, v. 68, n. 8, p. 1751-1756, 2011.