## **Research Article**

# Evaluation of Chemical Content and Phytochemical Composition of Yemeni Almond Cultivars

Yaser Aljawfi<sup>1</sup>, Muneer Mohammed Saleh Alsayadi<sup>2,3</sup>, Mansour Al dalas<sup>4</sup>, Hazem Alashwal<sup>4</sup>, Ibrahim Al hakami<sup>4</sup>, Sharifah Aminah Syed Mohamad<sup>5</sup>, Nor Hadiani Ismail<sup>5</sup>, Mufeda Ahmed Hazea Gazaem<sup>6</sup>

Independent researcher;
 Atta-ur-Rahman Institute for Natural Products Discovery, Universiti Teknologi Mara, Malaysia;
 Department of Food Science and Technology, Faculty of Agriculture, Ibb University, Ibb, Yemen;
 Faculty of Pharmacy, Sana'a University, Sana'a, Yemen;
 Faculty of Applied Sciences, Universiti Teknologi Mara, Malaysia;
 Universiti Teknologi Mara, Malaysia;

The almond plant is considered one of the most important crops economically, nutritionally, and health-wise; its properties and chemical composition differ according to the places of its cultivation. This study aims to estimate the chemical and phytochemical content of some local cultivars of almonds in Yemen. The chemical components, minerals, and secondary metabolites were estimated in the seeds. The results of the chemical analysis showed that the studied cultivars of Yemeni almonds contained a higher percentage of nutrients compared to the imported samples. The percentages of moisture, protein, fat, carbohydrates, and ash were 2.88–3.39%, 12.7–19.15%, 36.87–65.34%, 13.23– 38.02%, and 2.55–3.56%, respectively, and they contained a high content of some minerals such as iron, zinc, copper, and potassium. The total phenols in the studied almond cultivars ranged from 9.9– 113 mg Gallic acid/g, and the total flavonoids ranged from 47.7–63.00 mg Rutin/g. It is concluded from the study that Yemeni almonds are characterized by unique properties that promise multiple uses in the food and pharmaceutical industries and in the prevention and treatment of many diseases.

Corresponding author: Muneer Mohammed Saleh Alsayadi, smuneer006@gmail.com

## Introduction

The seeds of the almond plant are one of the nuts that are a source of high nutritional value, whether for macronutrients (protein, fat, and carbohydrates) or micronutrients (minerals and vitamins), in addition to other vital compounds such as secondary metabolites, including phenolic acids, flavonoids, tannins,

and alkaloids<sup>[11][2][3][4]</sup>. Almonds *Prunusdulcis* have great health and nutritional benefits, as they are a source of protein, polyunsaturated fatty acids, fiber, vitamin E, riboflavin, and essential minerals, in addition to phytosterols and polyphenols<sup>[5]</sup>. Consuming almonds regularly has many health benefits, especially for heart disease, obesity, high blood pressure, and diabetes<sup>[6]</sup>, in addition to having an antioxidant effect against cancer and atherosclerosis<sup>[7]</sup>, and they play a role in enhancing immunity and anti-inflammatory properties<sup>[8]</sup>. The high nutritional value of the almond kernel stems mainly from its high content of fat, which is an important source of calories. Despite this, it does not contribute to raising blood cholesterol due to its high content of unsaturated fatty acids<sup>[9]</sup>. The seeds of the almond plant are one of the nuts that are a source of high nutritional value. Almonds are also considered one of the most produced nuts in the world, as the amount of production worldwide reached 1.2 million tons in 2017. The US state of California is the most productive region, with almond production reaching about 81% of global production, followed by Australia at 7%, Spain at 4%, Iran at 1%, and Tunisia at 1%<sup>[10]</sup>. Almonds were cultivated in the Middle East 4000 years ago, and sweet almonds are an important and valuable specialty crop grown in many temperate and subtropical regions of the world, including South Australia and the United States of America.

In Yemen, almonds are one of the most important economic and monetary crops, and one of the most important crops competing with Qat is that its economic yield is high due to its high price, which is a very large difference compared to imported almonds, due to its high quality. It is considered one of the most important crops that qualify for expansion in its consumption and export levels as a type due to its high quality and its great competitiveness in the domestic and foreign markets. Moreover, its requirements are low as it can be successfully grown in marginal lands and under the rainfed system in areas with an annual rainfall of 200-300 mm or more under rain harvesting systems. Almonds are grown in Yemen on an area of 5,986 hectares, which represents 7% of the area cultivated with fruit crops. Its cultivation is mainly concentrated in Sana'a Governorate, with an estimated area of 5,959 hectares. It is also grown in both the governorates of Dhamar and Taiz, and the production reached 10,483 tons in 2018<sup>[11]</sup>. And the fact that pollination in almonds is mixed, the seed multiplication leads to the production of wide genetic variations, as there are many cultivars of almonds spread in Yemen, including pure cultivars bearing local designations that differ among themselves in quality, in phenotypic traits, and environmental needs. Almond cultivars also vary in their content of chemical compounds or minerals due to genetic factors as well as different environmental conditions and agricultural treatments, in addition to the ripeness of the fruits at harvest time<sup>[6]</sup>, and this is the most important factor affecting quality. The lack of a taxonomic key and a national record for the different varieties or breeds to preserve the rights and the national identity of these varieties/cultivars, especially for a national crop such as almonds, is one of the most important problems that should be overcome. In addition to that, the scarcity of studies and research concerned with characterizing the local genetic origins of almonds, whether phenotypic, chemical, or molecular characterization, as well as the scarcity of studies and research concerned with studying the chemical content of Yemeni almonds and whether the chemical content of Yemeni almonds is richer than others in compounds and minerals and has earned it high quality, means that such studies may contribute significantly to the issue of promoting and enhancing the competitiveness of Yemeni almonds locally and regionally. Therefore, this study aimed to characterize the phenotypic traits and estimate the chemical and mineral content and phytochemical composition of the most famous cultivars of Yemeni almonds.

## **Materials and Methods**

## Study area and sample sources

The study was conducted during June and July 2021, and plant tree samples were studied in the two regions of Khirbet Mohib in BaniMatar District and Jabal Al Lawz in Khawlan District, Sana'a Governorate, Yemen. The most prevalent cultivars were identified in the two regions with the help of almond farmers in the studied areas; three cultivars were identified in the BaniMatar region and six cultivars in the Khawlan region. The sources, geographical location data, and local names of the studied samples are listed in Table 1.

Sample	Cultivars Local	District	Region	Height	Geographical location	
NU	name		(111)	longitude	Latitude	
1	Large fit Razky		Khirbet Mohib - Jabal al- NabiShuaib	2877	43.98916	15.32038
2	Middle Razky	BaniMatar	Khirbet Mohib	2853	43.98186	15.3121
3	Small Raziki		Khirbet Mohib	2814	43.2036	15.31213
4	Large Razky		Murbk - Jabal Al-Lawz - Al- Tayyal	2684	44.2867	15.2334
5	Twined Jelly		Murbk- tayyal	2664	44.88634	15.23383
6	Shahty	Khawlan	Murbk- Jabal Al-Lawz - Al- Tayyal	2691	44.28669	15.2339
7	Rounded Razky		Murbk- Jabal Al-Lawz - Al- Tayyal	2687	44.2871	15.2337
8	FlatRazky		Murbk- Al-tayyal	2677	44.28635	15.2339
9	Oblong Razky		Murbk- Al-tayyal	2691	44.28713	15.23369

Table 1. The sources, geographical location data, and the local names of almond samples

## Almond seed sample collection and preparation

Nine samples of almond seeds, each weighing 1 kg, were collected from each of the identified trees. They were kept in sterile polyethylene bags, then transported to the laboratory, peeled, and kept in sterile polyethylene bags at 4 °C until use. They were transported to laboratories and preserved in the same way for chemical analysis. Almond seed samples were ground and preserved in sterile polyethylene vacuum-sealed bags in the dark at 4°C and used for analysis on the same day. The local cultivars used in this work (Fig. 1)



Figure 1. The variation of the studied cultivars of almonds in some of the phenotypical characteristics of the almond: 1) large fit Razky, 2) Medium Razky, 3) Small Razky, 4) Large Razky, 5) Rounded Gly, 6) Shahty, 7) Rounded Razky, 8) Flat Razky, 9) An oblong Razky

## Chemical analysis

#### Estimation of the chemical composition of almond seed samples

The almond seed content of moisture, protein (Keldahl method), total fat (Sacholite method), and ash were estimated by the Association of Official Analytical Chemists AOAC methods<sup>[12][13][14][15]</sup>.

## Determination of the mineral content of almond seeds

Minerals (Ca, Mg, Na, Al, Fe, Cu, Zn, Mn, and Boron B) were determined in almond seeds using ICP emission spectrometry according to the method used by the Association of Official Analytical Chemists AOAC No. 985.01.

#### Samples extraction and preparation for phytochemical analysis

The cold method extraction method was used for each sample. Fifty grams of seeds were milled, then extracted with 250 ml of methanol for 24 hours with continuous shaking. Then the extracts were filtered through Whatman paper No. 4, and the solvent was removed using a rotary evaporator at 40 °C. The extracts were then kept at 5 °C until the total phenols and flavonoids determination<sup>[13]</sup>.

## Determination of the phytochemical content of almond seeds

Secondary metabolite compounds (total phenols and flavonoids) were determined in almond seed extracts.

## Determination of Total Phenols

Total phenols were determined by the Folin–Ciocalteu reagent colorimetric method<sup>[16][13][14]</sup>. Briefly, 1 ml of sample extract (0.1%, w/v), 0.5 ml of Folin–Ciocalteu reagent (1:2 v/v), and 2 ml of 5% sodium carbonate were vortexed and allowed to stand at 30°C for 1 h. Absorbance was measured with a UV-VIS spectrophotometer at a wavelength of 765 nm. The total phenolic content of almond extracts was expressed as gallic acid equivalents per g of dry weight (mg GAE/g).

## Determination of Total Flavonoids

The total flavonoid contents (TFC) of the samples were determined using the aluminum chloride method that was previously described  $\frac{[17]}{1}$  with few modifications. Briefly, the sample extract (1.0 gm) was mixed

with distilled water (4 ml), followed by sodium nitrite (0.3 ml), and after 5 min, aluminum chloride (3 ml) solution was added. After 6 min, NaOH (2 ml, 1.0 M) and distilled water (10 ml) were added at 25°C. The absorbance was determined by a spectrophotometer at 510 nm. The results were expressed as rutin equivalents (mg RE/g sample).

## Statistical Analysis

The mean, standard deviation, and correlation coefficient for the results of the chemical and phytochemical composition of almond seeds were estimated. ANOVA was performed using One Way ANOVA and Tukey Tests, and significant differences were calculated at the level of  $P \le 0.05$  using SPSS V. 21.

## **Results and Discussions**

## **Chemical Content**

#### **Basic Chemical Compounds**

Table 2 shows the chemical composition of the studied almond samples. It is noted that the moisture content ranged from 2.88% for the sample of Small Razaki to 3.39% in the sample of "Shahty." This result is consistent with what was found by <sup>[18]</sup> while studying samples of Turkish almonds. It should be noted that the humidity rate is suitable for preserving almond kernels for long periods. On the other hand, the protein content ranged from 12.7% in a sample of flat Razaki almonds from the Khawlan region to 19.57% in a sample of large proportionate Razaki from the BaniMatar region, and these results are considered less than what was found by <sup>[18]</sup> in Turkish almond samples and Sylvia et al. <sup>[5]</sup> in samples of American almonds from California, both of which reached about 20%. In contrast, these results are consistent with what was found by <sup>[19]</sup> when studying samples of Greek almonds, where it ranged between 10–29%, while <sup>[20]</sup> and <sup>[19]</sup> found that the percentage of protein in Italian almond samples ranged between 16-25%, and <sup>[21]</sup> found that the percentage of protein in Spanish almond samples from France, Italy, and Greece, the protein content ranged between 10–29%. This was indicated by <sup>[20]</sup>, <sup>[23]</sup>, and <sup>[19]</sup>, who indicated that the chemical composition has an important role in the difference between almond samples in protein ratios.

almond cultivars	Moisture%	Protein (N×5.18)%	Total Fat%	Total Ash%	Carbohydrates%
Large fit Razky	3.38±0.05	19.57±0.30	44.88± 1.05	3.2±0.01	28.97±1.25
Middle Razky	3.39±0.04	17.6±0.78	37.95±1.02	3.56±0.95	37.46±1.73
Small Raziki	2.88±0.12	15.97±0.83	56.46±1.06	3.02±0.95	21.65±0.99
Large Razky	3.44±0.05	19±0.81	36.87±1.11	2.7±0.02	38.02±1.98
Twined Jelly	2.91±0.05	14.6±1.77	43.84±1.30	2.55±0.01	36.09±0.47
Shahty	3.73±0.06	18.88±0.43	53.53±1.09	3.27±0.33	20.57±0.81
Rounded Razky	3.27±0.05	16±0.2	64.71±1.13	2.84±0.03	13.23±0.90
Flat Razky	2.92±0.02	12.7±0.51	65.34±1.15	2.7±0.01	16.18±0.73
Oblong Razky	2.95±0.01	15.64±0.15	52.55±0.67	2.68±0.02	26.18±0.73

Table 2. Chemical content of protein, total fat, ash, and carbohydrates of the studied almond cultivars

From the results in Table 2, it is clear that the percentage of fat varied between the different samples, as it ranged from 36.87% in the Large Razqi from the Khawlan region to 65.34% and 64.71% in each of the almond samples, Razqi flat and Razqi round, respectively. It should be noted here that there is a worldwide variation in the fat content of almonds, ranging between 25–67%. In a study conducted by <sup>[24]</sup> on some samples of Turkish almonds, the fat content ranged between 25 and 65%, while in other studies on Spanish almonds, <sup>[25][26][27][28][29][30][31]</sup>, and <sup>[32]</sup>, the percentage of fat ranged between 40 and 67%, and in a study by Sylvia et al.<sup>[5]</sup> on American almonds from California, the percentage of total fat ranged between 49 and 50%. It should also be noted that the high nutritional value of the almond kernel arises mainly from its high-fat content, which is an important source of calories. Despite this, it does not contribute to raising blood cholesterol due to its high content of unsaturated fatty acids<sup>[9]</sup>.

As for the percentage of total ash, it ranged between 2.55% in the Jali Mabroum sample from the Khawlan region and 3.56% in the average Razqi sample from the BaniMatar region (Table 2). These results agree with what was found by <sup>[33]</sup> and <sup>[20]</sup>, where the percentage of total ash ranged between 2.3-3.7%. As for the carbohydrate content, it ranged in a relatively wide range between 13.24% in the sample of Razqi

Medawar and 38.02% in the sample of Razaki Kabeer, as shown in Table 2, and by comparing these results with the results of other studies, <sup>[34]</sup> found that the percentage of total carbohydrates ranged from 14% to 21%, while <sup>[35]</sup> found in another study that the proportion of carbohydrates reached between 23.5 to 27%, as also found by <sup>[36]</sup> that the proportion of carbohydrates was up to 28%.

## Mineral contents

Table 3 shows the results of mineral analysis in the studied almond cultivars. It is noted that the highest value was in the cultivar of Razqi large, which originated from the Khawlan region, where it amounted to 6.207 mg/100 grams, followed by each of the cultivars of Razqi medium and Razqi small from the BaniMatar region, which amounted to 5.419 and 5.003 mg/100 grams, respectively, followed by the two cultivars Razqi large matcher from BaniMatar and Shahti from Khawlan, with values of 3.572 and 3.415 mg/100 grams, respectively. The previous values were higher than the value achieved by the control, which amounted to 3.378 mg/100 grams, while the lowest value for zinc content was in the Razqi flat sample, reaching 2.814 mg/100 g. These results indicate that the zinc content of Yemeni almonds was high in three samples: large, medium, and small, while the rest of the samples were within the accepted range, according to international studies, which ranges between 2-4 mg/100 grams<sup>[9]</sup>. This is considered a comparative advantage for some types of Yemeni almonds over external almonds, which have a higher nutritional value and more health benefits.

The same is the case concerning the content of the iron element in the samples. The highest value was found in a medium Razqi sample, which came from BaniMatar, with a value of 7.261 mg/100 grams, followed by a small Razqi sample from BaniMatar as well, then a rectangular Razqi sample from Khawlan, with values of 6.622 and 5.001 mg/100 grams, respectively, compared to the lowest value in the Yemeni almond samples, which amounted to 4.221 mg/100 grams in the Jelly Mabroum sample. These results show that three samples of Yemeni almonds—medium Razqi, small Razqi from BaniMatar, and oblong Razqi from Khawlan—had their content of iron higher than the accepted range in the results of international studies of external almond samples, which ranged between 2.6 and 4.9 mg/100 grams<sup>[9]</sup>.

As for the copper component, the highest samples in their copper content were a medium Razqi and a small Razqi, both from the BaniMatar region, where the values of their copper content were 1.844 and 1.731 mg/100 grams, respectively, followed by a matching large Razqi sample from the BaniMatar region, also with a value of 1.731 mg/100 grams. These three samples were higher than the control sample, which amounted to 1.417 mg/100 grams, while the two samples of Razqi Round and Razqi Kabir were the least in

their content of copper, as they reached 0.917 and 0.948 mg/100 grams, respectively (Table 3). According to what was reported by Rafel et al.<sup>[9]</sup>, the copper content reported in many international studies ranged between 0.9 and 1.3 mg/100 g. From this, it is clear that Yemeni almonds are distinguished by their high content of copper compared to other types of almonds.

As for the manganese mineral, it is noted from Table 3 that its value ranged between 2.271 and 1.232 mg/100 grams in each of Shahti and Razqi flats, respectively. The studied samples were higher in copper content, except for flat Razqi, Jali twisted, and oblong Razqi, all of which were from the Khawlan region. As indicated by Rafel et al.<sup>[9]</sup> regarding the recognized range of manganese, it ranges between 1.2 and 3.04 mg/100 g, and all studied samples were within this range.

almond gultivare	Zn	Fe	Cu	Mn
		mg/1	Cu           mg/100g           3         1.731           31         1.844           32         1.765           39         0.948           21         0.974           34         1.124           39         0.917           29         1.142           31         1.081	
Large fit Razky	3.572	4.393	1.731	1.735
Middle Razky	5.419	7.261	1.844	1.798
Small Raziki	5.003	6.622	1.765	1.913
Large Razky	6.207	4.969	0.948	1.883
Twined Jelly	3.196	4.221	0.974	1.611
Shahty	3.415	4.804	1.124	2.271
Rounded Razky	3.305	4.789	0.917	1.809
Flat Razky	2.814	4.429	1.142	1.232
Oblong Razky	3.303	5.001	1.081	1.585

Table 3. Content of studied almond cultivars of some trace minerals (zinc, iron, copper, manganese) inmg/100g

Concerning the microelements in Table 4, the potassium content of the studied samples ranged between 2000 and 1130 mg/100 in each of the middle Razqi samples from the BaniMatar region and the

rectangular Razqi sample from the Khawlan region, which was less than the control sample, which amounted to 1200 mg/100 grams, and it should be noted here that the content of the studied samples was higher than the range reported by Rafel et al.<sup>[9]</sup>, which ranges from 430 to 940 mg/100 g. As for the element calcium, the range was relatively narrow, as it ranged between 87.71 and 105.80 mg/100 grams for Razky Round and Razky Small, respectively. As for the magnesium element, it ranged between 428.76 mg/100 grams in the Razqi Medawar sample from the Khawlan region and 484.52 mg/100 grams in the middle Razqi sample from the BaniMatar region. These results are consistent with what was found by Davide et al.<sup>[6]</sup>. The content of the studied samples of the sodium element ranged between 37.8 and 52.05 mg/100 grams in each of Razqi Kabir from the BaniMatar region and Razqi Rectangle from the Khawlan region, respectively. It should be noted here that the sodium content in all the studied samples was high compared to what was mentioned by Rafel et al.<sup>[9]</sup>, where he showed that the range of sodium in the results of some studies ranged between 1 and 20 mg/100 g.

On the other hand, it is noted from the results shown in Table 4 that the content of the studied samples of the aluminum element ranged between 0.434 and 2.497 mg/100 grams in each of the samples of a large proportional Razqi from the BaniMatar region and a large Razqi sample from the Khawlan region, respectively. It is also noted that the boron content of the samples ranged between 1.398 and 5.055 mg/100 grams in each sample of a large proportional Razqi from the BaniMatar region and a rectangular Razqi from the Khawlan region, respectively. These results are not consistent with what was reported by Cesarettin et al.<sup>[4]</sup>, who found that almonds do not contain aluminum and boron, while Davide et al. <sup>[6]</sup> found that in a study conducted on more than twenty types of almonds, the presence of aluminum and boron in one sample amounted to 0.39 and 1.85 mg/100 g, respectively.

almond gultivara	К	Ca	Mg	Na	Al	В	
almond cultivars	mg/100g						
Large fit Razky	1608	93.35	477.7	52.05	0.434	1.398	
Middle Razky	2000	98.88	484.5	48.3	0.807	2.001	
Small Raziki	1840	105.8	478.8	49.3	0.709	2.078	
Large Razky	1220	95.5	481.6	40.9	2.497	3.572	
Twined Jelly	1200	97.55	436.2	46.7	1.418	2.169	
Shahty	1430	93.44	473.7	44.1	0.632	2.826	
Rounded Razky	1390	87.71	428.8	39.2	0.694	1.883	
Flat Razky	1330	89.9	458.8	45.2	0.861	2.411	
Oblong Razky	1130	100.64	450.8	37.8	0.888	5.055	

**Table 4.** The content of the studied cultivars of almonds of some minerals (potassium, calcium, magnesium,sodium, aluminum, boron) in milligrams / 100 grams

## Phytochemical compounds content

Table 5 shows that the content of total phenolic compounds in the samples ranged between 9.9 and 113 mg of gallic acid/gram extracted in each of the two samples of Razqi Flat from the Khawlan region and RazqiSaghir from the BaniMatar region, respectively. Comparing the local almond samples with the control, it is noted that five samples mathematically outperformed the control sample: large razky, small razqi, large razqi, mabroum jelly, and oblong razqi. These results are consistent with what was found by <sup>[37]</sup>, where it ranged in the studied almond samples between 9.22 and 163.71 mg Gallic acid/gram of extract. Likewise, concerning the content of flavonoids, it is noted from Table 6 that there were only two samples that contained these compounds, a sample of small Razqi and a sample of large proportional Razqi, both from the BaniMatar region, as they reached 63 and 47.7 mg rutin /gram, respectively, while the rest of the samples did not contain flavonoids. Table 5 also shows the percentage of extraction yield, where the highest extraction rate was in the samples of Razqi Flat and Jali Mabroum, which amounted to

8% as the highest value, while the lowest value was in each of the samples of Razqi Medium, Razqi Small, and Razqi Large from BaniMatar, which amounted to 5%. This finding was similar to that found by <sup>[37]</sup>.

Almond cultivars	Total Polyphenols (mg GAE/g)	T Flavonoids (mg rutin/g)	Yield
Large fit Razky	66.5	47.7	5.1
Middle Razky	26.5	-	5
Small Raziki	113	63	5
Large Razky	36.6	-	5.5
Twined Jelly	37.6	-	8
Shahty	18.4	-	5.5
Rounded Razky	9.9	-	7.1
Flat Razky	19	-	8.1
Oblong Razky	33.4	-	7.3

 Table 5. The studied almond cultivar's content of phenolic and flavonoid compounds and the yield of phenolic

 materials extraction

## Results of cluster analysis for chemical content

A cluster analysis was carried out for the characteristics of the aforementioned 17 chemical components after giving each characteristic a score from 1 to 9. Figure 2 shows the results of the cluster analysis, where it is noted that the studied cultivars of almonds differ among themselves according to their variation in chemical content and their content of minerals, phenolic, and flavonoid compounds. The studied cultivars/samples were divided into two main groups. It is also noted that this corresponds to the collection area. The cultivars collected from the BaniMatar region formed one group, and the cultivars collected from the Khawlan region formed another group. This division achieved a high correlation coefficient of 0.9012 at a significant level of 0.011, and this is due to the existence of correlations between some chemical components, which are illustrated in Figure 3. The first group included the "BaniMatar cultivars group" of three cultivars: large compatible Razaki, medium Razaki, and small Razaki. This

group was characterized by a higher content of both ash and protein, carbohydrates, zinc, manganese, iron, copper, calcium, magnesium, sodium, potassium, phenolic compounds, and flavonoids, and a lower content of each of total fats, aluminum, and boron compared to the second group, the "Khulan region group." It is also noted from Figure 2 that there is a discrepancy between the cultivars within the groups, as the Razqi medium cultivars are located at a greater distance than the other two cultivars within the first group as a result of its higher content of some elements, lower content of fats, lower content of phenolic compounds, and lack of flavonoids. On the other hand, the sample Razqi Kabir differed from the rest of the cultivars within the framework of the second group because it contains more protein, carbohydrates, zinc, aluminum, boron, iron, and magnesium and contains less fat.



**Figure 2**. Tree of convergence or divergence between the studied cultivars of almonds according to their chemical and mineral content



**Figure 3.** The correlation between the studied chemical components: the blue color is a positive correlation, the red color is a negative correlation, and the circles surrounded by a square mean that there is a significant correlation at a significant level of 0.05.

Through the results discussed previously, it is noted that there is no relationship between the results of the cluster analysis of the phenotypic traits and the results of the cluster analysis of the chemical and mineral content, as it is likely that most of the chemical components of the almond cultivars are greatly affected by the environmental factor and the crop management factor. The cultivars collected from the BaniMatar region seemed to be richer in their chemical content for most of the chemical components except for fats, aluminum, and boron. Perhaps this is attributed to the environmental factor as well as better crop management processes in the BaniMatar region, and this can be attributed to the intensity of interventions in the BaniMatar region, especially the recent interventions. Within the project of decentralizing water management in the Sana'a Basin, the BaniMatar district, especially the Khirbet Moheib area, was a center for the project's interventions, especially in the almond cultivation sector. However, training programs were accompanied by training programs for farmers in managing the almond crop, such as fertilization, irrigation, and various agricultural operations.

# Conclusion

The results showed that the content of some minerals, such as zinc, copper, iron, and potassium, in some studied cultivars of Yemeni almonds was higher than the range known from international studies conducted on types of external almonds, which means that Yemeni almonds have a higher nutritional value and more health benefits compared to external almonds. The studied cultivars of almonds varied among themselves according to their variation in chemical content and were divided into two main groups according to the collection region. The cultivars collected from the BaniMatar region were characterized by a higher content of ash, protein, carbohydrates, zinc, manganese, iron, copper, calcium, magnesium, sodium, potassium, phenolic, and flavonoid compounds, and the content was lower in each of the total fats, aluminum, and boron compared to the second group, "the group of cultivars of the Khawlan region." Some types of Yemeni almonds have a higher nutritional and health value than the external types of almonds due to their high content of chemical components and minerals. Some of these benefits can be summarized as follows: Yemeni almonds, in some cultivars, contain a high percentage of fat, up to 65%, and are an excellent source of energy. In addition, the quality of the fat present in almonds is of a beneficial type that lowers cholesterol in the blood and preserves the integrity of the heart and blood vessels from sclerosis. The study showed that most of the almond cultivars contain a higher percentage of minerals compared to studies conducted on other types of almonds worldwide. Accordingly, it is recommended to conduct additional studies to evaluate the phytochemical content and identify the components of Yemeni almond varieties and evaluate their antioxidant activity. Their components can be utilized in the production of pharmaceutical products and functional foods due to their health-promoting properties.

## **Statements and Declarations**

## Data Availability Statement

The datasets generated for this study are available on request to the corresponding author.

#### Author Contributions

Conceptualization: Y.A. (Yaser Aljawfi), M.M.S.A. (Muneer Mohammed Saleh Alsayadi); Methodology: Y.A.; Formal Analysis: Y.A.; Investigation: Y.A., M.M.S.A.; Resources: M.M.S.A.; Data Curation: Y.A.; Writing – Original Draft Preparation: Y.A.; Writing – Review & Editing: Y.A., M.M.S.A.; Supervision: M.M.S.A.; Project Administration: M.M.S.A.

# References

- 1. <sup>^</sup>Moshfegh A, Ingwersen I, Goldman J (2007). "Nut Consumption on the U.S. and the Contribution to Nutrie nt Intakes." USDA, ARS, Beltsville, MD.
- 2. <sup>△</sup>Lichtenstein AH, Appel LJ, Brands M, Carnethon M, Daniels S, Franch HA, Franklin B, Kris-Etherton P, Harri s WS, Howard B, Karanja N, Lefevre M, Rudel L, Sacks F, Van Horn L, Winston M, Wylie-Rosett J (2006). "Diet and lifestyle recommendations revision. A scientific statement from the American Heart Association Nutriti on Committee." Circulation. 114:82–96.
- 3. <sup>^</sup>Sabate J, Ros E, Salas-Salvado J (2006). "Nuts: Nutrition and health outcomes." Br J Nutr. 96(Suppl 2):1S−2 S.
- 4. <sup>a</sup>, <sup>b</sup>Alasalvar C, Shahidi F (2009). Tree Nuts Composition, Phytochemicals, and Health Effects. CRC Press, Ta
   ylor & Francis Group. pp. 40, 137.
- 5. <sup>a, b, c</sup>Yada S, Huang G, Lapsley K (2013). "Natural variability in the nutrient composition of California-grow n almonds." J Food Compos Anal. **30**:80–85.
- 6. <sup>a, b, c, d</sup>Barreca D, Nabavi SM, Sureda A, Rasekhian M, Raciti R, Silva AS, Annunziata G, Arnone A, Tenore G C, Süntar İ, Mandalari G (2020). "Almonds (Prunus dulcis Mill. D. A. Webb): A source of nutrients and health -promoting compounds." Nutrients. **12**:672.
- 7. <sup>^</sup>Oliveira I, Meyer AS, Afonso S, Sequeira A, Vilela A, Goufo P, Trindade H, Gonçalves B (2020). "Effects of diff erent processing treatments on almond (Prunus dulcis) bioactive compounds, antioxidant activities, fatty a cids, and sensorial characteristics." Plants. 9:1627.
- 8. <sup>^</sup>Rabadán A, Álvarez-Ortí M, Pardo J (2019). "A comparison of the effect of genotype and weather condition s on the nutritional composition of most important commercial nuts." Sci Horti. **244**:218–224.
- 9. a. b. c. d. e. f. g. h. Sociasi Company R, Gradziel TM (2017). Almonds: Botany, Production and Uses. Wallingford: CAB International. pp. 429–440.
- 10. <sup>△</sup>Franklin LM, Mitchell AE (2019). "Review of the sensory and chemical characteristics of almond (Prunus d ulcis) flavor." J Agric Food Chem. 67:2743–2753.
- 11. <sup>A</sup>AASB (2018). Annual Agricultural Statistics Book. Ministry of Agriculture and Irrigation. pp 24,31.

- 12. <sup>△</sup>AOAC (1997). "Official Methods of Analysis. Analysis of Total Fat. Official Method 948.22." 18th Ed. Gaither sburg, MD: Association of Official Analytical Chemists.
- 13. <sup>a, b, c</sup>Al jawfi Y, Alsayadi M, Binmansor A, Atikbekkar F (2014). "Chemical Composition, Phytochemical Con stituents, and Antioxidant Activities of the Seeds Extract of Apiumgraveoleus L. from Yemen." RJPBCS. 5(1):1 074-1087.
- 14. <sup>a, b</sup>Aljawfi Y, Alsayadi M, Benmansour A, Chaban sari D, Lazoni H (2013). "Chemical and phytochemical an alysis of some antidiabetic plants in Yemen." Int Res J Pharm. 4(9):72-76. doi:<u>10.7897/2230-8407.04915</u>.
- <sup>^</sup>AOAC (2000). "Official Methods of Analysis. Ash of Fruits and Fruit Products. Official Method 948.22." 17th Ed. Gaithersburg, MD: Association of Official Analytical Chemists.
- 16. <sup>△</sup>BinMowyna MN, Alsayadi M (2020). "Assessment of the antioxidant and antimicrobial activities of Carall umadeflersiana growing in the South of Saudi Arabia." African Journal of Pharmacy and Pharmacology. 14 (8):331-338. doi:10.5897/AJPP2020.5183.
- 17. <sup>△</sup>AlMousa LA, AlFaris NA, Alshammari GM, ALTamimi JZ, Alsyadi MM, Alagal RI, Yahya MA (2022). "Antio xidant and antimicrobial potential of two extracts from Capparis spinosa L. and Rumex nervosus and mole cular docking investigation of selected major compounds." Saudi J Biol Sci. **29**(8):103346.
- 18. <sup>a, b</sup>Mustafa N, Nizamlioglu S (2017). "Physical and Chemical Properties of a Type of Almond Called 'Akbade m' Grown in the Aegean Region in Turkey." Int J Sec Metabolite. 4(3):134–141.
- 19. <sup>a, b, c</sup>Drogoudi PD, Pantelidis G, Bacchetta L, De Giorgio D, Duval H, Metzidakis I, Spera D (2013). "Protein an d mineral nutrient contents in kernels from 72 sweet almond cultivars and accessions grown in France, Gre ece, and Italy." International Journal of Food Sciences and Nutrition. **64**:202–209.
- 20. <sup>a, b, c</sup>Ruggeri S, Cappelloni M, Gambelli L, Nicoli S, Carnovale E (1998). "Chemical composition and nutritive value of nuts grown in Italy." Italian Journal of Food Science. **3**:243–251.
- 21. <sup>△</sup>Riquelme F, Romojaro F, Gimenez JL, Llorente S (1985). "Study on protein fraction in some almonds varieti es of the Spanish south-east." Fruit Science Reports. 12:55–59.
- 22. <sup>△</sup>Drogoudi P, Pantelidis G, Bacchetta L, De Giorgio D, Duval H, Metzidakis I, Spera D (2012). "Protein and mi neral nutrient contents in kernels from 72 sweet almond cultivars and accessions grown in France, Greece, a nd Italy." Int J Food Sci Nutr. doi:10.3109/09637486.728202.
- 23. <sup>△</sup>Sathe SK (1993). "Solubilization, electrophoretic characterization, and in vitro digestibility of almond (Pru nusamygdalus) proteins." Journal of Food Biochemistry. **16**:249–264.
- 24. <sup>△</sup>Agar IT, Kafkas S, Kaska N (1998). "Effect of cold storage on the kernel fatty acid composition of almonds."
   Acta Horticulturae. 470:349–358.

- 25. <sup>Δ</sup>García-López C, Grané-Teruel N, Berenguer-Navarro V, García-García JE, Martín-Carratalá ML (1996). "Ma jor fatty acid composition of 19 almond cultivars of different origins. A chemometric approach." Journal of Agricultural and Food Chemistry. 44:1751–1755.
- 26. <sup>△</sup>García-Pascual P, Mateos M, Carbonell V, Salazar DM (2003). "Influence of storage conditions on the qualit y of shelled and roasted almonds." Biosystems Engineering. 84:201–209.
- 27. <sup>△</sup>Kodad O, Sociasi Company R, Prats MS, López Ortiz MC (2006). "Variability in tocopherol concentrations i n almond oil and its use as a selection criterion in almond breeding." Journal of Horticultural Science & Biot echnology. 81:501–507.
- 28. <sup>△</sup>Kodad O, Sociasi Company R (2008). "Variability of oil content and major fatty acid composition in almon d (PrunusamygdalusBatsch) and its relationship with kernel quality." Journal of Agricultural and Food Che mistry. 56:4096–4101.
- 29. <sup>△</sup>López-Ortiz CM, Prats-Moya S, Beltrán Sanahuja A, Maestre-Pérez SE, Grané-Teruel N, Martín-Carratalá ML (2008). "Comparative study of tocopherol homologue content in four almond oil cultivars during two c onsecutive years." Journal of Food Composition and Analysis. 21:144–151.
- 30. <sup>△</sup>Romojaro F, Riquelme F, Giménez JL, Llorente S (1988). "Study on carbohydrate fractions in some almonds cultivars of the Spanish south-east." Fruit Science Reports. **15**:1–6.
- 31. <sup>△</sup>Sánchez-Bel P, Egea I, Martínez-Madrid MC, Flores B, Romojaro F (2008). "Influence of irrigation and orga nic/inorganic fertilization on the chemical quality of almond (Prunusamygdalus cv. Guara)." Journal of Agr icultural and Food Chemistry. 56:10056–10062.
- <sup>A</sup>Soler L, Cañellas J, Saura-Calixto F (1988). "Oil content and fatty acid composition of developing almonds s eeds." Journal of Agricultural and Food Chemistry. 36:695–697.
- 33. <sup>△</sup>Barbera G, Di Marco L, La Mantia T, Schirra M (1994). "Effect of rootstock on the productive and qualitativ e response of two almond varieties." Acta Horticulturae. **373**:129–134.
- 34. <sup>△</sup>Barreira J, Casal S, Ferreira I, Peres A, Pereia J (2012). "Supervised chemical pattern recognition in almond (Prunusdulcis) portugueses PDO cultivars: PCA-and LDA-based triennial study." J Agric Food Chem. 60:969 7-9704.
- 35. <sup>△</sup>Ahrens S, Venkatachalam M, Mistry A, Lapcley K, Sathe S (2005). "almond (Prunusdulcis) protein quality." Plant Foods Hum Nutr. 60:123-128.
- 36. <sup>△</sup>Akpambang V, Amoo I, Izuagie A (2008). "Comparative compositional analysis on two varieties of melon (colocynthiscitrullus and Cucumeropsisedulis) and a variety of almonds." Res J Agric Biol Sci. 4:639-642.

37. <sup>a. b</sup>Barreira JCM, Ferreira ICFR, Oliveira MBPP, Pereira JA (2008). "Antioxidant activity and bioactive compo unds of ten Portuguese regional and commercial almond cultivars." Food and Chemical Toxicology. **46**:223 0–2235.

## Declarations

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.