

# Review of: "Ecological diversity, structure and exploitation of rattan stands according to a disturbance gradient around the Nkoltang forest, Estuary province of Gabon"

Ismael Hernández-Valencia<sup>1</sup>

<sup>1</sup> Universidad Central de Venezuela

Potential competing interests: No potential competing interests to declare.

ASSESSMENT OF THE MANUSCRIPT Ecological diversity, structure and exploitation of rattan stands according to a disturbance gradient around the Nkoltang forest, Estuary province of Gabon

## ABSTRACT

Substitute mining by logging

## INTRODUCTION

- *E. macrocarpa* which are found in all agroecological zones (AEZs) of Cameroon. Indeed, the species *C. deerratus* was found in EAAs 2 and 5 while *L. secundiflorum* and *L. robustum* were found in EAAs 3, 4 and 5. In contrast, *E. wendlandiana* was found only in AEZ 4.

What are EAAs 2 and 5, EAAs 3, 4 and 5, and AEZ4?

## CHOICE OF STUDY SITES

- Rattan, a species of NTFP

What it means NTFP?

- Figure 1.

For homogeneity, you must present a map in English

Sampling was made in rainy or dry season?

The terrain is flat or hilly?

How far are the plots from towns or villages?

## DATA ANALYSIS AND PROCESSING

- The statistical processing was carried out on Ri386 version 3.4.0. The data were subjected to the graphical normality test.

Reference?

## RESULTS

- Figure 2.

The types of environments must be presented in English

### **Distribution of abundance of rattan species as a function of habitats**

- Figure 3

Titles must be presented in English

### **Rattan Cutting Pressure Level in Different Media**

- Figure 4

Titles must be presented in English

### **Regeneration of rattan stands in different environments**

- Figure 5

Titles must be presented in English

### **Abundances of rattan stands as a function of habitats**

- Nevertheless, the gender diversity observed

Substitute gender by genera