

Research Article

Two Intrinsic Formulae Generated by the Jones Polynomial

Brian Jonathan Wolk¹

1. Independent Colleges and Universities of Florida, Tallahassee, United States

This paper notes and derives two simple, intrinsic mathematical relations generated by the Jones polynomial.

Corresponding author: Brian Jonathan Wolk, Brian.Jonathan.Wolk@gmail.com

“Out of clutter, find simplicity.” – John A. Wheeler (on Albert Einstein)¹

1. Introduction

This paper notes two nice and simple intrinsic relations generated by the Jones polynomial for knots that, though the relation's *terms* can be generated by the Skein relations and Kaufmann bracket [\[1\]\[2\]](#) (once the relation is noticed), the relations themselves must be actually perceived and discovered and first noticed in order to then realize that the Skein relations and Kaufmann bracket generate their terms.

This author, in studying knots for the purpose of modeling the elementary particles of physics [\[3\]\[4\]\[5\]\[6\]](#), has not found through due diligence these specific relations recognized or even alluded to in any journal article or book on knots or otherwise by any author. Perhaps they are too trivial to note? Or perhaps they just simply weren't noticed until now. They are thus noted herein and derived, but in a novel way.

I contemplate the polynomials:

$$t^{-2} - t^{-1} + t^0 - t + t^2$$

$$t^2 - t^3 + 2t^4 - 2t^5 + 3t^6 - 2t^7 + t^8 - t^9$$

$$t^3 + t^5 - t^8$$

$$\begin{aligned}
& -t^{-11} + 3t^{-10} - 5t^{-9} + 6t^{-8} - 8t^{-7} + 8t^{-6} - 6t^{-5} + 5t^{-4} - 2t^{-3} + t^{-2} \\
& \quad -t^{-4} + t^{-3} + t^{-2} \\
& \quad t^0 - 2t + 4t^2 - 5t^3 + 6t^4 - 6t^5 + 6t^6 - 4t^7 + 2t^8 - t^9 \\
& \quad \quad -t^{1/2} - t^{5/2} \\
& \quad \quad t^{-5} - 2t^{-4} + 3t^{-3} - 3t^{-2} + 3t^{-1} - 3t^0 - 2t - t^2
\end{aligned}$$

...Clutter. But staring long enough, an utterly simple (and so perhaps profound?) pattern emerges. Two of these polynomials are not like the others; in a very certain specific sense, they are not the same.

One of the polynomials is a Jones polynomial which signals a link with two components, whereas the other polynomials (excepting one other polynomial among them) are Jones polynomials which signal only one-component links. And one of the polynomials isn't a Jones polynomial at all.

We know the distinguished two by simply reading the next sentence. These are the respective Jones polynomials for the 4_1 , the 7_3 , the 8_{19} , the 9_{23} , the trefoil (3_1), the 9_{36} , and the Hopf (2_1^2) links [\[7\]\[1\]\[2\]](#), along with the last polynomial which, we will see, is not even a Jones polynomial.

But what if one wasn't given that information? Can one, by scanning the polynomial and doing a simple operation, determine whether or not it can be a Jones polynomial in the first instance? And if so, then with the same operation, can one determine how many components its signaled link must have? What about the following list:

$$\begin{aligned}
f_1 &= t^{-9} - t^{-8} + 6t^{-7} - 6t^{-6} + 16t^{-5} - 15t^{-4} + 21t^{-3} - 15t^{-2} + 15t^{-1} - 10t^0 + 5t^1 - t^2 \\
f_2 &= -t^{-9} + 3t^{-8} - 5t^{-7} + 6t^{-6} - 7t^{-5} + 7t^{-4} - 6t^{-3} + 4t^{-2} - 2t^{-1} + t \\
f_3 &= -t^{-7/2} - t^{-3/2} - 2t^{-1/2} - 2t^{1/2} - t^{3/2} - t^{7/2} \\
f_4 &= t^{9/2} - 2t^{13/2} + 3t^{15/2} - 3t^{17/2} + t^{21/2} \\
f_5 &= -t^{7/2} - t^{11/2} + t^{13/2} - t^{15/2} + t^{17/2} - t^{19/2} + t^{21/2} - t^{23/2} \\
f_6 &= -t^{-9/2} + 3t^{-7/2} - 5t^{-5/2} + 5t^{-3/2} - 6t^{-1/2} + 5t^{1/2} - 4t^{3/2} + 2t^{5/2} - t^{7/2} ?
\end{aligned}$$

The answer is that f_4 is not a Jones polynomial. All the rest potentially are (and happen to be). How can one tell?

2. The Link Class Invariant

Designate the class containing all links L with the same number n of components as \mathcal{L}_n . We include in \mathcal{L}_n both the link L and its Jones polynomial J . Thus, for example, \mathcal{L}_2 is the class of all Jones polynomials and their referent links having 2 and only 2 components. For instance:

$$7_3 \in \mathcal{L}_1 \quad 2_1^2 \in \mathcal{L}_2 \quad 8_3^4 \in \mathcal{L}_4,$$

with the link L implicitly carrying its Jones polynomial J into the class with it.

Let $\{J\}_{L_n}$ be the sequence of coefficients of the Jones polynomial of any link L with n components. For example, for the Hopf link 2_1^2 we have [7][11]: $\{J\}_{2_1^2} = \{-1, 0, -1\}$. For the 8_1^2 link we have [11]: $\{J\}_{8_1^2} = \{-1, 0, -1, 1, -1, 1, -1, 1, -1\}$. Call $|J|_{L_n}$ the sum of the coefficients of any link L with n components - a member of \mathcal{L}_n :

$$|J|_{L_n} = \sum \{J\}_{L_n} : L_n \in \mathcal{L}_n.$$

The link class conjecture is:

$$|J|_{L_n} = X_n \quad n \in \mathbb{N},$$

where $X_n \in \mathbb{Z}$ and is completely determined and calculable for all n , to be shown infra. Therefore, one can state that any two Jones polynomials J and J' belong to the same \mathcal{L} -class iff $|J| = |J'|$. One can also state that given any polynomial p , if $|p| \neq X_n$ for some $n \in \mathbb{N}$ then p is not a Jones polynomial.

3. The Link Class Iteration

One can sit down and go through all the links with 1 component [7][11][8], and calculate $|J|_{L_1}$ for each. One will find:

$$\sum \{J\}_{L_1} = 1 \quad \forall L \in \mathcal{L}_1.$$

Do this for those links with 2, 3, and 4 components, and one will find:

$$\sum \{J\}_{L_2} = -2 \quad \forall L \in \mathcal{L}_2$$

$$\sum \{J\}_{L_3} = 4 \quad \forall L \in \mathcal{L}_3$$

$$\sum \{J\}_{L_4} = -8 \quad \forall L \in \mathcal{L}_4.$$

That is to say, we have for the Jones polynomial link class:

$$\sum \{J\}_L = (-2)^{n-1} \quad \forall L \in \mathcal{L}_n,$$

or what is the same:

$$|J|_{L_n} = (-2)^{n-1}.$$

If the sum of the coefficients of a purported Jones polynomial does not equal one of the numbers given by (9), then it's not a Jones polynomial. Looking back to (2), one finds that $|f_4| = 0$, and therefore f_4 is not a Jones polynomial. The others are, and are given by [8]:

$$L_{f_1} = L11a548$$

$$L_{f_2} = 9_{20} \quad L_{f_3} = 8_3^4$$

$$L_{f_5} = 8_1^2 \quad L_{f_6} = 8_{10}^2.$$

Likewise, looking back to (1), we see that the last polynomial p has $|p| = -4$, and therefore it cannot be a Jones polynomial.

Equation (9) applies as well if the orientation of the component(s) is changed, applies to alternating and non-alternating L , and applies to unoriented L . For example, reversing the orientation of one component of the 2-component Hopf link gives the following Jones polynomial and coefficient sum:

$$-t^{-5/2} - t^{-1/2} \quad |J|_L = -2.$$

Consider the Jones polynomial for unoriented (u) 7_1^3 [9]:

$$t^{-5/2} - t^{-3/2} + 4t^{-1/2} - 3t^{1/2} + 4t^{3/2} - 3t^{5/2} + 3t^{7/2} - t^{9/2},$$

which is strikingly different from the Jones polynomial for oriented 7_1^3 [11]:

$$-t^{-3} + 3t^{-2} - 3t^{-1} + 4t^0 - 3t^1 + 4t^2 - t^3 + t^4.$$

Nevertheless, we have $|J|_{7_1^3(u)} = 4 = |J|_{7_1^3}$.

Considering the pattern in (6)-(7), it is straightforward to deduce that the sum of the coefficients for any $n: J \in \mathcal{L}_n$ is determinable and can be calculated by utilizing the following iteration:

$$\sum \{J\}_{L_{n+1}} - \sum \{J\}_{L_n} = (-1)^n (2^n + 2^{n-1})$$

$$n \in \mathbb{N}, \quad \sum \{J\}_{L_1} = 1.$$

4. Conclusion

Number (14) is a nice, simple intrinsic relation generated by the Jones polynomial, just as (8) is. Their application? One never knows.

Statements and Declarations

Data Availability

Data sharing is not applicable to this article as no new data were created or analyzed in this study. The work relies on established mathematical definitions and previously published Jones polynomials referenced herein.

Footnotes

¹Cosmic Search Magazine, Vol. 1 No. 4, Forum: John A. Wheeler, From the Big Bang to the Big Crunch, <http://www.bigear.org/vol1no4/wheeler.htm>

References

1. ^{a, b, c, d, e, f}Adams C (2022). *The Knot Book*. Providence: American Mathematical Society.
2. ^{a, b}Kauffman LH (2013). *Knots and Physics*. 4th ed. Hackensack: World Scientific Publishing.
3. ^ΔWolk B (2023). "Building the Standard Model particles and fields within a sphere fiber bundle framework." *Phys. Open*. 15.
4. ^ΔWolk B (2020). "The underlying geometry of the Standard Model of particle physics: $SU(2) \otimes U(1)$." *Int. J. Mod. Phys. A*. 35.
5. ^ΔWolk B (2021). "The underlying geometry of the Standard Model of particle physics: $SU(3)$." *Adv. Appl. Clifford Algebras*. 31.
6. ^ΔBaez J, Muniain JP (2013). *Gauge Fields, Knots and Gravity*. Hackensack: World Scientific Publishing.
7. ^{a, b, c}Doll H, Hoste J (1991). "A tabulation of oriented links." *Mathematics of Computation*. 57:196.
8. ^{a, b}The Knot Atlas. http://www.katlas.org/wiki/Main_Page.

9. [△]Ganzell S, Huffman J, Mavrakis L, Tademy K, Walker G (2019). "Unoriented links and the Jones polynomial." *Involve: A Journal of Mathematics*. **12**(8).

Declarations

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.