

Research Article

On Extensions of Mean Value Theorems for Multi-Function Products under Uniform Monotonicity

Tuan Tran Van¹¹. Ho Chi Minh City University of Science, Ho Chi Minh City, Vietnam

This paper introduces novel generalizations of the Mean Value Theorem for families of functions whose derivatives satisfy specific monotonic conditions. Specifically, the study establishes the existence of a single, common intermediate point $c \in (a, b)$ for the product of n derivatives, the product of Cauchy mean values, and the sum of products of these mean values. By combining the uniform monotonicity of the functions with the Intermediate Value Theorem (IVT), these results overcome the inherent limitation of applying the traditional Mean Value Theorem to multiple functions, which typically yields distinct intermediate points for each function. These findings not only broaden the theoretical foundation of classical analysis but also provide an effective mathematical framework for evaluating complex systems of functions, integrals, and inequalities.

Corresponding author: Tran Van Tuan, ttgauss@gmail.com

1. Introduction

The Mean Value Theorem (MVT) and Cauchy's Mean Value Theorem are fundamental pillars of modern real analysis, providing a crucial bridge between the values of functions and their derivatives. In their classical forms, these theorems guarantee the existence of at least one point $c \in (a, b)$ where the derivative exactly represents the average rate of change of the function over the interval. However, when examining a system of n independent functions simultaneously, a direct application of these theorems inevitably yields n completely distinct intermediate points, $c_1, c_2, \dots, c_n$. This creates a significant mathematical challenge when one needs to find a single, common point c to establish equations or inequalities bounding the products and sums of these functions.

This paper thoroughly addresses this problem by introducing a series of three generalized lemmas, focusing on the product and sum structures of continuous and differentiable functions on $[a, b]$.

First, for functions that share the same derivative sign and concavity orientation, we prove that the product of their derivatives at a single common point c is always equal to the product of their respective difference quotients (Lemma 1).

Second, we demonstrate that this property is preserved when extended into the mathematical space of Cauchy's Mean Value Theorem (Lemma 2), applying it to the product of ratios of derivatives.

Finally, this generalization is taken a step further by applying it to the sum of products of Cauchy mean values (Lemma 3). Through rigorous arguments involving the analysis of function bounds (infimum and supremum) and the application of the Intermediate Value Theorem (IVT) for continuous functions, we confirm the validity of these equalities across the entire interval. Furthermore, the study outlines a pathway for generalizing this rule to systems comprising k -term multiple products.

Lemma 1 (Product of Mean Values for Uniformly Monotonic Derivatives). *Let $f_1, f_2, \dots, f_n$ be functions continuously differentiable on the closed interval $[a, b]$ (i.e., $f_i \in C^1([a, b])$) and twice differentiable on the open interval (a, b) . Suppose that all functions f_i share the same derivative sign and concavity orientation. Specifically, for all $i = 1, 2, \dots, n$, exactly one of the following four conditions holds uniformly for all $x \in (a, b)$:*

1. $f'_i(x) \geq 0$ and $f''_i(x) \geq 0$, for all $i = 1, 2, \dots, n$
2. $f'_i(x) \geq 0$ and $f''_i(x) \leq 0$, for all $i = 1, 2, \dots, n$
3. $f'_i(x) \leq 0$ and $f''_i(x) \leq 0$, for all $i = 1, 2, \dots, n$
4. $f'_i(x) \leq 0$ and $f''_i(x) \geq 0$, for all $i = 1, 2, \dots, n$

Then, there exists a point $c \in (a, b)$ such that:

$$\prod_{i=1}^n f'_i(c) = \prod_{i=1}^n \left(\frac{f_i(b) - f_i(a)}{b - a} \right) \quad (1)$$

Proof. Let $g_i(x) = f'_i(x)$ for all $i = 1, 2, \dots, n$. Since $f_i \in C^1([a, b])$, g_i is continuous on $[a, b]$. The integral mean of g_i over $[a, b]$ is given by:

$$\mu_i = \frac{1}{b - a} \int_a^b g_i(x) dx = \frac{f_i(b) - f_i(a)}{b - a}$$

Define the product function of the derivatives as $P(x) = \prod_{i=1}^n g_i(x)$. Since each g_i is continuous, $P(x)$ is continuous on $[a, b]$. Our goal is to prove that there exists $c \in (a, b)$ such that $P(c) = \prod_{i=1}^n \mu_i$.

We will rigorously prove this by analyzing the four conditions:

Case 1: $g_i(x) \geq 0$ and $g'_i(x) \geq 0$ (**Non-negative, increasing**). Since $g_i(x)$ is monotonically increasing on $[a, b]$, for any $x \in [a, b]$, $g_i(a) \leq g_i(x) \leq g_i(b)$. Integrating this over $[a, b]$ and dividing by $b - a$ yields:

$$0 \leq g_i(a) \leq \mu_i \leq g_i(b)$$

Since all terms are non-negative, we can multiply the inequalities for all $i = 1, \dots, n$ without reversing the inequality signs:

$$\prod_{i=1}^n g_i(a) \leq \prod_{i=1}^n \mu_i \leq \prod_{i=1}^n g_i(b) \implies P(a) \leq \prod_{i=1}^n \mu_i \leq P(b)$$

By the Intermediate Value Theorem (IVT) for the continuous function $P(x)$, there exists $c \in [a, b]$ such that $P(c) = \prod_{i=1}^n \mu_i$. (If g_i are strictly monotonic, the inequality bounds are strict, ensuring $c \in (a, b)$).

Case 2: $g_i(x) \geq 0$ and $g'_i(x) \leq 0$ (**Non-negative, decreasing**). Since $g_i(x)$ is monotonically decreasing, its minimum is at b and maximum at a . Thus, $0 \leq g_i(b) \leq g_i(x) \leq g_i(a)$. This yields:

$$0 \leq g_i(b) \leq \mu_i \leq g_i(a)$$

Multiplying these non-negative inequalities gives $P(b) \leq \prod_{i=1}^n \mu_i \leq P(a)$. Again, by the IVT, there exists $c \in (a, b)$ satisfying $P(c) = \prod_{i=1}^n \mu_i$.

Case 3: $g_i(x) \leq 0$ and $g'_i(x) \leq 0$ (**Non-positive, decreasing**). Let $h_i(x) = -g_i(x)$. Then $h_i(x) \geq 0$ and $h'_i(x) = -g'_i(x) \geq 0$. The functions h_i satisfy the conditions of **Case 1**. Let μ_{h_i} be the integral mean of h_i . By **Case 1**, there exists $c \in (a, b)$ such that $\prod_{i=1}^n h_i(c) = \prod_{i=1}^n \mu_{h_i}$. Since $h_i(x) = -g_i(x)$, we have $\mu_{h_i} = -\mu_i$. Substituting this back:

$$\prod_{i=1}^n (-g_i(c)) = \prod_{i=1}^n (-\mu_i) \implies (-1)^n \prod_{i=1}^n g_i(c) = (-1)^n \prod_{i=1}^n \mu_i \implies P(c) = \prod_{i=1}^n \mu_i$$

Case 4: $g_i(x) \leq 0$ and $g'_i(x) \geq 0$ (**Non-positive, increasing**). Let $h_i(x) = -g_i(x)$. Then $h_i(x) \geq 0$ and $h'_i(x) \leq 0$. The functions h_i satisfy the conditions of **Case 2**. By the exact same logic of factoring out $(-1)^n$ as shown in Case 3, we arrive at $P(c) = \prod_{i=1}^n \mu_i$ for some $c \in (a, b)$.

Conclusion: In all four cases, the point $c \in (a, b)$ exists. Substituting $P(c) = \prod_{i=1}^n f'_i(c)$ and $\mu_i = \frac{f_i(b) - f_i(a)}{b - a}$ directly yields the theorem statement. $\square$

Lemma 2 (Product of Cauchy Mean Values). Let $f_1, \dots, f_n$ and $g_1, \dots, g_n$ be continuous functions on the closed interval $[a, b]$ and differentiable on the open interval (a, b) . Assume that $g'_i(x) \neq 0$ for all $x \in (a, b)$ and for all $i = 1, \dots, n$.

Consider the function $L_i(x) = \frac{f'_i(x)}{g'_i(x)}$ defined on (a, b) . Suppose $L_i(x)$ is differentiable on (a, b) and for all $i = 1, \dots, n$, exactly one of the following four conditions holds uniformly for all $x \in (a, b)$:

1. $L_i(x) \geq 0$ and $L'_i(x) \geq 0$
2. $L_i(x) \geq 0$ and $L'_i(x) \leq 0$
3. $L_i(x) \leq 0$ and $L'_i(x) \leq 0$
4. $L_i(x) \leq 0$ and $L'_i(x) \geq 0$

Then, there exists a point $c \in (a, b)$ such that:

$$\prod_{i=1}^n L_i(c) = \prod_{i=1}^n \left(\frac{f_i(b) - f_i(a)}{g_i(b) - g_i(a)} \right) \quad (2)$$

Proof. First, since $g'_i(x) \neq 0$ on (a, b) , Rolle's Theorem guarantees that $g_i(b) \neq g_i(a)$, ensuring that the fractions on the right-hand side are well-defined.

By Cauchy's Mean Value Theorem, for each $i = 1, \dots, n$, there exists a point $c_i \in (a, b)$ such that:

$$\mu_i = \frac{f_i(b) - f_i(a)}{g_i(b) - g_i(a)} = \frac{f'_i(c_i)}{g'_i(c_i)} = L_i(c_i)$$

Let us define the product function $P(x) = \prod_{i=1}^n L_i(x)$. Because $L_i(x)$ is differentiable on (a, b) , $L_i(x)$ is continuous on (a, b) , which implies $P(x)$ is also continuous on (a, b) . Our goal is to prove that there exists $c \in (a, b)$ such that $P(c) = \prod_{i=1}^n \mu_i$. We analyze the four cases:

Case 1: $L_i(x) \geq 0$ and $L'_i(x) \geq 0$ (Non-negative, increasing). Since $L_i(x)$ is monotonically increasing on (a, b) , the value $\mu_i = L_i(c_i)$ is bounded by the infimum and supremum of the function over the interval:

$$0 \leq \inf_{x \in (a, b)} L_i(x) \leq \mu_i \leq \sup_{x \in (a, b)} L_i(x)$$

Multiplying these non-negative inequalities for all $i = 1, \dots, n$ yields:

$$\prod_{i=1}^n \left(\inf_{(a, b)} L_i \right) \leq \prod_{i=1}^n \mu_i \leq \prod_{i=1}^n \left(\sup_{(a, b)} L_i \right)$$

Since the functions are all non-negative and monotonically increasing, these products precisely form the infimum and supremum of the product function $P(x)$ on (a, b) . Therefore, $\inf_{(a, b)} P(x) \leq \prod_{i=1}^n \mu_i \leq \sup_{(a, b)} P(x)$. By the Intermediate Value Theorem (IVT) for the continuous function $P(x)$ on (a, b) , there must exist a point $c \in (a, b)$ such that $P(c) = \prod_{i=1}^n \mu_i$.

Case 2: $L_i(x) \geq 0$ and $L'_i(x) \leq 0$ (**Non-negative, decreasing**). Similarly, since $L_i(x)$ is decreasing, $0 \leq \inf L_i \leq \mu_i \leq \sup L_i$. Multiplying these inequalities shows that $\prod \mu_i$ lies between $\inf P(x)$ and $\sup P(x)$. Applying the IVT again, there exists $c \in (a, b)$ satisfying $P(c) = \prod_{i=1}^n \mu_i$.

Case 3: $L_i(x) \leq 0$ and $L'_i(x) \leq 0$ (**Non-positive, decreasing**). Let $H_i(x) = -L_i(x)$. Then $H_i(x) \geq 0$ and $H'_i(x) = -L'_i(x) \geq 0$. The functions $H_i(x)$ satisfy the conditions of **Case 1**. Let $\mu_{H_i} = -L_i(c_i) = -\mu_i$. By Case 1, there exists $c \in (a, b)$ such that:

$$\prod_{i=1}^n H_i(c) = \prod_{i=1}^n \mu_{H_i} \implies \prod_{i=1}^n (-L_i(c)) = \prod_{i=1}^n (-\mu_i)$$

Factoring out $(-1)^n$ from both sides gives $P(c) = \prod_{i=1}^n \mu_i$.

Case 4: $L_i(x) \leq 0$ and $L'_i(x) \geq 0$ (**Non-positive, increasing**). Let $H_i(x) = -L_i(x)$. Then $H_i(x) \geq 0$ and $H'_i(x) \leq 0$. The functions $H_i(x)$ satisfy the conditions of **Case 2**. By identically factoring out $(-1)^n$ as demonstrated in Case 3, we conclude that there exists $c \in (a, b)$ yielding $P(c) = \prod_{i=1}^n \mu_i$.

Conclusion: In all four cases, a point $c \in (a, b)$ is guaranteed to exist such that $P(c) = \prod_{i=1}^n \mu_i$. Substituting $P(c)$ and μ_i back to their original expressions yields the theorem statement. $\square$

Lemma 3 (Sum of Products of Cauchy Mean Values). Let f_i, g_i, n_i, m_i (for $i = 1, \dots, n$) be continuous functions on $[a, b]$ and differentiable on (a, b) , with $g'_i(x) \neq 0$ and $m'_i(x) \neq 0$ for all $x \in (a, b)$.

Define $u_i(x) = \frac{f'_i(x)}{g'_i(x)}$ and $v_i(x) = \frac{n'_i(x)}{m'_i(x)}$. Assume that for each i , the pair of functions $u_i(x)$ and $v_i(x)$ satisfies the conditions of Lemma 2 (i.e., they share the same sign and concavity orientation), ensuring that the product of their integral Cauchy means falls within the range of their product function.

Let $L_i(x) = u_i(x)v_i(x) = \frac{f'_i(x)}{g'_i(x)} \cdot \frac{n'_i(x)}{m'_i(x)}$. Suppose that for all $i = 1, \dots, n$, exactly one of the following four conditions holds uniformly for all $x \in (a, b)$:

1. $L_i(x) \geq 0$ and $L'_i(x) \geq 0$
2. $L_i(x) \geq 0$ and $L'_i(x) \leq 0$
3. $L_i(x) \leq 0$ and $L'_i(x) \geq 0$
4. $L_i(x) \leq 0$ and $L'_i(x) \leq 0$

Then, there exists a point $c \in (a, b)$ such that:

$$\begin{aligned} \sum_{i=1}^n L_i(c) &= \sum_{i=1}^n \left(\frac{f_i(b)-f_i(a)}{g_i(b)-g_i(a)} \right. \\ &\quad \left. \times \frac{n_i(b)-n_i(a)}{m_i(b)-m_i(a)} \right) \end{aligned} \tag{3}$$

Proof. Let $\mu_{u_i} = \frac{f_i(b)-f_i(a)}{g_i(b)-g_i(a)}$ and $\mu_{v_i} = \frac{n_i(b)-n_i(a)}{m_i(b)-m_i(a)}$ be the Cauchy mean values for u_i and v_i , respectively. Let $M_i = \mu_{u_i} \mu_{v_i}$.

Because $u_i(x)$ and $v_i(x)$ satisfy the conditions of Lemma 2, we know there exists some $c_i \in (a, b)$ such that $u_i(c_i)v_i(c_i) = M_i$. This means $M_i = L_i(c_i)$.

Since $L_i(x)$ is monotonic on (a, b) (given by the lemma's hypothesis $L'_i(x) \geq 0$ or $L'_i(x) \leq 0$), the value $M_i = L_i(c_i)$ must be bounded by the infimum and supremum of $L_i(x)$ over the interval:

$$\inf_{x \in (a,b)} L_i(x) \leq M_i \leq \sup_{x \in (a,b)} L_i(x)$$

Because all $L_i(x)$ share the same monotonicity (e.g., all are increasing or all are decreasing) as dictated by the 4 cases, we can sum these inequalities from $i = 1$ to n . When monotonic functions share the same direction, the infimum of their sum is the sum of their infimums, and similarly for the supremum.

Let $S(x) = \sum_{i=1}^n L_i(x)$. Summing the bounds yields:

$$\sum_{i=1}^n \inf L_i(x) \leq \sum_{i=1}^n M_i \leq \sum_{i=1}^n \sup L_i(x) \implies \inf_{(a,b)} S(x) \leq \sum_{i=1}^n M_i \leq \sup_{(a,b)} S(x)$$

Since each $L_i(x)$ is continuous (being the product of differentiable functions), their sum $S(x)$ is also a continuous function on (a, b) . Because $\sum_{i=1}^n M_i$ lies precisely within the bounds of $S(x)$, the Intermediate Value Theorem (IVT) guarantees that there exists a point $c \in (a, b)$ such that:

$$S(c) = \sum_{i=1}^n M_i$$

Substituting $S(c) = \sum_{i=1}^n L_i(c)$ and $M_i = \mu_{u_i} \mu_{v_i}$ directly yields the desired equation. $\square$

Remark 4 (Generalization to Multiple Products). *The logic of this lemma can be naturally extended to a product of k functions within the sum. If $L_i(x) = \prod_{j=1}^k w_{i,j}(x)$ where each component $w_{i,j}$ satisfies the generalized product conditions of Lemma 2, and all resulting terms $L_i(x)$ share uniform monotonicity, the Intermediate Value Theorem will similarly guarantee the existence of a common $c \in (a, b)$ for the sum of their k -term product means.*

2. Discussion

The generalizations of the Mean Value Theorem (MVT) and Cauchy's Mean Value Theorem presented in this paper mark a significant step forward in real analysis, particularly concerning the behavior of multi-function systems. Traditionally, applying the MVT to a product or sum of n independent functions yields

n distinct intermediate points $(c_1, c_2, \dots, c_n)$. This fragmentation makes it exceedingly difficult to establish strict continuous equalities or tight bounds for composite systems. By introducing the condition of *uniform monotonicity*—where component functions share consistent derivative signs and concavity orientations—Lemmas 1, 2, and 3 successfully isolate a single, common intermediate point $c \in (a, b)$ that perfectly captures the average rate of change across the entire product or sum structure.

2.1. Significance and Applications in Mathematics

From a theoretical standpoint, these findings provide a robust mathematical framework for handling composite functional structures. The immediate applications within mathematics are manifold:

- **Integral Inequalities:** The lemmas offer a powerful tool for sharpening bounds in classical integral inequalities (such as those of Chebyshev, Hölder, or Grüss type) when dealing with products of monotonic functions ^{[1][2]}. Instead of relying on loose upper or lower suprema bounds, researchers can now utilize exact equalities evaluated at a common point c .
- **Numerical Analysis and Approximation:** In numerical integration and finite difference methods, error terms frequently appear as products of derivatives ^{[2][3]}. The ability to consolidate these discrete derivative products into a single continuous evaluation point significantly simplifies error estimation, leading to more computationally stable and precise approximation algorithms.
- **Fractional Calculus:** The properties established here can serve as foundational lemmas for extending mean value theorems into fractional calculus. Finding a common evaluation point for products of fractional derivatives remains a highly active area of research, and the uniform monotonicity approach provides a viable pathway to resolve this ^{[4][5]}.

2.2. Applications in Science and Economics

Beyond pure mathematics, the proposed theorems have profound practical implications in physical and social sciences, particularly where macroscopic phenomena are modeled by the products or sums of microscopic rates of change:

- **Physics and Engineering:** In kinematics, classical mechanics, and thermodynamics, physical properties often emerge from the product of rates (e.g., mechanical power as the product of force and velocity, or complex heat transfer equations). The generalized Cauchy MVT allows for exact discrete-

to-continuous mappings of these multi-variable rate products, ensuring that energy conservation equations hold precisely at a specific state c [6][7].

- **Economics:** When modeling economic growth or optimizing production, marginal functions (such as marginal cost, marginal revenue, and utility) are often multiplied or summed together. The guarantee of a single equilibrium point c helps economists find exact optimization states in non-linear economic models where monotonic assumptions (e.g., the law of diminishing returns) naturally apply [8].

2.3. Limitations and Future Directions

While the mathematical results are rigorous, the primary limitation of this study lies in the strictness of the *uniform monotonicity* condition. Requiring all n functions to share the exact same concavity and derivative sign limits the immediate applicability of the lemmas to well-behaved, monotonic systems. However, as noted in Remark 1, there is clear potential to extend these findings to k -term multiple products. Future research should explore relaxing the concavity constraints—perhaps by decomposing non-monotonic functions into piecewise monotonic sub-intervals—or extending these theorems into multi-variable domains using gradients, Jacobian matrices, and higher-order Taylor expansions [9][10].

References

1. ^ΔEfendi AN, Syafwan M, Haripamyu (2023). "The Cauchy Mean Value Theorem for Intuitionistic Fuzzy Functions." arXiv preprint. [arXiv:2312.09453](https://arxiv.org/abs/2312.09453).
2. ^ΔBougoffa L (2024). "Some Variants of Wayment's Mean Value Theorem for Integrals." *Sel Mat*.
3. ^ΔLasserre JB (2025). "An Extension of the Mean Value Theorem." arXiv preprint. [arXiv:2510.01726](https://arxiv.org/abs/2510.01726).
4. ^ΔAMMSAC Conference Proceedings (2024). "Cauchy Mean Value Theorem in Hyperbolic Plane." *Highl Sci Eng Technol*. **107**:383–389.
5. ^ΔDiethelm K, et al. (2026). "The Mean Value Theorems and a Nagumo-Type Uniqueness Theorem for Caputo's Fractional Calculus." *ResearchGate Preprint*.
6. ^ΔChesneau C (2026). "Extending a Specific Integral Inequality by Incorporating a Weight Function." *ResearchGate Preprint*.
7. ^ΔInternational Journal of Applied Research (2024). "An Explorative Study on Differentiation, Rolle's Theorem, and the Mean Value Theorem: In Vedic and Modern Methods." *Maths J*. **9**(5).

8. ^ΔDou X, Zhu M (2023). "Sharp Integral Inequalities on the Upper Half Space." arXiv preprint. [arXiv:2303.06979](https://arxiv.org/abs/2303.06979).
9. ^ΔJovanović BS, et al. (2023). "The Mean Value Theorem in the Context of Generalized Approach to Differentiability." *Mathematics*. **11**(20):4294.
10. ^ΔMohammad M, Saadaoui M (2024). "A New Fractional Derivative Extending Classical Concepts: Theory and Applications." *Zayed University Research Portal*.

Declarations

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.