

Open Peer Review on Qeios

Stage III Ovarian Cancer AJCC v8

National Cancer Institute

Source

National Cancer Institute. <u>Stage III Ovarian Cancer AJCC v8</u>. NCI Thesaurus. Code C139971.

Stage III includes: IIIA1: T1/2, N1, M0; IIIA2: T3a, N0/N1, M0; IIIB: T3b, N0/N1, M0; IIIC: T3c, N0/N1, M0. T1: Ovarian cancer with tumor limited to ovaries (one or both). T2: Ovarian cancer with tumor involving one or both ovaries with pelvic extension below pelvic brim. T3a: Ovarian cancer with microscopic extrapelvic (above the pelvic brim) peritoneal involvement with or without positive retroperitoneal lymph nodes. T3b: Ovarian cancer with macroscopic peritoneal metastasis beyond pelvis 2 cm or less in greatest dimension with or without metastasis to the retroperitoneal lymph nodes. T3c: Ovarian cancer with macroscopic peritoneal metastasis beyond the pelvis more than 2 cm in greatest dimension with or without metastasis to the retroperitoneal lymph nodes (includes extension of tumor to capsule of liver and spleen without parenchymal involvement of either organ). N0: No regional lymph node metastasis. N1: Positive retroperitoneal lymph nodes only (histologically confirmed). M0: No distant metastasis. (AJCC 8th Ed.)

Qeios ID: GYBSNB · https://doi.org/10.32388/GYBSNB