

Classical Explanation of Absorption Spectra

G. H. Jadhav

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.

Abstract

This paper first discusses the difficulties in interpreting the absorption spectra of a gas of particular atoms filled in a chamber with perfectly reflective inner walls. It is expected that if an atom produces a light wave of frequency ν , the electron that generated this wave due to its transition must oscillate at one place in the atom with that frequency which must be the natural frequency of that electron at that place. But no atomic model is in a position to support this. When a light wave of the same frequency interacts with the same electron of the atom, it will oscillate at the same frequency, which is called resonance. But if there is a difference of 180 degrees between the light wave created by these oscillations of that electron and the light wave interacting that electron, no net wave can be created or come out, so a black line will appear at the frequency ν at absorption spectrum of gas of such atoms. No matter how long light is shone on that gas, this condition will remain forever. By considering the conservation of energy the same thing cannot be explained using the concept of absorption of photons, which seems to be ignored forever.

Ghanshyam Jadhav

Shri Chhatrapati Shivaji College, Oemrga-413606, India

Email: ghjadhav@rediffmail.com

1. Introduction

Absorption spectroscopy is primarily used as an analytical chemistry tool to determine the presence or quantity of a particular substance in a sample. Among them, Atomic Absorption Spectroscopy (AAS), a simple but relatively expensive technology, is widely used to identify elements in elemental solutions. Technique used in absorption spectroscopy is that when a radiation interacts with a sample, it absorbs some part of it which can be function of frequency or wavelength. The chemical information of that compound can be extracted from it. A substance that emits radiation also absorbs the same kind of radiation, so the emission spectrum gives as much information about the substance as the absorption spectrum. Kirchhoff first showed in 1859 that the Fraunhofer lines seen in the sun were actually atomic absorption lines and that they were produced by various elements in the sun's atmosphere. Further Kirchhoff and Bunsen showed that different elements in a flame can give characteristic emission and absorption spectra. This gave rise to emission spectroscopy. For

the next hundred years or so, much emphasis was placed on emission spectroscopy, perhaps Kirchhoff and Bunsen's measurements were used only for visible light. But later it was noticed that atomic absorption methods could offer many potential advantages over emission methods ^{[1][2]}. This initially posed a problem in automating the sample from which emission and absorption spectra were to be obtained, but over time research has evolved into different methods of automating the sample and improving the application of spectroscopy ^{[3][4][5]}. But one thing to note is that no matter how advanced the techniques in emission and absorption spectroscopy are, there is still no clarity as to what exactly is going on where photons are emitted or absorbed. It is unfortunate that when an atom emits a light wave, the electron that generated the light wave must oscillate to a place in that atom at the frequency of the light wave. That is, the natural frequency of oscillations of that electron should be at the place in the atom where the corresponding electron oscillates and creates the light wave. This means that when a light wave of the same frequency interacts with that electron, it will oscillate at the same frequency which is its natural frequency of oscillations which can be called as resonance. Therefore, an attempt has been made in this paper to find out what its overall effect can be.

A hypothetical experiment and its expected results are discussed in next section. Section 3 discusses in detail how and why the expected results of the relevant experiments are obtained through aspects of quantum mechanics and classical mechanics. Section 4 includes the expected conclusions.

2. Experimental arrangement and results

For this a gas chamber can first be constructed as shown in Fig. 1, with two windows facing each other. The surface inside the gas chamber will be perfectly reflective and is filled with a gas at a proper pressure to obtain its absorption spectra. White light will enter through window W1 and pass straight through the gas and exit through window W2. The emitted light can be analyzed using a spectrometer and a prism. Suppose, according to quantum mechanics, it absorbs photons of the frequency ν , then in the light coming out of the gas, these photons will be absent. So, in the dispersion spectra produced by the prism, called the absorption spectra of the gas, a black line will appear whose wavelength is $\lambda=c/\nu$. Now, no matter how long the light through window W1 is put on the gas, there will be no difference in the absorption spectra, i.e. the black line will remain.




Fig. 1. Experimental arrangement to obtain absorption of a gas.

3. Discussion

In the above experiment, no matter how long the white light through window W_1 is thrown on the gas, if the black line in its absorption spectra remains, it means that the gas atoms in that chamber are constantly absorbing photons of ν frequencies. Suppose there are a total of n atoms of gas in that gas chamber, then any of the following possibilities can occur.

- All the n atoms of the gas in the chamber will absorb n number of photons of frequency ν from the incoming white light and go into excited states after which no photon of the frequency f will be absorbed and the black line will disappear again from the absorption spectra.
- n atoms in the gas will absorb photons of frequency ν from the incoming white light, according to their local conditions, and go into excited state and re-emit photons of the same frequency and come to ground state and this process continues continuously. Even if the re-emitted photons are assumed to go in any direction, since the internal wall of the chamber is perfectly reflective, either the photons will be reabsorbed by the gas atoms and some will also exit through window W_1 and some out of window W_2 . But since the surface area of the windows is much less than the surface area of the wall, the number of photons exiting the window W_1 will not affect the results of the experiment very much. Now either because the re-emitted photons are absorbed by the gas atoms again, photons of frequency ν coming from window W_1 will not be absorbed or photons of frequency f coming from window W_1 will be absorbed as much as re-emitted photons of frequency ν will come out of window W_2 which means that in the absorption spectra that black line will disappear again.

But experience so far suggests that the black line in the absorption spectra will remain constant no matter how long white light is thrown on the gas. This means that gas atoms are constantly absorbing photons of frequency ν . This is against law of energy conservation, and there is no explanation anywhere, and hardly anyone seems to be thinking about why this is happening.

According to the expectations of the classical theory, light is an electromagnetic wave and if the frequency of a light wave

is f , it is a universal truth that the electron transaction in the atom that created this wave should oscillate with this frequency at a place in the atom. There is no other option. But no one seems to be thinking about this. Each one relies on the equation $E_1 - E_2 = h\nu$. But no one knows anything about how that electron oscillates at the frequency ν and it is a matter of surprise that there is no mention of it.

The cause of this situation is the prevailing atomic models. Two atomic models seem to be accepted till date, one is the Rutherford-Bohr atomic model and the other is the Cloud atomic model. In the Rutherford-Bohr atomic model, the electrons are assumed to be moving continuously in orbit so that their centripetal and centrifugal forces are balanced. Therefore, there is no question of an electron oscillating in a stationary position in an atom, so this model cannot throw any light on how the energy difference of the electron is converted into a light wave. Despite the many problems with this atomic model, it is surprising that it is featured in the logos of many International Atomic Energy Agencies or Institutes. The first thing is that if an electron moves in an orbit, it will radiate energy because an electron has its own electric field, so it will create electric field ripples while moving in an orbit, so its speed will slow down and eventually it will fall into a nucleus, which no one can stop. Yet Bohr introduced the concept of stationary orbit which was not to be accepted under any circumstances and it was to be decided that the electron could not move in an orbit under any circumstances [6]. Further, no one can explain how the net orbital magnetic moment of the paired electrons becomes zero, or how the centripetal force and the centrifugal force balance immediately after an electron jumps from one orbit to another. This means that electrons cannot move in orbits under any circumstances in atoms. So, another option should have been considered for how the electrons can stay away from nucleuse in atoms but it did not happen. Einstein later proposed light quanta to explain the photoelectric effect [7]. In fact, at the time it was known that light was an electromagnetic wave and how its electric and magnetic fields could apply forces on electrons in the photoelectric effect. Why no one thought of the force is an unsolved puzzle. Everyone stuck on the same thing that as the intensity of light increases, the energy in it should increase. So, the classical solution of photoelectric effect could not be found. If the force was considered at that time, it would have been realized that the asymmetric electric field in the light wave is responsible for the photoelectric effect. Not only this, but it is responsible for creating the magnetic field effect and the answer to why magnetic monopoles are not found in the universe would have been answered and physics today could have been seen following a different path. But still, no one seems to consider the force to solve the photoelectric effect. However, due to Einstein's proposal of light quanta, light came to be regarded as a particle rather than a wave. From this, de Broglie proposed that if a particle is in motion, it can be expressed as a wave. Using this, Schrödinger predicted where the electron might be in the hydrogen atom, assuming the electron to be a wave rather than a particle, which is called the cloud atomic model. This further complicates the subject of the atom, which is not in a position to predict how an electron will oscillate at that frequency while emitting a light wave. It is a universal truth that the corresponding electron in that atom must oscillate at the frequency at which the light wave emits. The first condition for this would be that all the electrons in the atom should not move in orbits. The only alternative left is that if the two electrons are in opposite spin motion about the same axis, their net spin magnetic moment becomes zero and magnetic repulsion is created between them [8]. If there is a positive core between them as a nucleus, the net attractive force between the nucleus and the electrons and the net electric and magnetic repulsion between the electrons and electrons can be balanced, and these two electrons can be fixed at a certain distance from the nucleus, which can be called the spin atomic model, which further must be refined. So that it will

be possible to get the information about how electrons settle in different shells and how much their total energy is and at what frequency it can oscillate at that place. But until then, this indicates that electrons are fixed in atoms due to their spin motion, which is why they can oscillate around their mean positions and produce light waves, i.e. electromagnetic waves. Another possibility arises from this that there should be a natural frequency of the electron at the place where it is fixed in the atom. How far it jumps to that point should not make a difference in the frequency, but the amplitude of the oscillations should make a difference. The natural frequency of oscillations of that electron at that point must depend on the forces that stabilize the electron at that point. From this one can make some predictions about what is happening in the absorption spectra.

Suppose an electron in an atom, where it is fixed, has the natural frequency of oscillations, and if an electromagnetic wave or light wave of the same frequency interacts with that electron, it will oscillate with the same frequency, it is called resonance. In fact, the magnetic field is an effect of the asymmetric electric field, so there is only the electric field in the light wave. Now when the light wave interacts with the electron, the displacement of the electron will be in the opposite direction to the electric field in the light wave. But the oscillations of electrons will also produce a light wave or an electric field wave. If the direction of the electric field of the applied light wave and the direction of the electric field created by the electron are opposite to each other, i.e. the two waves have a phase difference of 180 degrees, then the effective electric field will be zero. No wave will come out of that process and the process will be continuous. Therefore, a black line will always appear in the absorption spectra at that frequency i.e. at the corresponding wavelength. There can be no other option. Of course, this will raise many questions that need to be answered. But if you want to understand this universe, you have to come down to earth.

4. Conclusion

When atom emits a light wave, the concerned electron, because transition of which the light wave is created, must vibrate at a place in the atom with the frequency of the light wave. That electron should have natural frequency of oscillations at that position in the atom. Therefore, the electrons in the atoms are not moving in orbits. They can maintain their safe distance from the nucleus by adjusting their spin motions. If light wave of natural frequency of the electron at that place in the atom interacts with the electron, then it will start to oscillate with this frequency otherwise it will not. It is called resonance. If the light wave generated by the oscillations of the electron and the light wave interacting with the electron are 180 degrees out of phase then there will be no resultant wave produced.

Other References

- B. J. Russel, J. P. Shelton and A. Walsh, *Spectrochim. Acta* 8, 317(1957)
- J. E. Allan, *Analyst* 83, 433(1958)

References

1. ^ C. T. J. Alkemade and J.M.W. Milatz, *Appl. Sci. Res. B4*, 289(1955)
2. ^ A. Walsh, *Spectrochim. Acta* 7, 108(1955)
3. ^ J. E. Allan, *Spectrochim. Acta* 18, 605(1962)
4. ^ J. B. Willis, *Nature* 107, 715(1965)
5. ^ A. Walsh, *Appl. Spectrosc.* 27, 335(1973)
6. ^ N. Bohr, *On the Theory of the Decrease of Velocity of Moving Electrified Particles on passing through Matter*, *Philosophical Magazine and Journal of Science*, 25, 10-31(1913)
7. ^ G. H. Jadhav, *Photoelectric effect: Magnetic force should increase with increase in frequency of EM waves*, *AIP Conf. Proc.* 2451, 020071 (2022)
8. ^ Ghanshyam Jadhav, *Spin Atomic Model: Role of Electron Spin in forming Atoms and Molecules*, *J. Phys.: Conf. Ser.* 2603 (2023)