

Research Article

Emergent Electromagnetism from the Gauged Logarithmic Superfluid Vacuum

Boris Kulangiev¹

1. Independent researcher

The relativistic logarithmic Klein–Gordon equation for the vacuum condensate possesses a global $U(1)$ phase symmetry. We show that promoting this to a local gauge symmetry—the minimal coupling $\partial_\mu S \rightarrow \partial_\mu S - eA_\mu$ —produces Maxwell’s equations as the Euler–Lagrange equations for the gauge field A_μ , with the condensate current $J^\nu = (em_{eff}/\hbar_{eff}^2)\rho u^\nu$ as source, on the same acoustic metric that carries the framework’s emergent gravity. The photon that results is exactly massless and achromatic: the gapless branch of the condensate is linear to all orders, so $c_{EM} = c_{grav} = c$ identically rather than by numerical suppression, and the source-free gauge action is conformally invariant on the acoustic metric, so the photon couples only to the gravitational flow and not to the conformal factor. The central result concerns *where* the gauge field may couple. We prove that the coupling cannot act on the bulk condensate: a gauged bulk would acquire a Planck-scale photon mass through the Anderson–Higgs mechanism and a uniform vacuum charge density for which Gauss’s law has no solution, both excluded by many orders of magnitude, and the two obvious escapes—bulk neutrality and phase inhomogeneity—are shown to fail. The coupling is therefore confined to the topological defect sector, the bulk remaining neutral. This yields an asymmetry between the two emergent interactions that general relativity postulates rather than explains: gravity couples universally, to the energy of every soliton, while electromagnetism couples selectively, only to the charged defects. London screening and the Meissner effect are correspondingly recovered not as a pathology of the vacuum but as the electrodynamics of charged matter. A microscopic account of why the bulk phase decouples from A_μ is identified as the principal open problem.

Correspondence: papers@team.qeios.com — Qeios will forward to the authors

1. Introduction

The logarithmic nonlinearity was introduced in nonlinear quantum theory ^[1]; its relativistic Klein-Gordon form, on which this work is built, originates with Rosen ^[2] and was analyzed by Bartkowski and Górka ^[3]. Emergent spacetime metrics from a logarithmic condensate have also been obtained in the separate superfluid-vacuum-theory programme ^[4]; that programme and the present logarithmic-superfluid-vacuum framework share the logarithmic nonlinearity but differ in their field ontology, rest-frame structure, and relativistic formulation. The acoustic-metric formalism through which long-wavelength excitations of a condensate experience an emergent Lorentzian geometry was established in the analogue-gravity programme ^{[5][6][7]}.

In previous work, we showed that the Madelung decomposition of the relativistic logarithmic Klein-Gordon equation produces an acoustic metric equivalent to the Schwarzschild geometry in the weak-field limit, yielding the PPN parameter $\gamma = 1$ ^{[8][6][7]} and an emergent gravitational constant $G_{eff} \sim c^2 / (\xi^2 \rho_0)$ ^{[9][10]}, with ξ the condensate healing length identified with the Planck length and ρ_0 at the Planck density. The vacuum condensate's density ρ and flow velocity $u_\mu = \partial_\mu S / m_{eff}$ generate the gravitational field; topological defects (vortex-Gaussons) play the role of massive particles.

This framework has a conspicuous unused structure. The logarithmic wave equation

$$\square\psi + \mu^2\psi - b\psi\ln\left(\frac{|\psi|^2}{\rho_0}\right) = 0 \quad (1)$$

is invariant under the global phase rotation $\psi \rightarrow \psi e^{i\alpha}$ for constant α . This $U(1)$ symmetry is associated, via Noether's theorem, with the conservation of the condensate particle number (the continuity equation $\partial_\mu(\rho\partial^\mu S) = 0$).

In the Standard Model, the fundamental interactions arise from *gauging* internal symmetries: promoting global invariances to local ones and introducing gauge fields to maintain covariance. Electromagnetism itself originates from gauging the $U(1)$ phase symmetry of the electron field ^[11].

In this paper, we apply the same principle to the vacuum condensate and follow its consequences. Promoting the global $U(1)$ of Eq. (1) to a local symmetry introduces a gauge field A_μ through the minimal coupling $\partial_\mu \rightarrow D_\mu = \partial_\mu - i(e/\hbar)A_\mu$; the Euler-Lagrange equations for A_μ are Maxwell's equations, with the condensate current as source, propagating on the same acoustic metric as the gravitational sector. Two features then follow. First, the emergent photon is exactly massless and achromatic, and

$c_{EM} = c_{grav}$ is structural rather than tuned. Second—the central result—the gauge coupling cannot act on the bulk condensate: a charged bulk would acquire a Planck-scale photon mass and a uniform vacuum charge density, both excluded, so the coupling is confined to the topological defect sector and the bulk is necessarily neutral. This forced restriction is the condensate origin of an asymmetry that general relativity postulates: gravity couples to all energy, electromagnetism only to charge.

2. The Gauged Lagrangian

The ungauged Lagrangian density for the logarithmic condensate is:

$$\mathcal{L}_0 = (\partial_\mu \psi^*)(\partial^\mu \psi) - \mu^2 |\psi|^2 + b |\psi|^2 \ln \left(\frac{|\psi|^2}{\rho_c} \right), \quad (2)$$

where ρ_c is the characteristic density of the logarithmic potential. The background density ρ_0 is defined by the stationarity condition $V'(\rho_0) = 0$, which gives $\rho_c = e\rho_0$. This is invariant under $\psi \rightarrow \psi e^{i\alpha}$ for constant α .

Promoting to local gauge invariance $\psi \rightarrow \psi e^{i\alpha(x)}$ requires replacing $\partial_\mu \rightarrow D_\mu = \partial_\mu - i(e/\hbar)A_\mu$ and adding the gauge field kinetic term. The gauged Lagrangian is:

$$\boxed{\mathcal{L} = (D_\mu \psi)^*(D^\mu \psi) - \mu^2 |\psi|^2 + b |\psi|^2 \ln \left(\frac{|\psi|^2}{\rho_c} \right) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}} \quad (3)$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the field strength tensor and e is the gauge coupling constant.

The term $-\frac{1}{4} F_{\mu\nu} F^{\mu\nu}$ is the unique Lorentz-invariant, gauge-invariant, parity-preserving kinetic term for A_μ with at most two derivatives [\[11\]](#). Its inclusion is not optional: without it, A_μ would have no dynamics and could be gauged away.

Two remarks fix the geometric setting of this term. First, the gauge field is an excitation of the same condensate whose density and flow define the acoustic metric $g_{\mu\nu}(\rho, u)$ (Sec. 3); it therefore propagates on that metric, not on a separate flat background. The kinetic term is understood in its covariant form

$$-\frac{1}{4} \sqrt{-g} F_{\mu\nu} F_{\rho\sigma} g^{\mu\rho} g^{\nu\sigma}, \quad (4)$$

with $g_{\mu\nu} = g_{\mu\nu}(\rho, u)$ the acoustic metric; the flat expression is its local-inertial-frame form, valid at the background where $g_{\mu\nu} \rightarrow \eta_{\mu\nu}$. Second, the source-free Maxwell action (4) is conformally invariant on the acoustic metric: under a rescaling $g_{\mu\nu} \rightarrow \Omega^2 g_{\mu\nu}$ the volume factor $\sqrt{-g}$ and the pair of inverse metrics contribute canceling powers of Ω , leaving the action unchanged. (This cancellation is a property of the

acoustic metric's Lorentzian $3 + 1$ structure, in which the emergent time direction enters with opposite signature to the three spatial directions; it is not an assumption that time is a fundamental dimension.) The acoustic metric's departure from flat is carried by an overall conformal factor ρ/c_s together with the Painlevé–Gullstrand flow term; the conformal factor drops out of the photon dynamics entirely, so the gauge sector is sensitive only to the flow. This is the geometric reason the emergent photon remains exactly achromatic (Sec. 5) while still being deflected by the gravitational flow (Sec. 7).

A scope qualification is required at the outset. As written, Eq. (3) assigns charge e to the entire field ψ , including the topologically trivial bulk condensate. Section 6 shows that this assignment is untenable—a charged bulk generates a Planck-scale photon mass and a uniform vacuum charge density—and that consistency requires the covariant derivative to act only on the charged, topological sector: the vortex–Gausson defects. Equation (3) is therefore to be read as the Lagrangian of the charged sector, with the neutral bulk governed by $\mathcal{L}_0$. The kinematic results of Secs. 3–4 concern the structure of the gauged equations and are unaffected by this restriction.

3. Madelung Decomposition of the Gauged Equation

3.1. The gauge-invariant velocity

Applying the Madelung transformation $\psi = \sqrt{\rho}e^{iS/\hbar_{eff}}$, the covariant derivative becomes:

$$D_\mu \psi = \left(\frac{\partial_\mu \sqrt{\rho}}{\sqrt{\rho}} + \frac{i}{\hbar_{eff}} (\partial_\mu S - eA_\mu) \right) \psi. \quad (5)$$

The gauge-invariant superfluid four-velocity is:

$$u_\mu = \frac{1}{m_{eff}} (\partial_\mu S - eA_\mu) \quad (6)$$

This is the central modification. In the ungauged theory, $u_\mu = \partial_\mu S/m_{eff}$; the gauge field shifts the velocity by $-eA_\mu/m_{eff}$. This is the relativistic generalization of the London relation in superconductors: $m\mathbf{v}_s = \hbar\nabla\phi - e\mathbf{A}$ ^[12].

3.2. The modified fluid equations

Separating the Euler–Lagrange equation for ψ into real and imaginary parts yields the gauged Madelung equations:

Continuity:

$$\partial_\mu(\rho u^\mu) = 0. \quad (7)$$

Hamilton-Jacobi:

$$\frac{m_{eff}^2}{\hbar_{eff}^2} u_\mu u^\mu = \mu^2 - b \ln\left(\frac{\rho}{\rho_c}\right) + Q[\rho], \quad (8)$$

where $Q = \square\sqrt{\rho}/\sqrt{\rho}$ is the quantum potential (Bohm potential) [13], unchanged from the ungauged case.

These equations are *identical in form* to the ungauged Madelung equations, with the single replacement $\partial_\mu S \rightarrow \partial_\mu S - eA_\mu$. The acoustic metric depends on ρ and u_μ , both of which are gauge-invariant. The entire gravitational sector—the PPN parameter $\gamma = 1$, the emergent G_{eff} , the Painlevé–Gullstrand flow—is preserved without modification.

4. Maxwell's Equations from the Gauge Field Variation

The Euler-Lagrange equation for A_μ is obtained by varying the Lagrangian (3) with respect to A^ν .

The condensate contribution:

$$\frac{\delta}{\delta A^\nu} [(D_\mu \psi)^* (D^\mu \psi)] = -\frac{em_{eff}}{\hbar_{eff}^2} \rho u^\nu. \quad (9)$$

The gauge field contribution, from the covariant kinetic term (4):

$$\frac{\delta}{\delta A^\nu} \left[-\frac{1}{4} \sqrt{-g} F_{\alpha\beta} F_{\rho\sigma} g^{\alpha\rho} g^{\beta\sigma} \right] = \sqrt{-g} \nabla_\mu F^{\mu\nu}. \quad (10)$$

Setting the total variation to zero gives Maxwell's equation on the acoustic metric,

$$\boxed{\nabla_\mu F^{\mu\nu} = \frac{1}{\sqrt{-g}} \partial_\mu \left(\sqrt{-g} F^{\mu\nu} \right) = J^\nu} \quad (11)$$

which at the background metric $g_{\mu\nu} \rightarrow \eta_{\mu\nu}$ reduces to the flat form $\partial_\mu F^{\mu\nu} = J^\nu$, while near a gravitating mass the metric factors encode the deflection, redshift, and time delay of electromagnetic waves by the Painlevé–Gullstrand flow (Sec. 7). The four-current is:

$$J^\nu = \frac{em_{eff}}{\hbar_{eff}^2} \rho u^\nu. \quad (12)$$

Here ρ and u^ν are the density and gauge-invariant velocity of the charged sector (Sec. 2); for the neutral bulk this current vanishes identically. In the far field of a vortex-Gausson, this reduces to the current of a localized charged defect.

Equation (11) is Maxwell's inhomogeneous equation. The homogeneous equations $\partial_{[\mu} F_{\nu\lambda]} = 0$ follow identically from the definition $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ (the Bianchi identity).

In three-vector notation, at the background metric (the flat reduction; curvature enters through the metric factors of Eq. 11):

$$\nabla \cdot \mathbf{E} = \rho_{ch}/\epsilon_0 \quad (\text{Gauss}), \quad (13)$$

$$\nabla \cdot \mathbf{B} = 0 \quad (\text{no monopoles}), \quad (14)$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \quad (\text{Faraday}), \quad (15)$$

$$\nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t} \quad (\text{Ampère-Maxwell}), \quad (16)$$

where $\rho_{ch} = J^0$ is the charge density and $\mathbf{J} = (J^1, J^2, J^3)$ is the current density. These are the full Maxwell equations, derived from the gauge field variation of the condensate Lagrangian.

The defects that source this current carry a conserved topological index. A vortex-Gausson of winding number n has phase increasing by $2\pi n \hbar_{eff}$ around any loop encircling its core, $\oint \partial_\mu S dx^\mu = 2\pi n \hbar_{eff}$ with $n \in \mathbb{Z}$, by single-valuedness of the order parameter. The integer n labels a conserved sector: $n = 0$ is the topologically trivial bulk, and $n \neq 0$ are the localized defects. As shown in Sec. 6, only the latter can carry the gauge coupling; the current (12) is nonzero only there.

5. Electromagnetic Waves and the Achromatic Photon

In the vacuum background the neutral bulk does not couple to A_μ (Sec. 6); with no charged defects present, $J^\nu = 0$ and the Maxwell equations (11) reduce to:

$$\square A_\mu = 0 \quad (\text{in Lorenz gauge}), \quad (17)$$

where $\square = \partial_t^2/c^2 - \nabla^2$ and $c = c_s(\rho_0)$ is the sound speed at the background density. Electromagnetic waves propagate at the speed of sound in the condensate.

This provides a unified explanation: the speed of light is the speed of sound because the photon is a gauge excitation propagating on the same condensate whose acoustic properties define c . The identity $c_{EM} = c_{grav} = c_{sound}$ is not a coincidence but a structural consequence of both fields emerging from the same substrate.

The photon is moreover *exactly* achromatic. The gapless excitation branch of the relativistic logarithmic condensate is linear, $\omega = ck$ to all orders [8]: the second-order relativistic kinetic operator admits no k^4 correction, in contrast to the non-relativistic Bogoliubov spectrum. Electromagnetic propagation in the bulk therefore carries no energy-dependent dispersion at any accessible scale, satisfying the

gravitational-wave and Lorentz-invariance bounds identically rather than by numerical suppression. Any detection of an energy-dependent photon velocity at sub-Planckian energies would falsify this structure.

Near a gravitating mass, the acoustic metric modifies the effective spacetime seen by all excitations equally. Electromagnetic waves are deflected, redshifted, and time-delayed by the Painlevé–Gullstrand flow, reproducing the standard general relativistic predictions. The Shapiro time delay, the gravitational redshift, and the bending of light are automatically consistent with $\gamma = 1$ because the photon (gauge excitation) and the phonon (density excitation) share the same acoustic metric. The conformal factor ρ/c_s of the metric drops out of the source-free gauge action (Sec. 2), so the photon responds only to the flow, not to the conformal part—the geometric statement behind its achromaticity.

6. The Gauge Coupling Cannot Act on the Bulk

We now establish the central structural result: a literal reading of Eq. (3), in which the bulk condensate itself carries the gauge coupling, is excluded.

In a region of uniform density ($\rho = \rho_0$) and smooth phase, the gauge-invariant velocity (6) contains the piece

$$u_\mu = -\frac{e}{m_{eff}} A_\mu, \quad (18)$$

which, substituted into the current (12), yields a response proportional to the gauge field itself (m_{eff} cancels):

$$J^\nu = -\frac{e^2 \rho_0}{\hbar_{eff}^2} A^\nu. \quad (19)$$

Combined with Maxwell's equation (11), this gives the Proca equation

$$\square A^\nu + \frac{1}{\lambda_L^2} A^\nu = 0, \quad (20)$$

with London penetration depth

$$\lambda_L = \frac{\hbar_{eff}}{e} \frac{1}{\sqrt{\rho_0}}. \quad (21)$$

This is the Anderson–Higgs mechanism: the gauge field absorbs the condensate phase and propagates with mass $m_\gamma = \hbar_{eff}/(\lambda_L c)$. With ρ_0 at the Planck density, λ_L is microscopic and m_γ is of order the Planck mass up to the dimensionless coupling—at least forty orders of magnitude above the

observational bound $m_\gamma < 10^{-18} eV$. A minimally gauged bulk would be a Planck-scale superconductor, opaque to electromagnetic radiation.

Independently, the same conclusion follows from the charge density. The background carrier phase $\partial_t S = -m_{eff}c^2$, structural throughout the framework [8], makes $u^0 \neq 0$: the gauged bulk would carry a uniform charge density $J^0 \propto e\rho_0$, for which Gauss's law admits no consistent solution in an infinite homogeneous medium. The photon-mass obstruction and the charge-density obstruction are independent, and either alone is fatal.

Two seemingly natural escapes fail. Electrical *neutrality* of the bulk does not help: the screening kernel in Eq. (19) is proportional to $e^2\rho_0$, not to the net charge, and laboratory superconductors—globally neutral systems—expel magnetic fields for precisely this reason. Nor does spatial variation of the macroscopic phase: far from gravitating masses $\nabla S \rightarrow 0$ and $\rho \rightarrow \rho_0$, so the screening would be *strongest* exactly where electromagnetic waves are observed to propagate freely. (Note also that $\lambda_L \propto 1/\sqrt{\rho}$: screening weakens, not strengthens, where the density drops toward a vortex core.)

The resolution is forced. The bulk condensate is topologically trivial, $n = 0$, and must lie outside the gauge-coupled sector; the covariant derivative acts only on the charged topological defects, while the neutral bulk is governed by the ungauged $\mathcal{L}_0$. With the bulk decoupled, the photon mass term vanishes identically, electromagnetic waves propagate freely at $\omega = ck$ (Sec. 5), and the Coulomb field of a charged particle is sourced by its charge through Gauss's law in the standard way. This is the reading of Eq. (3) announced in Sec. 2, now seen to be not a choice but the only consistent option.

The London response then reappears where it belongs. Wherever charged defects condense into a coherent macroscopic state—a laboratory superconductor—the charged-sector density ρ_{ch} is nonzero over a macroscopic region, and Eqs. (18)–(21) apply with $\rho_0 \rightarrow \rho_{ch}$: a finite penetration depth and the Meissner effect. The screening that consistency forbids in the vacuum is recovered as the correct electrodynamics of superconducting matter: the vacuum is a transparent dielectric; a charged condensate of defects is a superconductor.

7. The Coupling Asymmetry of the Two Interactions

The gauged logarithmic condensate produces two emergent interactions from a single Lagrangian, coupled through the condensate but not symmetric in character.

Gravity is the emergent geometry itself: the acoustic metric $g_{\mu\nu}(\rho, u)$ on which all condensate excitations propagate. The soliton energy density (mass) sets the scale of the metric perturbation, and self-consistency of the metric selects the Painlevé-Gullstrand form equivalent to Schwarzschild, with coupling $G_{eff} \sim c^2/(\xi^2 \rho_0)$, $\xi \sim \ell_P$, ρ_0 at the Planck density ^{[9][10]}. The microscopic origin of the collective flow that realizes this metric remains an open problem ^[9].

Electromagnetism arises from the gauge field A_μ required by local $U(1)$ invariance. Its source is the charge of the topological defect sector. Its field equation is Maxwell's $\nabla_\mu F^{\mu\nu} = J^\nu$. Its coupling constant is e .

The two are coupled through the condensate: a charged vortex-Gausson creates both a gravitational (acoustic-metric) and an electromagnetic (gauge) field; within charged matter the electromagnetic field modifies the charged-sector flow through the London relation (18), while the neutral bulk does not couple; and the gravitational flow deflects electromagnetic waves, giving the bending of light.

The essential result of this paper is the difference in *how* they couple. The geometric coupling is universal: every soliton carries energy, so every soliton sources the acoustic metric, and gravity acts on all matter. The gauge coupling is selective: only defects with $n \neq 0$ carry charge, the trivial bulk is neutral, and the charge of ordinary matter cancels. This asymmetry is not imposed—it is forced by the obstruction of Sec. 6, which excludes any universal (bulk) electromagnetic coupling. It is the condensate origin of a fact that general relativity postulates without explanation: gravity couples to all energy, electromagnetism only to charge.

8. Discussion

8.1. What is derived and what is assumed

The results rest on four inputs:

- i. The logarithmic wave equation (1), which defines the condensate dynamics.
- ii. The requirement of local $U(1)$ gauge invariance, which forces the introduction of A_μ .
- iii. The gauge field kinetic term $-\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$, which is the unique allowed term.
- iv. The restriction of the gauge coupling to the topological sector, leaving the bulk neutral.

Of these, (i) is the foundation of the entire framework. Input (ii) is a symmetry principle—the same one used in the Standard Model to derive electromagnetism from the electron field's $U(1)$ symmetry ^[11]. Input (iii) is uniquely determined by Lorentz invariance, gauge invariance, and the requirement of at

most two derivatives. Input (iv) is not assumed but *derived*: Sec. 6 shows a bulk-gauged condensate is excluded by the photon mass and vacuum charge it would generate, and that the topological sector is the only place the coupling can consistently reside. What remains open is one level deeper—a microscopic account of *why* the bulk phase decouples from A_μ , rather than the demonstration *that* it must. That microscopic account is the principal open problem of this work.

The coupling constant e is *not* determined by the condensate parameters. Deriving its value (equivalently, the fine structure constant $\alpha = e^2/(4\pi\epsilon_0\hbar c) \approx 1/137$) would require computing the self-energy of a vortex-Gausson in the gauged condensate and matching it to the observed electron mass. This remains an open problem.

8.2. Comparison with the Standard Model

In the Standard Model, electromagnetism arises from gauging the $U(1)_Y$ hypercharge symmetry of the fermion and Higgs fields. The photon is the massless gauge boson associated with the unbroken $U(1)_{EM}$ after electroweak symmetry breaking, and the electron's charge is a coupling constant.

In the present framework, electromagnetism arises from gauging the $U(1)$ phase symmetry of the vacuum condensate. The photon is the massless gauge excitation propagating on the condensate, and the electron's charge is the same coupling constant e , carried by the topological defect sector rather than by the trivial bulk. The key structural difference is that here the condensate provides both the gravitational field (acoustic metric) and the electromagnetic field (gauge connection), whereas the Standard Model treats gravity as external—and the coupling asymmetry between them (Sec. 7) emerges from the condensate rather than being postulated.

8.3. Limitations

The framework produces only a $U(1)$ gauge field. The Standard Model requires $SU(3) \times SU(2) \times U(1)$. Extending to non-Abelian gauge groups would require a multi-component order parameter (e.g., a spinor or matrix-valued condensate), which is beyond the scope of the present single-component model. All observed charges are integer multiples of e ; quarks carry fractional charges but are confined, so all observable states carry integer charge. Deriving the charge spectrum, and quark confinement, from the defect structure would require that non-Abelian extension and is left open.

9. Conclusion

Gauging the $U(1)$ phase symmetry of the logarithmic superfluid vacuum produces Maxwell's equations alongside the acoustic metric equations from a single Lagrangian, with an exactly massless, achromatic photon whose speed equals the gravitational sound speed as a structural identity. The central result is that the gauge coupling cannot act on the bulk condensate—a bulk-gauged vacuum is excluded by an Anderson-Higgs photon mass and an unsolvable vacuum charge density, with the obvious escapes closed—so the coupling is confined to the topological defect sector and the bulk is necessarily neutral. This yields, from the condensate, the asymmetry that general relativity postulates: gravity couples universally to energy, electromagnetism selectively to charge. London screening and the Meissner effect are recovered as the electrodynamics of charged matter rather than a property of the vacuum.

The coupling constant e (the fine structure constant) is not determined by the condensate parameters, and a microscopic account of why the bulk phase decouples from A_μ remains open; both are identified as the principal directions for further work. Extension to non-Abelian gauge groups requires a multi-component condensate and is deferred.

Acknowledgments

The methodological stance of this paper—treating the order parameter as a real classical field whose hydrodynamic behavior underlies emergent phenomena—is indebted to the Madelung-de Broglie-Bohm tradition, particularly as developed in P. R. Holland's exposition of the quantum theory of motion. Claude (Anthropic) was used as an editorial and computational tool during manuscript preparation. All scientific content, equations, and interpretations are the author's own.

References

1. ^ΔBiałynicki-Birula I, Mycielski J (1979). "Gaussons: Solitons of the Logarithmic Schrödinger Equation." *Phys Scr.* 20:539.
2. ^ΔRosen G (1969). "Dilatation Covariance and Exact Solutions in Local Relativistic Field Theories." *Phys Rev.* 183:1186.
3. ^ΔBartkowski K, Górka P (2008). "One-Dimensional Klein-Gordon Equation with Logarithmic Nonlinearities." *J Phys A: Math Theor.* 41:355201.

4. ^AZloshchastiev KG (2023). "Derivation of Emergent Spacetime Metric, Gravitational Potential and Speed of Light in Superfluid Vacuum Theory." *Universe*. 9:234.
5. ^AVolovik GE (2003). *The Universe in a Helium Droplet*. Oxford: Oxford University Press.
6. ^A^BVisser M (1998). "Acoustic Black Holes: Horizons, Ergospheres and Hawking Radiation." *Class Quant Gra* v. 15:1767.
7. ^A^BUnruh WG (1981). "Experimental Black-Hole Evaporation?" *Phys Rev Lett*. 46:1351.
8. ^A^B^CKulangiev BB (2026). "Conditions for Emergent Gravitational Light Bending from a Logarithmic Superfluid Vacuum." Preprint. <https://kulangiev.github.io/#paper1>.
9. ^A^B^CKulangiev BB (2026). "Emergent Newtonian Gravity from the Collective Acoustic Metric of a Logarithmic Superfluid Vacuum." Preprint. <https://kulangiev.github.io/#paper2>.
10. ^A^B^CKulangiev BB (2026). "Singularity Regularization and the Emergent Planck Scale in the Logarithmic Superfluid Vacuum." Preprint. <https://kulangiev.github.io/#paper5>.
11. ^A^B^CWeinberg S (1996). *The Quantum Theory of Fields, Vol. 2*. Cambridge: Cambridge University Press.
12. ^ALondon F, London H (1935). "The Electromagnetic Equations of the Supraconductor." *Proc Roy Soc A*. 149:71.
13. ^AHolland PR (1993). *The Quantum Theory of Motion*. Cambridge: Cambridge University Press.

Declarations

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.