

Review of: "Evaluation of Diabetes Risk Score Tool for Detecting Undiagnosed Type 2 Diabetes Mellitus in Attendees of Referral Clinics at Primary Health Care Centers, Sudan"

Mitaw Girma¹

1 Wollo University

Potential competing interests: No potential competing interests to declare.

Your update manuscript comment

- 3. What is the benefit of doing Hemoglobin A1C and RBS at the same time? See carefully this part
- 4. Exclusion Criteria: Already, patients who use metformin are known DM patients, but your intention is to study undiagnosed Type II DM patients; due to this reason, avoid patients who use metformin. B/c exclusion criteria are taken from inclusion criteria
- 5. **Sampling of the study participants:** Attendees at the RPHCCs were chosen using a systematic random sample (SRS), as you think, but you did not show how to calculate the Kth value and also how to calculate each facility's study participant Kth value. See carefully this part (major comment)
- 6. Inclusion Criteria: Why did you include in your study individuals previously diagnosed with diabetes? B/c your study focuses on tool evaluation for undiagnosing Type II DM (major comment) and avoids males and females, instead saying, Adult
- 8. **General comment:** Your sample size is not representative of the evaluation of tools, and your sampling also lacks reliability
- 9. By communicating with me and your justification being satisfied, your manuscript will be accepted

Qeios ID: IX3OK9 · https://doi.org/10.32388/IX3OK9