


A single hidden variable interpretation of the quantum wave function

David Gentile



Preprint v9

Oct 2, 2023

<https://doi.org/10.32388/J3PVM1.9>

Author: David Gentile

Location: Riverside, IL, USA

ORCID: 0000-0001-5777-761X

E-mail: gentile.david31@yahoo.com

Keywords: Measurement problem, Single hidden variable interpretation, Subjective and objective probability, Nonlocality, Decoherence, Delayed choice quantum eraser.

Statements and Declarations: No competing interests or financial support are declared.

A single hidden variable interpretation of the quantum wave function.

Abstract

An interpretation of quantum mechanics is presented in which there is one hidden variable. While this number of hidden variables is insufficient to explain Bell nonlocality, it can help solve the measurement problem. It also provides an explanation of the delayed choice quantum eraser experiment that is more intuitive than current explanations. In addition, a theory of nonlocality emerges directly from its assumptions. The key idea is that from the perspective of an outside observer, not correlated with an observed system, we treat all interactions within the system as unitary. However, from the perspective of an “observer” inside the system, we treat all new entanglements as “micro-observations” and as a projection onto some basis of measurement. One particle can “observe” another. Thus, particles will have a defined value on one measurement basis from an inside perspective. This means that the wave function, from the outside view, will at times describe an objective uncertainty, at other times a subjective uncertainty and at yet other times a combination of both. The standard wave function describes the total uncertainty, the minimal uncertainty that is present for any observer outside of the system, while for an observer inside the system, only the objective part of the uncertainty remains, and this can be described by an inner wave function. The hidden variable is invisible to the outside observer.

Contents

1. Introduction.
 2. Types of uncertainty.
 3. The basic idea and the delayed choice quantum erasure experiment.
 4. Nonlocality addressed.
 5. The measurement problem.
 - 5.1 What happens in a micromasurement?
 - 5.2 What happens in a macroscopic measurement?
 - 5.3 When do observations become nonerasable?
 - 5.4 Decoherence.
 - 5.5 The new Wigner’s friend thought experiment.
 6. Nonlocal effects and the measurement problem.
 - 6.1 Not erased, just rehidden.
 - 6.2 Destruction of entanglements.
 - 6.3 How should we define a measurement?
 7. Summary.
- References.
- Table of Figures.

1. Introduction.

We begin in section 2 by defining two types of uncertainty, objective and subjective. We then assert that the standard or outer wave function represents the total uncertainty to an outside observer

not correlated with the system, a combination of objective and subjective uncertainty. From this outside perspective, all interactions are treated as unitary interactions. From a perspective inside the system, all entanglements are treated as projections onto some basis. The particles “observe” each other and give each other definite values on some (unknown) measurement basis. From this inside perspective, only objective uncertainty remains, and this perspective can be described by an inner wave function. Thus, from the outside perspective, there is one hidden variable – a definite value on some measurement basis that is unknown. One hidden variable is insufficient to explain Bell nonlocality; however, it is helpful with other problems. We attempt to show, for example, that it offers a solution to the measurement problem. In section 3, after the most basic example, we first discuss a configuration in which, from the outside perspective, only subjective uncertainty remains. We then turn to a situation in which measurement by one observer, Alice, will resolve subjective uncertainty, but another, by Bob, will resolve objective uncertainty. In this configuration, we attempt to show that the results of the delayed choice quantum eraser experiment can be addressed and that this explanation is more intuitive than existing explanations. Then, in section 4, we turn to cases in which there is objective uncertainty for both experimenters. These examples will involve accounting for nonlocal correlations in order to explain their results. Here, a specific theory of how nonlocal correlations occur emerges directly from the assumptions of the interpretation and existing experimental results. Then, in sections 5 and 6, we attempt to tackle the measurement problem. In section 5, we address the questions of “what happens in a micromasurement?”, “what happens in a macroscopic measurement?”, and “when do observations become nonerasable?” Then, we look at the phenomenon of decoherence, which we argue can be reinterpreted, and we take a brief look at the new Wigner’s friend thought experiment. Finally, in section 6, we introduce nonlocal effects to elucidate the measurement problem. We argue that observations are not truly erased; they are just rehidden. Additionally, we address the question of when entanglements are broken, creating separate subsystems. We assert that “interactions” on the sterile path that particles do not follow are critical to this process. We conclude section 6 with a short discussion of how measurements should be defined, and then we summarize our work in section 7.

2. Types of uncertainty

Let us define two different types of uncertainty. One we will call intrinsic or objective uncertainty, and the other is epistemic or subjective uncertainty. The first represents real uncertainty in the universe, and the other is about what information we have available. In statistics, historically, these conceptions of probability divided mathematicians into two camps, classicists and Bayesians (Jaynes, 2003), although today, both are mostly accepted as two valid but different approaches to probability. One simple example of a coin flip is enough to illustrate the difference. Suppose someone flips a coin and holds the result behind their back. To a classical statistician, the coin is represented as a random number generator. The odds are 50/50 heads/tails before the flip. However, when the coin is flipped but still hidden, the probability is now 1 or 0. We just don’t know which. To the Bayesian, the odds are still 50/50 until we learn the result because Bayesian statistics is concerned with the information available to us.

While completely epistemic interpretations of quantum mechanics exist (Barzegar & Oriti, 2022), here, we assume that quantum mechanical systems can and do exhibit real, objective, intrinsic uncertainty. This can be illustrated by many experiments, but the most well known may be the classic 2-slit experiment (Ananthaswamy, 2018). The photons in this experiment clearly seem to pass through both slits in order to interfere with themselves. Both states exist in superposition with each other. This

kind of uncertainty is more than us just not knowing which path the photon took. We might say that the universe does not even “know”. Our macroscopic world does not exhibit this sort of behavior, nor does classical physics. The coin flip in classical physics would not be described as intrinsically random, but rather, we simply lack sufficient information to make predictions. Somehow in moving from the quantum world up to the macroscopic world, intrinsic uncertainty is lost.

Here, we assert that the standard wave function can represent objective uncertainty, subjective uncertainty, or a combination of both, depending on the circumstances, and that the transformation from one type of uncertainty to another is invisible from outside the system. From the perspective of an “outside” observer, not correlated with an observed system, we treat all interactions within the system as unitary. The standard, or outer, wave function then gives a perfect description of the system from this perspective, or at least gives us probabilities that accurately predict all experimental results, but what portion of it represents objective versus subjective uncertainty is unclear.

From the perspective of an observer “inside” the system¹, correlated with the system, we treat all new entanglements as “micro-observations”. One particle can “observe” another. This transforms some objective uncertainty into subjective uncertainty, at least temporarily², with every new entanglement. Assuming a new measurement is not completely compatible with the previous measurement, a “coin is flipped” when this happens. A nondeterministic event takes place. That is, from the perspective inside the system, projection onto some (unknown) basis has occurred, and a variable measured on that basis now has a definite value, while from an outside perspective, subjective uncertainty regarding that variable’s value remains. Objective uncertainty also persists, however. Values for other variables that would have to be measured on an incompatible orthogonal basis remain objectively undetermined. An inner wave function will describe this interior perspective.

The standard wave function describes the total uncertainty, objective and subjective combined. One might ask “Exactly who’s subjective uncertainty?” So, to be more precise, the outer wave function describes the minimal uncertainty that is present for ANY observer outside of the system. An individual observer could have greater uncertainty, for some idiosyncratic reason, unrelated to physical laws, but not less uncertainty³. We label the subjective part of the wave function “subjective” because the uncertainty is due to a lack of information. The information exists but is unavailable. Similarly, the objective uncertainty, described by the inner wave function, represents the minimal uncertainty that any observer must have, even if correlated with the system. We label it “objective” because the information needed to resolve this sort of uncertainty does not exist. Thus, from an inside perspective, every

¹ This outside/inside language is borrowed from (Ormrod, Vilasini, & Barrett, 2023) where it is used to describe macroscopic observers with inside and outside perspectives in the new Wigner’s friend thought experiment.

² We use the term “micro-observations” to describe entanglements, because like observations in standard interpretations, the claim here is that they transform the wave function by projection onto some basis, from an inside perspective. However, they differ from observations in standard interpretations and also differ from what we here will call macroscopic observations in that they are not performed by a human, but rather by other particles, and in that they can be easily erased. Performing a new observation on an orthogonal basis to an existing micromasurement will erase the information gathered by the previous measurement and create new objective uncertainty regarding the previously measured values.

³ (Colbeck & Renner, 2011) show that any extension of QM cannot yield improved predictions. In the interpretation presented here, knowledge of the hidden variable could yield better predictions, if it could be known from outside the system, but it cannot be known.

entanglement pushes the amount of uncertainty present down toward the minimum allowable limit⁴ (Heisenberg, 1925) (Ozawa, 2003) (Bastos, et al, 2015).

3. The basic idea and the delayed choice quantum erasure experiment.

Perhaps it is best to start with the simplest possible case as an example. Suppose we have one single random photon that we are not correlated with. From this outside perspective, we might write the wave function as:

$$|\psi\rangle = |\psi_{Initial}\rangle \quad (1)$$

Representing a superposition of all possible states. However, here we assert that there is a hidden variable, only visible from an inside view, correlated with the system. The photon already has a determined value on some basis. This value was determined by its most recent interaction.

$$|\psi\rangle = |U_+\rangle \quad (2)$$

This indicates a positive value on some unknown basis of measurement. But then, of course, on some orthogonal basis of measurement, say V, we could also write the wave function as a superposition.

$$|\psi\rangle = \frac{|V_+\rangle + |V_-\rangle}{\sqrt{2}} \quad (3)$$

In equations (2) and (3), we see the minimum possible uncertainty that must exist for any observer, even if they are correlated with the system. There is no uncertainty on some basis and objective uncertainty on some orthogonal basis. If we then consider equation (1), we can see that while it correctly predicts the results we will see if we measure multiple photons – namely, a random result on any basis on which we measure – it represents a mix of subjective and objective uncertainty. If we happen, by accident, to measure on the U basis, then our measurement will only resolve our subjective uncertainty as to the preexisting value. However, if we happen to measure on basis V, orthogonal to U, then our result is objectively undetermined until we measure, and a nondeterministic event occurs.

Let us now turn to a more complex example. In (Hobson, 2022), a pair of entangled photons or a biphoton is considered. Suppose two experimenters, Alice and Bob, each receive one of the pair. Each of the photon paths has been split into two paths they can follow with 50% probability. Because the states are correlated, if Alice receives a photon via path 1, then Bob will as well and the same for path 2. We can write the wave function from an outside perspective, which represents the minimum uncertainty we must have from that perspective as:

$$|\psi_{AB}\rangle = \frac{|A1\rangle|B1\rangle + |A2\rangle|B2\rangle}{\sqrt{2}} \quad (4)$$

Suppose Bob puts a phase shifter on one path and tries to get his photon to interfere with itself. He will not be successful. As (Hobson, 2022) points out, experiments have shown that this does not happen. (Hobson, 2022), citing (Horne, Shimony, & Zeilinger, 1990) and (Horne, Shimony, & Zeilinger,

⁴ Technically, there could be more than one hidden variable, so long as they were all compatible, since this is allowed by the uncertainty principle.

1989), also says that the theoretical reason why interference does not appear is that the nonlocal photon's contribution needs to be considered. When this is done, the phases always line up so that, as (Hobson, 2022) puts it, “the photons decohere each other”, that is, they prevent each other from creating an interference pattern. Thus, all experimental evidence of superposition has been eliminated in this configuration. (Hobson, 2022) contends that this represents a measured state⁵. The experimental setup discussed is diagrammed below (fig. 1):




Figure 1 – A diagram of the experimental apparatus discussed here and in (Hobson, 2022).

We can write the two-point nonlocal quantum field amplitudes at the detectors as:

$$\psi_{A1,B1} = \psi_{A2,B2} = \frac{1 + \exp(i\phi_B)}{2\sqrt{2}} \quad (5)$$

$$\psi_{A1,B2} = \psi_{A2,B1} = \frac{1 + \exp(i\phi_B + \pi)}{2\sqrt{2}} \quad (6)$$

where ϕ_B is the phase change imposed by Bob, and we have assumed that Alice is not altering her photon and that the various phase changes imposed by the experimental setup have been subsumed into a zero phase shift in eq. (5) and a shift of π in eq. (6).

The joint probabilities are then given by:

$$P(A1, B1) = P(A2, B2) = |\psi_{A2,B2}|^2 = \frac{1 + \cos(\phi_B)}{4} \quad (7)$$

and

$$P(A1, B2) = P(A2, B1) = |\psi_{A2,B1}|^2 = \frac{1 + \cos(\phi_B + \pi)}{4} \quad (8)$$

Then, we have $P(B1) = P(A1, B1) + P(A2, B1) = 0.5$ regardless of the phase. And in general, $P(A1) = P(A2) = P(B1) = P(B2) = 0.5$ regardless of any phase change added by Bob.

⁵ This is not the first suggestion that decoherence can solve the measurement problem (Bacciagaluppi, 2020), and while we do not completely agree with the presentation in (Hobson, 2022), it a good jumping off point for our discussion here, since it was while trying to understand the claim in (Hobson, 2022) that we had the idea for this paper.

In this specific configuration, where Alice happens to measure her photon on the basis that the photons measured each other and Bob has set his phase shift to zero, only subjective uncertainty remains. Let us suppose that the paths to detectors A1 and B1 represent reality. Then, to an outside observer uncorrelated with the system, who has subjective uncertainty, equation (4) still appears to be the correct description of the system and still describes their minimum possible uncertainty. Over many observations of many photons with unknown preestablished values on unknown bases, it will give correct statistical predictions. However, for a “micro-observer” entangled with the system, in this specific case, equation (9) is correct.

$$|\psi_{AB}\rangle = |A1\rangle|B1\rangle \quad (9)$$

The “coin” has already been flipped in this case. A nondeterministic event occurred as the photons separated. Thus, one crucial component of what we need a measurement to accomplish has already happened: projection onto a basis of measurement. One single entanglement counts as a “micro-observation”. We can legitimately speak of Alice’s photon as the observed system and Bob’s photon as our measuring device, which we have not yet queried. Note that this is not yet a macroscopic measurement; we can still do things such as erase the micromasurement and reintroduce objective uncertainty. However, for the moment, Bob’s photon has measured Alice’s photon and vice versa.

One might object that having this information preexisting when the photons are still together constitutes hidden variables. However, Bell’s inequality (Bell, 1964), (Maccone, 2013) only tells us that it is not possible for values on multiple incompatible orthogonal measurement bases to be preexisting (Napolitano & Sakurai, 2021). It is not possible in this example that a value on some unknown measurement basis, U, is preexisting and then also have preexisting values on an incompatible orthogonal basis such as V or W. Hidden variables sufficient to explain Bell nonlocality have been ruled out experimentally; however, that does not mean there cannot be an ANY hidden variable⁶.

Let us now have Bob change the phase by $\pi/2$.

$$P(B1) = P(A1, B1) + P(A2, B1) = \frac{1 + \cos(\pi/2)}{4} + \frac{1 + \cos(\pi/2 + \pi)}{4} = 0.25 + 0.25 = 0.5 \quad (10)$$

$$P(B2) = P(A2, B2) + P(A1, B2) = \frac{1 + \cos(\pi/2)}{4} + \frac{1 + \cos(\pi/2 + \pi)}{4} = 0.25 + 0.25 = 0.5 \quad (11)$$

We now have introduced new objective uncertainty because Bob’s basis of measurement has changed. Bob’s measurement is now completely uncorrelated with Alice’s, and the result is intrinsically uncertain before measurement. We still have a 50% subjective chance of A1 or A2, and in each case, there is a 50% objective chance of B1 or B2. That objective uncertainty will be resolved when Bob’s photon first becomes entangled with his measuring device. This distinction between types of uncertainty is invisible, however, in our equations. We might instead wish to write something like the following, where P_T , P_O , and P_S represent the total probability, the objective probability, and the subjective probability, respectively.

$$P_T(B1) = P_S(A1) * P_O(B1|A1) + P_S(A2) * P_O(B1|A2) \quad (12)$$

⁶ This idea takes the middle ground in the historical Einstein, Bohr debate where Einstein thought there must be sufficient hidden variables to avoid any nonlocality (Einstein, Podolsky, & Rosen, 1935), and Bohr believed the wave function was a complete description of the system.

$$P_T(B1) = 0.5 * \frac{1 + \cos(\phi_B)}{2} + 0.5 * \frac{1 + \cos(\phi_B + \pi)}{2} \quad (13)$$

$$P_T(B1) = 0.5 * 0.5 + 0.5 * 0.5 = 0.5 \quad (14)$$

A well-known result in quantum mechanics is that if we perform a measurement and then measure again on an incompatible orthogonal basis, the information gained from the first measurement is erased (Napolitano & Sakurai, 2021). Alice's photon measured Bob's photon, and Bob's photon was in a definite state where only subjective uncertainty existed until Bob erased this information and reintroduced objective uncertainty.

Some readers may notice that in this configuration, we are very close to the configuration of the delayed choice quantum erasure experiment (Kim, Yu, Kulik, Shih, & Marlan, 2000). We only need to have Alice direct her two beams each into one hole of a two-slit experiment and place Bob significantly farther from the photon source than Alice, and Bob then plays the role of the eraser. Standard interpretations with no hidden variables struggle to explain the results of this experiment without resorting to hypothesizing retroactive erasure of "which way" information⁷, and multiple hidden variables are ruled out by Bell's inequality. Our "one hidden variable interpretation", however, has no difficulty explaining the results of the experiment in an intuitively straight-forward way.

⁷ We do not want to imply that in standard interpretations the results of this experiment are entirely inexplicable without retrocausal effects. (Fankhauser, 2017) (Qureshi, 2020). However, we do try to show that the account here is more intuitive.




Figure 2 – Diagram of the delayed choice quantum eraser experiment. In this diagram, “Alice” would be the detector at D_0 , and “Bob”, perhaps temporarily renamed “Dan”, would be the detectors at D_1 and D_2 . We will, however, continue to refer to our hypothetical Alice and Bob experiment.

The experiment will of course involve many pairs of entangled photons, and each photon will measure the other half of its pair on some random basis. For our simple example here, we will assume they either measure each other’s path information or phase information rather than all possible bases on which they could measure each other⁸. There will then be populations of biphotons with preexisting values on some measurement basis. Those that happen to be predetermined to be on path A1 we can call “population A1”, and they will be objectively undetermined between the B detectors – these will only go through Alice’s slit number one. Additionally, those biphotons that happen to have measured each other’s phase information rather than path information and are now predetermined to arrive at detector B1 will be objectively undetermined on the A paths. Those photons will go through both of Alice’s slits and interfere with each other. This means that if we only look at the results on Alice’s detection screen that correspond to photons that were measured at detector B1, we will see an interference pattern. Suppose for simplicity, rather than every possible population, we have only four

⁸ Section 4 takes up the issue of what happens when the preexisting measurement basis does not match at least one of the experimenters’ bases.

populations - those biphotons that are 100% determined to be destined for detectors A1, A2, B1, or B2. From the inside perspective, we can write:

$$|\psi_A\rangle = |A1\rangle, |\psi_B\rangle = \frac{|B_1\rangle + |B_2\rangle}{\sqrt{2}} \quad (15)$$

$$|\psi_A\rangle = |A2\rangle, |\psi_B\rangle = \frac{|B_1\rangle - |B_2\rangle}{\sqrt{2}} \quad (16)$$

$$|\psi_B\rangle = |B2\rangle, |\psi_A\rangle = \frac{|A_1\rangle - |A_2\rangle}{\sqrt{2}} \quad (17)$$

$$|\psi_B\rangle = |B1\rangle, |\psi_A\rangle = \frac{|A_1\rangle + |A_2\rangle}{\sqrt{2}} \quad (18)$$

From an outside perspective, the system is still described as a superposition of all the possible states, and the wave function describes the minimum uncertainty that must be present from that outside perspective and still makes correct statistical predictions from that perspective. However, from the inside perspective, these different predetermined populations exist.

Detector B1 will pick up all the population B1 biphotons, of course, and then also half of each of the A1 and A2 populations. The A1 and A2 populations in the mix will smear the pattern out slightly on Alice's detection screen, but an interference pattern will remain. This is a heuristic argument, but it shows that one hidden variable is all we need to explain the results of this experiment intuitively. One "coin flip" already happened as the photons were created, so it does not matter how delayed Bob's measurement choice is. The population B1 biphotons were always going to be objectively undetermined on the A paths and interfere with themselves at Alice's detector. Population B2 biphotons interfere with themselves as well; however, the pattern is 180 degrees out of phase with the pattern produced by the B1 population. If viewed together, the patterns wash each other out. All Bob does, after Alice's results have been recorded, choose to look at some of the existing populations and not others. He does not retroactively erase "which way" information.




Figure 3 – Results of (Kim, Yu, Kulik, Shih, & Marlan, 2000) relabeled for our Alice and Bob thought experiment.

4. Nonlocality addressed

Now, let us suppose that both Alice and Bob alter their basis of measurement by $\pi/2$.

$$P(B1) = P(A1, B1) + P(A2, B1) = \frac{1 + \cos(\phi_B - \phi_A)}{4} + \frac{1 + \cos(\phi_B - \phi_A + \pi)}{4} = 0.5 \quad (19)$$

$$P(B2) = P(A2, B2) + P(A1, B2) = \frac{1 + \cos(\phi_B - \phi_A)}{4} + \frac{1 + \cos(\phi_B - \phi_A + \pi)}{4} = 0.5 \quad (20)$$

Both have now measured on a basis orthogonal to the basis on which the photons measured each other, and both results, taken individually, are objectively undetermined, but they perfectly correlate, even nonlocally, when considered together. Assuming we do not allow superdeterminism, this nonlocal correlation has been conclusively experimentally demonstrated (Hensen, Bernien, & Dréau, et al, 2015), (Storz, Schär, & Kulikov, et al., 2023). The description of the system in this case would be no different than in standard interpretations. From both the inside and outside perspectives, equation (4) describes the state of the system. There is, based on the discussion thus far, only objective uncertainty in this case, as knowledge of the hidden variable adds no useful information.

$$|\psi_{AB}\rangle = \frac{|A1\rangle|B1\rangle + |A2\rangle|B2\rangle}{\sqrt{2}} \quad (4)$$

The perfect correlation makes it appear as if Alice and Bob were somehow forced to measure on the preexisting basis, or alternately, the basis altered itself to match their measurement basis. The latter is what we will now assert happens. In order to match all experimental results, we propose the following: The biphoton has a “head” and a “tail”. One of the two photons, randomly, is the head or “control photon”. If Alice receives the control photon, she rotates the basis of the biphoton to match her basis of measurement. Bob’s half gets “dragged along for the ride” even nonlocally. If the photon is not a control photon, Alice leaves its basis unchanged. Those will have their basis altered to match Bob’s measurement basis, and then Alice’s photon gets dragged along with it to some new basis. She then simply measures relative to the new basis according to the Born rule. The analysis then proceeds just as in the previous section but on the new bases as if at least one of them had been the preexisting basis all along.

This has a number of benefits. For example, in the quantum eraser experiment, in the nonlocal case, there will ONLY be the 4 populations shown. Alice will reorient half of all biphotons to be populations A1 and A2, and Bob will reorient the other half to be populations B1 and B2. Additionally, a distinct benefit is that nonlocality becomes slightly less mysterious. One difficulty with instant correlation at a distance is that SR tells us that either Bob or Alice could be regarded as having acted first, so who influenced whom is a difficult question⁹. Our answer is that 50% of the time Alice influences Bob nonlocally and 50% of the time it is the other way around. Either way, we have only one nondeterministic event. If Alice reorients the basis of her photon, her result is predetermined,¹⁰ and Bob’s result for that photon is nondeterministic.

Note that assigning half the photons to Alice and half to Bob is not arbitrary or done merely to attempt to demystify nonlocality a little bit. The even division is needed in order to reproduce the results in experiments such as the quantum eraser. Assuming we exclude superdeterminism and assuming that micromasurements occur and create a hidden variable as we have asserted, and assuming the hidden polarization affects the next measurement in a standard way, according to the Born rule, then the rotations to the new basis of measurement we’ve proposed must take place, and they must be distributed half to each observer, and there must be a predetermined head and tail. Without the rotations, we would obtain incorrect probabilistic predictions. And, if Alice rotated all of the biphotons, we would only have populations A1 and A2, and if there were no predetermined head and tail and random “decisions” to rotate were made on each end, then some photons might be missed completely.

Let us suppose, as an example, that Alice sets her phase adjustment to $-\pi/3$ and Bob sets his to $\pi/3$. Without nonlocal effects, the single hidden variable interpretation would yield incorrect predictions in this case. It would say that Alice and Bob should agree with the value on the hidden measurement basis 75% of the time. They could then disagree no more than 50% of the time. However, standard theory predicts, and experiments show, they will disagree 75% of the time in this configuration. Thus, we need to bring nonlocal effects into the picture.

⁹ (Gillis, 2019) writes, “In general, it is very difficult to construct a coherent account of effects that are both nonlocal and nondeterministic without assuming some underlying sequence.”

¹⁰ To be more precise, it is predetermined following the rotation. We assume that the choice of which direction to rotate is partly nondeterministic and follows the standard Born rule.

Let us say that Alice receives the control photon. She reorients it to her basis of measurement. It is now as if, all along, she was measuring on the predetermined basis. Bob's photon rotates its basis with hers so that Bob is now at $2\pi/3$ relative to the predetermined measurement basis. Alice will find A1 half the time and A2 half the time in these cases, since this outcome is subjectively uncertain on the premeasurement basis. Bob's results can then be calculated to match hers 25% of the time and disagree 75% of the time.

$$P(B1|A1) = \frac{1 + \cos(\phi_B)}{2} = 0.25, \quad P(B1|A2) = \frac{1 + \cos(\phi_B + \pi)}{2} = 0.75 \quad (21)$$

The fact that only the heads of the biphotons cause nonlocal correlations leads to an obvious suggestion. Rather than treating the head and tail as Parity transformations of each other, we could treat them as CT transformations of each other. This would yield a future input dependent interpretation (Warton & Argaman, 2020) where the head of each biphoton actually originates at its respective detector. That is, population A1 photons originate at detector A1 and travel backward in time, etc. This would also yield a continual action interpretation, (Warton & Argaman, 2020). Communication between the photons is mediated by a chain of events. Basis rotation takes place over the entire trajectory of the biphoton. At least in the nonlocal case, treating the head of the biphoton as a CT transformation of the tail is certainly an attractive idea. At the very least, the biphoton behaves as if this were true. And while theoretically this involves particles traveling into the past, the measurable effects are only manifested nonlocally, not in the past light cone.

Finally, let us look again at the delayed choice quantum eraser experiment. In the delayed choice case, Alice acts first. In the cases where Alice receives the head of the biphoton, she will rotate those biphotons to match her path-information basis of measurement. These will be populations A1 and A2. We then have two options to explain what happens to the tails of the biphotons on Alice's side. One choice is to suppose that it is no different than the nonlocal case. Bob rotates the heads of these biphotons to his basis of measurement, and Alice measures relative to her new predetermined basis. However, this would involve Bob's measurement affecting Alice's measurement in his past light cone. This is the simpler option, but not the better option, in our view.

Alternately, treating the head and tail as CT transformations of each other might just be a useful way of treating the biphoton mathematically, without asserting that retrocausality is what actually happens. In the delayed choice case, it is possible to explain the results without supposing future input dependence. The time-forward alternative is that Alice performs all the basis rotations, although heads and tails are still treated differently. This preserves the standard time order of cause and effect. How would the tails "know" they needed to behave differently than in the nonlocal case? They would not yet have received a signal from the head that entanglement was broken. Similarly, the heads would "know" they had to behave differently once they received a signal from the tail that entanglement was broken.

Alice's unaltered tail photons will pass through both slits in various proportions and have their information determined by the recording screen in nondeterministic events. As the biphotons are measured at the detector, their other half will be rotated so that they continue to have values exactly opposite to each other on some basis. The information recorded on Alice's detection screen will be enough to largely separate the photons into populations B1 and B2 (see equations (17) and (18) and fig.

3).¹¹ Bob receives population A1 and A2 photons that are objectively undetermined on his basis and then also receives Alice's initially unrotated photons that were largely undetermined before measurement on her end and are now largely predetermined on his end to be either population B1 or population B2.

Note that we are still asserting that head and tail photons behave differently. The heads (populations A1 and A2) are rotated to specifically match her "which path" basis of measurement, as if they always had that basis, and the tails (populations B1 and B2) are not. The tails are just measured relative to whatever basis they happen to have, according to the Born rule, and then they drag their counterparts along for the ride so that anticorrelation between the photons is maintained. The tails' information will still be transmitted along the biphoton's entire trajectory, but in this case, the only measurable effect appears as communication into the future light cone at luminal or subluminal speeds between Alice's photon and Bob's photon.

A concrete example of the difference in behavior between head and tail particles seems needed here. Suppose we have one half of a stream of entangled electrons, and we put them into a magnetic field on the z-axis as in the Stern-Gerlach experiment. Each of the electrons will already have a definite spin on some random basis. Suppose one particular electron has a spin angle of 60 degrees from the z+ axis. If it is a tail electron, it will behave just like an unentangled electron in the magnetic field. It will split itself into two streams: 75% into the spin-up stream and 25% into the spin-down stream. The two parts will have the same net z-axis spin as the original electron. When the tail electron hits a detector, if it is still part of an entangled pair, it will rotate its counterpart, at that point in time, to maintain perfect anticorrelation.

The head electrons will behave differently. When they enter the magnetic field, they will not split into two beams. They will take one path or the other with 100% probability, destroying coherence, (Thiago De Oliveira & Caldeira, 2006), as in the photon experiments. To maintain the anticorrelation and the same net z-axis spin, they will drag their distant partner along to a new basis of rotation. They respond to the experimenter's preparation to measure on some basis. In the time reverse picture, then, they will appear to have originated 100% at one detector or the other.

One might be concerned that this would allow signaling. However, the experimenter can only control the hidden basis (in a random half of all cases) and not choose values on that basis. Bob's outer wave function and his probabilities are unchanged by Alice changing the basis of the hidden variable. If he could see the basis of the hidden variable, signaling would be trivial to accomplish. However, any attempt Bob might make to see it will just cause projection onto some new basis. The basis Bob finds will always be the one he chooses to find.

In theory, this is something a very sensitive experiment could test. Half of the electrons should cause a torque, in the apparatus, in a random direction, large enough to rotate the biparticle. The net average torque would be zero. Interestingly, if this torque could be measured exactly and added to the information acquired at the final detector, it would allow Alice to identify the value of the hidden variable on the basis on which the entangled electrons measured each other. This would not be possible with the tail electrons or with unentangled electrons.

¹¹ (Fankhauser, 2017) goes into extensive detail here.

Let us take stock of what we claim to have accomplished thus far. If this analysis is correct, the single hidden variable interpretation will replicate all the predictions of any standard quantum mechanical interpretation regarding the results of QM experiments. This would make choosing between interpretations mostly a philosophical matter. However, now we turn to the issue of the measurement problem where we believe the single hidden variable interpretation has a distinct advantage.

5. The measurement problem

In this section, we first attempt to answer three questions.

- 1) What happens in a micro-observation?
- 2) What happens in a macroscopic observation?
- 3) When do observations become nonerasable?

We then take a look at decoherence, which we assert can be reinterpreted, and a brief look at the new Wigner's friend thought experiment.

5.1 What happens in a micro-observation?

We simply suppose that the particles trade some bits of hidden information. Two electrons might, for example, trade hidden spin axes. Particles may also exchange information about position, momentum, or other variables. This is invisible from outside the system, but in effect, the particles have measured each other on some basis. In the case of the electrons trading spin axes, they “know” what the spin axis of the other particle was, and they changed it in the process of measurement. They must obey basic rules such as conservation laws and must not exchange more information than allowed by the uncertainty principle. In addition, the Born rule will play its standard role¹².



Figure 4 – Two particles exchanging information.

As a simple example, suppose an incoming electron is in a superposition of a $|z_+\rangle$ and a $|z_-\rangle$ spin state in a Stern-Gerlach experiment. An electron in the $|z_+\rangle$ branch is measured by an electron in a target. We would suppose that there would be a 50% chance that the target electron would acquire the

¹² This is admittedly vague, but models of exactly what information is exchanged under what circumstances are topics for future research. A starting point for this research would be the various currently existing models of information exchange during the process of decoherence, such as those described in (Schlosshauer, 2019). The following simple example should suffice for this paper, however.

$|z_+\rangle$ state, as the Born rule predicts. The incoming electron would then acquire the state of the target electron, and a new entanglement is formed. This simple trading of spin axes constitutes a micromasurement.

However, this seems to lead to information loss. Suppose the incoming electron had been in a pure $|x_+\rangle$ state before it acquired the $|z_+\rangle$ state. Was the information regarding the original state completely destroyed? Also, what happens if the resolution is that the incoming electron is in the $|z_-\rangle$ state, in which case the electron will be determined to not even be on the path where it is interacting? Additionally, when are old entanglements broken? We will return to these questions and what happens on the “sterile” path that the electron does not take in section 6. Here, we note that projection on to a new basis has occurred, from the point of view described by the inner wave function, and objective uncertainty has been resolved. From the point of view described by the outer wave function, a unitary interaction has taken place, and subjective uncertainty regarding this result will persist.

5.2 What happens in a macroscopic observation?

Let us now turn to macroscopic observations. After the photons in the original Alice and Bob experiment measure each other, we have a couple more steps to consider. First, one of the photons, let us say Bob’s, must become entangled with Bob’s measuring device, which then becomes correlated with Bob. We then have, from the point of view of an outside observer, not correlated with the system:

$$|\psi_B\rangle \otimes |Device\rangle \otimes |Bob\rangle \quad (22)$$

From Alice’s outside perspective, this still describes her probabilities. However, at this point, it represents 100% subjective uncertainty. Standard accounts would say that decoherence destroyed any possible evidence of superposition long ago. We assert that a myriad of micromasurements have destroyed the hypothetical global entanglement and created separate subsystems. No matter how clever an experiment Alice designs, she will not be able to coax Bob into interfering with himself. And obviously, Bob does not experience himself in a superposition; so, what is happening from the inside perspective? Something important happens in each of the three steps implied by eq. (22). First, objective uncertainty was removed at the particle level. Then, once entanglement with the macroscopic device takes place, the measurement can become thermodynamically irreversible. Whereas we could have erased the “memory” of a single photon, the result has now left an indelible mark on the universe. Finally, Bob looks at the measurement. Let us say that he finds that the photon took path 1. Now, we can finally write:

$$|\psi_B\rangle = |B1\rangle \quad (23)$$

With no uncertainty. No wave function collapse is needed here. Since Bob is now correlated with the system, he now knows which path has been taken. The outer wave function persists and describes the subjective uncertainty from the outside perspective for Alice until Bob chooses to inform her of the result. All that has happened is that Bob is now allowed to take the inside perspective, which has existed all along. Bob can now see a value for the formerly hidden variable on some basis he chose to measure.

To discuss what happens in the steps between single particle interactions and human beings looking at results, we need to distinguish between two different types of bases for measurement. There is an intended basis of measurement by the experimenter, and there are the bases of measurement used

by the micro-observers. The experimenter cannot completely control on which bases the micromasurements are made. However, she can control the experimental design, which will allow her to extract the desired sort of information. In a Stern-Gerlach experiment, for example, magnets separate electron streams according to spin, and then positional detectors are used to record the electrons. Even though Alice cannot control the exact basis of measurement for all the micromasurements, whatever positional information they do acquire will be enough to give her the desired information about the spin of the electron, since the paths are macroscopically separated. When multiple entanglements take place at her detector, any positional information the micromasurements acquire will be distinctly different than any positional data that would be acquired at the other detector. Thus, even if an individual micromasurement does not completely localize the particle, it will definitively identify which detector it has arrived at. The energy absorbed by the detector then sets off a chain of events, which Alice then interprets as a result regarding the electron spin. She does control something at the micromasurement level, however. If the electron hitting the detector is in a superposition of spin states, it will have its spin measured on the basis she chose when it interacts with the target, and it will be resolved into one of its two component states. Alice chose, via her experimental design, that it would be measured on this basis but did not have exact control of what other information was exchanged.

The actual physical process, in at least one of the actual quantum eraser experiments, involved a silicon avalanche photodiode (Ananthaswamy, 2018). The detected photon is absorbed and causes an electric signal involving billions of electrons. This process is too complex to analyze in detail, so let us turn to another thought experiment and a toy model. Suppose we have a “detector” consisting of just two particles. Alice is trying to determine the position of an incoming particle, so that is her chosen basis of measurement. However, Alice cannot control the basis on which each micro-observation will be made. Each of the particles will cause their own separate collapse of the inner wave function and project it onto some basis¹³. Let us suppose that particle 1 measures momentum more precisely and particle 2 measures position more precisely¹⁴. If we assume for this thought experiment that Alice can somehow directly read these results, she will have two separate positional estimates, and one will have less uncertainty than the other. She can then use some standard statistical procedure, such as the maximum likelihood method, to come up with her best possible estimate of the position. Thus, Alice does not actually obtain an eigenvalue as a result. She gets a messy classical-looking result. If this were a classic two-slit experiment, she obtains a finite-sized dot on her photographic plate, not an eigenvalue. She gets a statistical summary of just the positional portion of whatever information each of the micromasurements recorded, which she then interprets.

¹³ Our assertion that a collapse is taking place might raise concerns about conservation of energy. However, collapse models that conserve energy exist. (Gao, 2013) is an example. A key feature of this model is that it does not assume that collapse to a positional eigenstate occurs, and neither do we. Rather than a “collapse” of the inner wave function, a “localization” of the inner wave function is perhaps a better term, particularly in the case of continuous observables. However, when we are dealing with observables with discrete eigenstates, then collapse of the inner wave function to a single eigenstate is a valid description of the process.

¹⁴ What we have in mind here, is something like the collisional decoherence models discussed in section 4.1 of (Schlosshauer, 2019). There, various environmental interactions slowly destroy coherence. In the models presented, a short wavelength interaction will carry away more positional information than a long wavelength interaction. We are identifying these sorts of interactions with micromasurements. Thus, in the case of observables with continuous eigenvalues we hypothesize that even after a micromasurement, some objective uncertainty regarding these values remains. Micromasurements in these cases have “error bars”.

In the Stern-Gerlach example, suppose Alice is interested in the z-axis spin of an electron. The inner wave function, describing the objective uncertainty regarding the z-axis spin, collapses to a single eigenstate when it becomes entangled with her detector. The spin state that is determined is recorded in the spin state of another particle, and multiple position and momentum estimates are also acquired by the detector. However, Alice does not see any of these individual results. Her detector absorbs the incoming energy and sets off a chain of events. Her experimental design then allows her to infer the spin result from the summary information she does receive.

5.3 When do observations become nonerasable?

Now, we turn to the question of when measurements become nonerasable. In section 6, we argue that in this interpretation, nothing is ever truly erased; instead, the results can be rehidden. However, in this section, we will continue to use the term “erased” in the standard way. The question of when measurements can be effectively erased is still a very important one for many purposes even if this interpretation is correct. Quantum computing is probably the prime example. In the context of this interpretation, the question is “When can we no longer change subjective uncertainty back into objective uncertainty?” or “When can we no longer place the system in superposition and make a measurement orthogonal to a previous measurement?”.

Let us start with the famous Schrödinger’s cat thought experiment (Schrödinger, 1937). The first thing to note is that we cannot even describe what sort of physical process we would use to perform an orthogonal measurement on a complex macroscopic system. Let us first assume that we can accomplish isolating the system from the rest of the universe. How do we then perform a measurement orthogonal to a previous measurement? If the box is spinning on the z-axis, it is not in a superposition of x-axis spin states. The cat is not in superposition. It is nonsensical.




Figure 5 – Schrödinger’s cat.

When dealing with subatomic particles, both the outer and inner wave functions are useful concepts. However, for this macroscopic system, the outer wave function is now too far removed from the inner wave function. The inner wave function is 100% objective uncertainty and still describes quantum objects, whereas the outer wave function is by now 100% subjective uncertainty and cannot be changed back to objective uncertainty. Trying to project the outer wave function onto some new basis is an invalid move. Superposition is no longer possible. The theoretical global entanglement was destroyed by a myriad of micromasurements. We can no longer use the outer wave function to describe any new global manipulations we might make. The outer wave function still represents the subjective uncertainty regarding the fate of the cat, but that is its only useful function now. It has no practical value for predicting new measurements. It is a matrix of uncountable dimensions for which we cannot even fill in values¹⁵. However, it will persist as long as we keep writing it down, and it will continue to describe the probability for anyone not yet correlated with the system until they open the box and look at the cat. Until then, it still describes their epistemological state but no longer describes a potential uninstantiated alternate reality. That path is now permanently closed. (So hopefully the cat is still alive).

Let us now think about a carefully controlled quantum state. The inner wave function will describe objective uncertainty and quantum phenomena such as superposition, and the outer wave function will not be far removed from the inner wave function, if it differs at all. Now, if we know what basis a previous measurement used, we will know what on what basis to perform an orthogonal measurement, and we will be able to erase the previous result. Problems will arise as soon as any interactions not fully under our control become involved, however¹⁶. We will not know the basis of measurement for a random micromasurement, so we will not know what basis is orthogonal to it. (Think of the position and momentum measurements made by the two-particle detector). And given that each micromasurement causes its own localization of the internal wave function, things can get out of control very quickly. We would now have to make a myriad of separate, individualized, micromasurements. We are now dealing with thermodynamic irreversibility. To erase a macroscopic result, we would have to, in effect, reassemble the dead version of Schrödinger's cat one subatomic particle at a time.

So, when can we no longer change subjective uncertainty back into objective uncertainty in this interpretation? In practice, this happens when a variable we are interested in is micromasured in a way we cannot control, and thus we no longer know how to perform an orthogonal measurement on the system or its parts. Roughly, as in standard interpretations, this happens when uncontrolled environmental interaction causes what is commonly known as decoherence, which destroys the phenomenon of superposition.

5.4 Decoherence.

¹⁵ (Nielsen & Chuang, 2016) points out that the state of a system with only 500 qbits, many orders of magnitude simpler than our hypothetical cat, would take 2^{500} amplitudes to specify. This number is larger than the number of atoms in the known universe. It seems unreasonable that nature would be keeping track of that quantity of information in such a small system. It seems more reasonable to suppose nature continually simplifies the wave function. This is accomplished by the destruction of global entanglements, as we discuss in section 6, and by the collapse/localization of the inner wave function.

¹⁶ This is, of course, the bane of quantum computing.

(Schlosshauer, 2019) extensively discusses the well-studied phenomenon of decoherence. He writes “effectively, the environment is performing nondemolition measurements on the system” and “This suggests that decoherence can indeed be understood as an indirect measurement—a monitoring—of the system by its environment”. (Schlosshauer, 2019) also points out that decoherence by itself cannot solve the measurement problem, in part because decoherence theory is based on the unitary development of the wave function. These comments fit very well with our interpretation, since we assert that the outer, or standard, wave function never collapses but also that “under the hood”, the environment can perform uncontrolled micromasurements, collapsing the inner wave function and transforming the outer wave function into pure subjective uncertainty.

A disagreement we have is that (Schlosshauer, 2019) states that after local coherence is destroyed, only a global measurement of the system that includes the environment could reveal it. We do not believe that such a global measurement, showing superposition, would be possible. The roots of this disagreement can be found in the simplest example in (Paz & Zurek, 2000). Here, three one-bit systems are used to represent a measured system, an apparatus, and the environment. The system and the apparatus are first entangled, and it is stated that this cannot constitute a measurement, just an entanglement.

$$(\alpha | \uparrow \rangle + (\beta | \downarrow \rangle)|A_0 \rangle \rightarrow \alpha | \uparrow \rangle|A_1 \rangle + \beta | \downarrow \rangle|A_0 \rangle = \Phi \quad (24)$$

Then, bringing in the third system, an environment bit that has premeasured the apparatus, allows us to look at a reduced density matrix representing the first two systems. This is accomplished by tracing over the environment.

$$\rho_{AS} = \text{Tr}_E |\Psi\rangle \langle \Psi| = \alpha^2 | \uparrow \rangle \langle \uparrow | |A_1 \rangle \langle A_1| + \beta^2 | \downarrow \rangle \langle \downarrow | |A_0 \rangle \langle A_0| \quad (25)$$

This allows us to see a classical “OR” instead of a quantum “AND” in this specific case. (Paz & Zurek, 2000) conclude section (2.4) by writing, “*Disappearance of quantum coherence because of a ‘one-bit’ measurement has been verified experimentally... A single act of quantum measurement we have discussed here should be regarded as an elementary discrete instance of continuous monitoring, which is required to bring about the appearance of classicality.*”

We agree that it looks like a measurement has taken place, but that is, we argue, because a micromasurement actually has taken place, causing a collapse of the inner wave function to a single eigenstate. We also agree, however, that no collapse of the outer wave function takes place. It just represents subjective uncertainty at this point. If this was only a micro-observation, then the outer wave function could be converted back into objective uncertainty. However, if a macroscopic apparatus was involved, then no global process, including bringing the environment into the picture, would allow superposition to be recreated.

We suggest a different interpretation of the same three bit experiment. Suppose the environment bit that is “monitoring” the system instead represents a micro-observer. The observer is pre-entangled with the apparatus. This allows the observer to take the view of the inner wave function, where in this case objective uncertainty has been resolved¹⁷. Equation (25) then reflects our subjective

¹⁷ In (Paz & Zurek, 2000) “if the states of the environment are correlated with the simple products of the states of the apparatus–system combination” then the result is that the reduced density matrix appears classical. We would say that only subjective uncertainty remains.

uncertainty about a result that already exists. With that one minor reinterpretation of the role of the environment/micro-observer bit, the reduced density matrix in decoherence theory can be reunderstood to describe the probable results of micromasurements, the actual results of which would be described by the inner wave function. We want to emphasize this stunning conclusion. All the work that has been done studying the phenomenon of decoherence can be easily reinterpreted to be about the collapse or localization of the inner wave function.

The difference in interpretation is that we assert that “coin flips” have taken place “under the hood” where we cannot see them, leaving us with only epistemological uncertainty about their results. A key insight in this interpretation is that the wave function can both collapse and persist. The outer wave function describing total uncertainty persists, while inner wave function collapse takes place, and projection occurs. However, the standard interpretation of decoherence theory says that we only have the “*appearance*” of classicality, with no collapse taking place. This does not lead to any differences in predictions, as far as we can tell at this point, except that we do not predict that half alive, half dead cats can be created by any sort of global procedure. That is, as long as we focus on only the reduced matrix, there will be no disagreement. The disagreement occurs only when we talk about the global matrix, which includes the environment. We assert that micromasurements destroy the hypothetical global entanglement, creating separate subsystems. Again, we return to the topic of entanglement destruction in section 6.

However, even when we look at a global system that is still entangled, before separate subsystems are formed, there are no differences in predictions. Let us look at an example involving continuous observables¹⁸. In section 7.6.2, “Decoherence and the destruction of cat states”, (Benenti, Casati, Rossini, & Strini, 2019) gives a standard example of two Gaussian wave packets moving along the x-axis in superposition.

$$\psi_{cat}(x) \equiv \frac{1}{\sqrt{2}} [\psi_+(x) + \psi_-(x)] \quad (26)$$

$$\langle x | \rho_{cat} | x' \rangle = \frac{1}{2} [\psi_+(x)\psi_+(x') + \psi_-(x)\psi_-(x') + \psi_+(x)\psi_-(x') + \psi_-(x)\psi_+(x')] \quad (27)$$

A photon that is much lighter than the system is scattered off the system. It is shown that this destroys the visible signs of superposition. The diagonal terms of the density matrix for the original cat states are almost unaffected, but the off-diagonal terms are eliminated. Standard interpretations would assert that the global system, including the photon, could still be described as a superposition. We assert that a micromasurement has taken place, and there is currently only subjective uncertainty as to the result. However, as in our original Alice and Bob example, this distinction does not offer any differences in predictions. The difference is invisible. If we take X to represent the position of the large system and x to represent the photon state correlated with the position of the measured system, then we assert that the current state, from the inner perspective, is:

$$|\psi_{xx}\rangle = |X_-\rangle|x_-\rangle \text{ or } |X_+\rangle|x_+\rangle \quad (28)$$

¹⁸ A detailed version of this interpretation that applies to continuous observables is another topic for future research. However, a point to note is that for continuous observables, micromasurements only reduce objective uncertainty by converting it to subjective uncertainty. They do not eliminate all objective uncertainty regarding continuous observables. We would also assert that the photon in this example measures a combination of position and momentum. (Gampel & Gajda, 2023) offers current insight into such simultaneous measurements.

Rather than:

$$|\psi_{xx}\rangle = \frac{|X_{-}\rangle|x_{-}\rangle + |X_{+}\rangle|x_{+}\rangle}{\sqrt{2}} \quad (29)$$

This is essentially the same state as the original state in our Alice and Bob experiment. Only subjective uncertainty exists, but objective uncertainty could theoretically be restored by orthogonal measurement. Entanglement will prevent evidence of superposition from appearing in this configuration, according to standard interpretations. There is no observable difference between the interpretations.

Given that for experiments we can actually do, there is no difference in observable predictions between the standard interpretation of decoherence and our interpretation, all we can do to illustrate our perspective is compare the differences in interpretation step by step. In the three bit experiment, in the two particle case, (Paz & Zurek, 2000) state that no measurement has taken place and that ambiguity regarding bases is present. We assert that a micromasurement has taken place but that we have subjective uncertainty as to what basis was used. When the third particle becomes involved, (Paz & Zurek, 2000) state that the basis uncertainty has been resolved. The environment bit determines what basis will be used for the apparent measurement or environmental monitoring. We would say that because the third bit is pre-entangled with the system, we now have access to the inner wave function and can see what measurement basis was used. And, if we could directly query our one bit apparatus, we would see that its hidden variable contains information about what value was discovered. Measurement has taken place, and if we are correlated with our measurement apparatus, we will have access to the result of the measurement.

5.5 The new Wigner's friend thought experiment.

Before moving on to the next section, let us take a brief look at the new Wigner's friend thought experiment (Frauchiger, 2018), (Bong, Utreras-Alarcón, & Ghafari, et al., 2020), (Ormrod, Vilasini, & Barrett, 2023). It treats a macroscopic observer as a quantum system in a larger experiment and arrives at a contradiction. The contradiction can be resolved but only at a cost. Different authors have published proofs that enumerate all the logical possibilities. This provides a couple of different taxonomical systems with which to classify interpretations. One issue with this thought exercise is that what constitutes a measurement is not defined. We will provide our definitions. It is then an interesting exercise to see where this interpretation falls in the taxonomies.

In short, in the thought experiment (Ormrod, Vilasini, & Barrett, 2023), Alice and Charlie have a spacelike separation from Daniella and Bob. All perform measurements. Daniella has an inside perspective and precedes Bob, who has an outside perspective. If Daniella is a macroscopic observer, then we would claim that in practice, Bob cannot perform an orthogonal measurement. It is possible in theory, but only if he has perfect knowledge and control of every subatomic particle and erases all the previous results, one by one. He has a myriad of orthogonal measurements to perform. Alternately, if Daniella is a micro-observer, then Bob can erase her result. Or as we clarify in the next section, his measurement, in effect, changes the answer, erases her memory, and hides her original answer away where no one can see it.

So, where does this interpretation fall on the taxonomies? We clearly avoid some difficult ideas, such as superdeterminism and many-world hypotheses. In other cases, it is more of a “yes and no” answer. Does the interpretation allow superluminal causation? Yes, but only hidden variables are affected, and there can be no signaling. Is there a wave function collapse? This is a complex question. No, for the standard wave function that describes the outer perspective, although in the next section we argue that when entanglements are broken, it is divided into separate descriptions of separate subsystems. However, it still persists and correctly describes subjective probabilities for external observers not correlated with the observed system, and it never collapses to a single eigenstate. The inner wave function collapses to a single eigenstate in the case of discrete observables, although it never collapses to a single positional eigenstate, so it might be better to say that it is just ‘localized’ in the case of continuous observables. And, in section 6, we argue that there is no information loss in this process, as nothing is actually erased.

Finally, are observations absolute? Yes, macroscopic observations are absolute and cannot be erased/rehidden in practice. Destruction of global entanglements and thermodynamic irreversibility prevent this. No, micro-observations are not absolute, as their results can be rehidden, but even they cannot be truly erased, and this is the topic to which we now turn.

6. Nonlocal effects and the measurement problem.

In this section, we introduce nonlocal effects, similar to those discussed in section 4, to explain some key features of the measurement process. Unlike the assertions in section 4, there is no one clear chain of deductive reasoning from our original hidden variable assumption to the assertions made in this section that motivates these assertions. However, we hypothesize that similar nonlocal effects might be involved in both entanglement and wave function collapse.

6.1 Not erased, just rehidden.

Until this point, we have addressed the question of erasability as a question of being able to measure on an orthogonal basis. That is, if the experimenter can manage a real-life orthogonal measurement of the system, then it is erasable; if she cannot, then it is nonerasable. Having multiple micromasurements instead of one macroscopic measurement to erase just makes this task significantly more difficult to accomplish than one might have expected¹⁹. However, now we ask if we should even talk about “erasure”. In the single hidden variable interpretation, all we really do is rehide information. We make the information unavailable in the description given by the outer wave function, but from an inner perspective, the information never truly goes away.

¹⁹ (Schlosshauer, 2019) writes “Decoherence is a genuinely quantum-mechanical effect, to be carefully distinguished from classical dissipation and stochastic fluctuations. One of the most surprising aspects of the decoherence process is its extreme efficiency...”.

Let us look at a prototypical example of erasure and see what this interpretation says is going on. Figure 4 shows a sequential Stern-Gerlach experiment.




Figure 6 – A sequential Stern-Gerlach experiment.

We initially prepare a steam of electrons (actually silver atoms, in practice) with a $z+$ spin. In this case, the inner and outer wave functions are now identical. We have:

$$|\psi\rangle = |z_+\rangle \quad (30)$$

We then put it through the SGx apparatus. Now, it is best described as:

$$|\psi\rangle = \frac{|x_+\rangle + |x_-\rangle}{\sqrt{2}} \quad (31)$$

This is both the exterior and interior wave function. Objective uncertainty exists on the x-axis. The input into the next step, however, is best written as:

$$|\psi\rangle = |x_+\rangle \quad (32)$$

Which is then subsequently best written as:

$$|\psi\rangle = \frac{|z_+\rangle + |z_-\rangle}{\sqrt{2}} \quad (33)$$

What happened between equation (31) and equation (32)? Measurement and projection of course. But where did the information regarding the previous measurement go? We assert that it “disappeared” into the wall hit by the $|x_-\rangle$ beam. Similar to entanglement, wave function collapse is also a nonlocal process. In fact, this is what Einstein originally meant by “spooky action at a distance” (Ananthaswamy, 2018). As in the case of entanglement, we assert that information is transmitted along both paths of the electron. When the $|x_-\rangle$ beam hits the wall, it changes from 50% $|x_-\rangle$ to 0% $|x_-\rangle$. It leaves hidden basis information with the particles in the wall. It then communicates this information back along its path to the moment it split from the $|x_+\rangle$ beam, and then the information propagates forward along the $|x_+\rangle$ path. Thus, the electron in the $|x_+\rangle$ beam “knows” it is now 100% $|x_+\rangle$.

So did the $|z_+\rangle$ spin information truly disappear? No, it just became unretrievable in the wall. To retrieve that information, we would have to have known the exact state of the hidden variables of the particles in the wall before they interacted with the electron. From an outer perspective, we don’t have

access to that information. Thus, we might wish to change terminology.²⁰ Rather than asking if the experimenter can erase the information via orthogonal measurement, we might ask instead if the experimenter can reconceal the discovered information via orthogonal measurement.

Our assertion that information is deposited in the wall by the $|x_{-}\rangle$ beam on a path where there is no energy exchange might be easier to accept if we consider the quantum bomb experiment (Elitzur & Vaidman, 1993). From that experiment, we know that quantum systems can discover information about a path the “particle” does not travel. It should not be surprising then that the system can communicate information to an object on the sterile path as well.

But what if the electron hits the wall and is resolved into an $|x_{-}\rangle$ state? It had been in a $|z_{+}\rangle$ state. Where does this information go if the $|x_{+}\rangle$ beam does not hit a wall anytime soon? We must suppose that the sterile beam continues to carry this information. Sometime in the future, it will deposit this information into a target that can accommodate it. The target would receive zero energy transfer, but a torque would be applied to it. To picture the situation, imagine that the real particle has a missing “ghost” particle that it would have if it were part of an entangled pair. The ghost electron would have, effectively, been in a $|z_{-}\rangle$ state:

$$|\psi\rangle = |z_{-}\rangle = \frac{|x_{+}\rangle - |x_{-}\rangle}{\sqrt{2}} \quad (34)$$

And would be rotated to an $|x_{+}\rangle$ state. All the hidden spin values would balance in this case. The incoming $|z_{+}\rangle$ and this $|z_{-}\rangle$ are both gone. The “live” branch created an $|x_{-}\rangle$ state in the wall it hit, where it also transferred energy and linear momentum when it traded states with an electron in the wall. The sterile branch would now be in an $|x_{+}\rangle$ state. There is no real electron on the sterile path, however. But we hypothesize that the torque to rotate the live branch will still come from this branch. The ghost electron would need to receive a positive y-axis torque, and therefore, a negative y-axis torque should appear in the target. As in the case of entangled photons, we hypothesize that a basis rotation then propagates along the entire length of both potential electron paths. The effect in the target would likely be manifested in other hidden particle spin states but also, perhaps, in a macroscopic object’s angular momentum.

This might be something a cleverly designed experiment could detect. A difficult aspect of this would be that there would be no other sign that a zero energy “particle” hit a target, other than a tiny torque. A negative result might mean it missed the target or “declined” to interact. In addition, the acquired angular momentum would likely only be found in a hidden variable in the target, making it undetectable. On the other hand, it does not seem impossible that a tiny target could show a measurable effect, particularly if it could be hit many times. (Galvez & Zhelev, 2007) describe an experiment involving photons imparting angular momentum to a latex sphere 5 μm in diameter trapped in an optical tweezer and suspended in oil. Something like this is what we have in mind since photons would be easier to work with than electrons. A difference is that their experiment involved photon orbital angular momentum, and this experiment would involve photon spin information. Suppose we split a vertically polarized photon into an LHC polarized beam and an RHC polarized beam. We block the RHC polarized beam and direct the other beam at a tiny target. If the live photons can be induced to produce a measurable change in angular momentum in the target, then perhaps the ghost photons can

²⁰ Although, ironically, the term “erasure” is likely permanent.

as well. The difference would be that the ghost photons, carrying no energy, could only alter the orientation of the angular momentum in the target, not increase the total angular momentum. For the most part, at least, we would expect that only hidden information would be affected. However, with enough interactions, a macroscopic effect might be produced. This would constitute experimental verification of a unique prediction of this interpretation.

6.2 Destruction of entanglements.

One final assertion we make here regards the question of exactly when old entanglements are broken. We assert that interactions on the sterile branches that particles do not follow are critical to breaking old entanglements without forming new ones. For example, between eqs. (31) and (32) the most obvious thing that happens is that the $|x_{-}\rangle$ beam interacts with a target. Given the other assertions of this interpretation, in particular the assertion that live branch interactions that form entanglements are the loci for micromasurements, sterile branch interactions are nearly the only place entanglements could be broken, thus this is what we propose takes place. To the best of our knowledge there is no experimental data that would contradict this assertion.

Breaking of entanglements is the final important piece of the measurement puzzle. Although, recall that breaking entanglements will not by itself guarantee that values discovered in a micromasurement of a system cannot be erased/rehidden, it will just create separate subsystems, guaranteeing that no global process can accomplish “erasure” and/or recreation of superposition. It will prevent us from creating a Schrödinger’s cat via some hypothetical global manipulation, and thermodynamic irreversibility will prevent us from creating a superimposed cat via a process that involves addressing one subsystem at a time.

While interactions on both the live and sterile branches can affect the values of hidden variables, this is the only proposal we have made here that would affect the outer or standard wave function in any way. All real particle interactions are still treated as unitary interactions, but we assert that sterile branch “interactions” should be treated as events that separate the system the wave function describes into separate subsystems. For example in the diagram below, the indicated source is a source of entangled particles in singlet states. Then, looking only at the particles that are received at Bob’s x-axis detectors:




Figure 7 – A diagram of an experiment used to show when entanglement is broken.

$$|\psi_{AB}\rangle = \frac{|Ax+\rangle|Bx-\rangle + |Ax-\rangle|Bx+\rangle}{\sqrt{2}} \quad (35)$$

Becomes:

$$|\psi_A\rangle = \frac{|x_+\rangle + |x_-\rangle}{\sqrt{2}} \text{ and } |\psi_B\rangle = \frac{|x_+\rangle + |x_-\rangle}{\sqrt{2}} \quad (36)$$

A and B are now independent, and we assert that the transition happened when an information transfer event occurred during the sterile branch's interaction with the target. From the moment entanglement ends, going forward, the outer wave function will continue to yield correct subjective probabilities regarding previous results. However, if it is used to describe any new manipulations of the system, it will be incorrect.

6.3 How should we define a measurement?

Finally, we can turn to the general question of “What constitutes a measurement?” If what we mean by a measurement is that it causes projection onto some basis and causes some objective uncertainty to be resolved in a nondeterministic event and that the information acquired cannot then subsequently be truly erased, then micromerasurements are measurements, period. If we “erase” a measurement, that does not change the fact that it happened, and the information gathered is not truly destroyed. We transform the system, and this hides the result from us. We change the old value in (30) to the new value in (32) and soon or later deposit information about the original answer somewhere we likely cannot retrieve it. This explains how macroscopic observations can be truly nonerasable. One might ask how even a myriad of erasable micromerasurements could add up to a nonerasable macroscopic measurement. The answer is that the micromerasurements are not truly erasable either.

However, if we want our definition of a measurement to also include the destruction of existing entanglements, then we must be sure to include sterile branch information transfer events as part of our definition of a measurement. Adding this component to the definition would yield a definition that says that measurement results cannot be rehidden via any global process.

Or if we want our definition of a measurement to stipulate that results cannot be rehidden via any process, including processes that address individual subsystems separately, then in practice, the question of what exactly constitutes a measurement will come down to a question of exactly how clever experimenters can be in concealing information from themselves in erasure experiments that do not, in general, replicate natural processes. In theory, nothing other than thermodynamic irreversibility prevents arbitrarily large systems from being treated in this manner.

Our preference is to use the term “measurement”, without qualification, to refer to events that include the destruction of global entanglements. Including this feature in the definition gives measurements an important form of irreversibility. We prefer to continue to use the term “micromerement” to refer to any new entanglement. The key feature here is that a nondeterministic event occurs, removing at least some objective uncertainty. Finally, we prefer to use the term “macroscopic measurement” once thermodynamic irreversibility has become a factor. In our view, the involvement of a human mind, when an observer becomes correlated with an observed system and can take the inside perspective, is not a critical part of the measurement process.

7. Summary

We treat all new entanglements as micro-observations from an inside perspective. These observations transform objective uncertainty into subjective uncertainty on the basis measured but

leave values on other potential bases of measurement objectively undetermined. If we look again at equation (4):

$$|\psi_{AB}\rangle = \frac{|A1\rangle|B1\rangle + |A2\rangle|B2\rangle}{\sqrt{2}} \quad (4)$$

Depending on the basis on which Alice and Bob choose to measure, it may represent completely subjective uncertainty in the case where they both measure on the same basis on which the photons measured each other. Alternately, it may represent completely objective uncertainty for an experimenter measuring on a basis orthogonal to the premeasurement basis. Alternatively, it may represent a combination of both if they measure on some other basis.

The standard or outer wave function represents the total probability, objective plus subjective, and the minimal uncertainty that any outside observer must have. The objective probability is what is uncertain to an observer that is part of the system, the minimal uncertainty that any observer must have. It is represented by an inner wave function. The difference between them is that from the inner perspective, one can see the value of hidden variables.

We have provided a theory of nonlocality that is a direct consequence of our hidden variable assumption and a desire to match all experimental data. In this theory, biparticles have heads and tails, and the heads behave differently than unentangled particles. They orient their preexisting basis of measurement to match the experimenter's basis of measurement and reorient the tail end of the biparticle along with them, even nonlocally. A very sensitive experiment might be able to test for this effect.

Micro-observations are simply particle interactions where the particles exchange allowable bits of hidden information. They might, for example, alter each other's basis of rotation. Nondeterministic events obeying the Born rule occur, which resolve at least some objective uncertainty. Macroscopic observations should not be treated as projections onto a basis. Rather, the observer simply gains access to the interior wave function and finds a statistical result built up from many micro-observations. Micro-observations can be "erased", and macroscopic observations cannot. However, we argue that "erasure" is truly the wrong term in this interpretation. Information is instead rehidden and replaced by a new value.

We also argue that the reduced density matrix in decoherence theory can be reunderstood to describe the probable results of micromasurements, the actual results of which would be described by the inner wave function. We also argue that interactions on sterile paths that particles do not follow are critical to destroying previous entanglements without creating new entanglements, thus creating separate subsystems rather than a continuing global entanglement. We also argue that the definition of a measurement, unqualified by "micro" or "macroscopic", should be that a measurement happens when global entanglement ends, and thus, no global procedure can then cause "erasure" of existing results.

This gives us an interpretation of quantum mechanics that avoids difficult ideas such as nonabsolute macroscopic events, many-world hypotheses, superdeterminism, information loss, and superluminal causation (affecting nonhidden variables). It intuitively explains the results of the quantum eraser experiment, and it offers a theory of nonlocality. It offers a proposed solution to the measurement problem and a reinterpretation of decoherence. Finally, it proposes an explanation for how entanglements are destroyed, and it banishes cats in superposition from QM.

A challenge for future research would be to attempt to verify this interpretation experimentally. Additionally, if this QM interpretation is successful, yet another challenge would be to develop a QFT version of it. Additional projects for future research have also been described in the text.

References

- Ananthaswamy, A. (2018). *Through Two Doors at Once: The Elegant Experiment That Captures the Enigma of Our Quantum Reality*. Penguin.
- Bacciagaluppi, G. (2020). The Role of Decoherence in Quantum Mechanics. *The Stanford Encyclopedia of Philosophy*. (E. N. Zalta, Ed.) Stanford, California, USA: The Metaphysics Research Lab. Retrieved from <https://plato.stanford.edu/entries/qm-decoherence/>
- Barzegar, A., & Oriti, D. (2022). Epistemic-Pragmatist Interpretations of Quantum Mechanics: A Comparative Assessment. *Physical Review*, 47, 777-780. Retrieved from arXiv:2210.13620
- Bastos, et al, C. (2015). Robertson-Schrödinger formulation of Ozawa's uncertainty principle. *Journal of Physics: Conference Series*, 626. Retrieved from <https://iopscience.iop.org/article/10.1088/1742-6596/626/1/012050/pdf>
- Bell, J. S. (1964). On The Einsein Podolsky Rosen Paradox. *Physics*, 1(3), 195-200. Retrieved from https://cds.cern.ch/record/111654/files/vol1p195-200_001.pdf
- Benenti, G., Casati, G., Rossini, D., & Strini, G. (2019). *Principles of Quantum Computation and Information*. Danvers, MA, USA: World Scientific Publishing.
- Bong, K., Utreras-Alarcón, A., & Ghafari, et al., F. (2020). A strong no-go theorem on the Wigner's friend paradox. *Nat. Phys.*, 16, 1199–1205. Retrieved from <https://doi.org/10.1038/s41567-020-0990-x>
- Colbeck, R., & Renner, R. (2011). No extension of quantum theory can have improved predictive power. *Nature Communications.*, 2(8), 411.
- Einstien, A., Podolsky, B., & Rosen, N. (1935, May 15). Can Quantum-Mechanical Description of Physical Reality Be Considered Complete? *Physical Review*, 47, 777-780.
- Elitzur, A. C., & Vaidman, L. (1993). Quantum mechanical interaction-free measurements. *Foundations of Physics*, 23(7), 987–997. doi:10.1007/BF00736012
- Fankhauser, J. (2017, July). Taming the Delayed Choice Quantum Eraser. *Quanta*, 8(1). doi:10.12743/quanta.v8i1.88
- Frauchiger, D. R. (2018). Quantum theory cannot consistently describe the use of itself. *Commun*, 9, 3711. Retrieved from <https://doi.org/10.1038/s41467-018-05739-8>
- Galvez, E. J., & Zhelev, N. (2007). Orbital Angular Momentum of Light in Optics Instruction. *Education and Training in Optics and Photonics*. doi:<https://doi.org/10.1364/ETOP.2007.ESB3>
- Gampel, F., & Gajda, M. (2023, Jan). Continuous simultaneous measurement of position and momentum of a particle. *Phys. Rev. A*, 107(1), 012420. doi:10.1103/PhysRevA.107.012420

- Gao, S. (2013, May 08). A discrete model of energy-conserved wave function collapse. *Proceedings of the Royal Society A*, 469(2153). doi:<https://doi.org/10.1098/rspa.2012.0526>
- Gillis, E. J. (2019, April 5). Wave Function Collapse and the No-Superluminal-Signaling Principle. *International Journal of Quantum Foundations*, 5(2). doi:10.48550/arXiv.1501.03198
- Heisenberg, W. (1925). Über quantentheoretische Umdeutung kinematischer und mechanischer Beziehungen. *Zeitschrift für Physik*, 33, 879–893.
- Hensen, B., Bernien, H., & Dréau, et al. (2015). Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres. *Nature*, 526, 682–686.
- Hobson, A. (2022, March). Entanglement and the Measurement Problem. *Quantum Engineering*, 2022. Retrieved from <https://doi.org/10.1155/2022/5889159>
- Horne, M. A., Shimony, A., & Zeilinger, A. (1989). “Two-Particle interferometry”. *Physical Review Letters*, 62(19), pp. 2209–2212.
- Horne, M. A., Shimony, A., & Zeilinger, A. (1990). “Introduction to two-particle interferometry”. In *Sixty-Two Years of Uncertainty* (pp. pp. 113–119). New York, NY, USA: Plenum Press.
- Jaynes, E. T. (2003). *Probability Theory*. (G. L. Bretthorst, Ed.) Cambridge, UK: Cambridge University Press.
- Kim, Y.-H., Yu, R., Kulik, S. P., Shih, Y. H., & Marlan, O. S. (2000, February 2000). A Delayed Choice Quantum Eraser. *Physical Review Letters*.
- Maccone, L. (2013). A simple proof of Bell's inequality. *American Journal of Physics*, 81(854).
- Monroe, C., Meekhof, D. M., King, B. E., & Wineland, D. J. (1996, May 24). A "Schrödinger Cat" Superposition state of an atom. *Science*, 272.
- Napolitano, J. J., & Sakurai, J. (2021). *Modern Quantum Mechanics*. Cambridge, UK: Cambridge University Press.
- Nielsen, M. A., & Chuang, I. L. (2016). *Quantum Computation and Quantum Information* (10th anniversary edition ed.). New Delhi, India: Cambridge University Press.
- Ormrod, N., Vilasini, V., & Barrett, J. (2023). Which theories have a measurement problem? *arXiv preprint*. Retrieved from arXiv:2303.03353 [quant-ph]: <https://physics.paperswithcode.com/paper/which-theories-have-a-measurement-problem>
- Ozawa, M. (2003). Universally valid reformulation of the Heisenberg uncertainty principle on noise and disturbance in measurement. *Phys. Rev A*, 67, 1-6.
- Paz, J. P., & Zurek, W. H. (2000, Oct. 2). Environment–Induced Decoherence and the Transition from Quantum to Classical. *Course lectures given by both authors at the 72nd Les Houches Summer School on "Coherent Matter Waves", July–August 1999*. Retrieved from <https://arxiv.org/pdf/quant-ph/0010011.pdf>
- Qureshi, T. (2020, August 13). Demystifying the delayed-choice quantum eraser. *European Journal of Physics*, 41. doi:10.1088/1361-6404/ab923e

- Schlosshauer, M. (2019, October). Quantum decoherence. *Physics Reports*, 831, 1-57.
doi:<https://doi.org/10.1016/j.physrep.2019.10.001>
- Schrödinger, E. (1937). The present situation in quantum mechanics: A translation. *Proceedings of the American Philosophical Society*, 124, pp. 323-338.
- Storz, S., Schär, J., & Kulikov, et al. (2023). Loophole-free Bell inequality violation with superconducting circuits. *Nature*(617), 265–270. Retrieved from <https://doi.org/10.1038/s41586-023-05885-0>
- Thiago De Oliveira, & Caldeira, A. O. (2006, September). Coherence and Entanglement in a Stern-Gerlach experiment. *arXiv:quant-ph/0608192*. Retrieved from https://www.researchgate.net/publication/2199225_Coherence_and_Entanglement_in_a_Stern-Gerlach_experiment
- Warton, N., & Argaman, K. B. (2020, May). Colloquium: Bell's theorem and locally mediated reformulations of quantum mechanics. *Reviews of Modern Physics*, 92(2), 021002.
doi:<https://doi.org/10.1103/RevModPhys.92.021002>

Table of figures

Fig. 1 – Diagram of the experimental apparatus discussed here and in (Hobson, 2022). Reprinted from (Hobson,2022). Copyright © 2022 Art Hobson. Open access article distributed under the Creative Commons Attribution License.

Fig. 2 – Diagram of the delayed choice quantum eraser experiment. Original work by Patrick Edwin Moran – CC BY-SA 4.0,
<https://commons.wikimedia.org/wiki/File:KimDelayedChoiceQuantumEraserGraphsSVG.svg>

Fig. 3 – Results of (Kim, Yu, Kulik, Shih, & Marlan, 2000) relabeled for our Alice and Bob experiment. Relabeled version of original work by Patrick Edwin Moran - CC BY-SA 3.0,
<https://commons.wikimedia.org/w/index.php?curid=31312077>

Fig. 4 Two particles exchanging information. Copyright Smithsonian Institute. Copyright allows for educational, noncommercial use. https://forces.si.edu/atmosphere/02_01_02.html

Fig. 5 Schrödinger's cat. By Dhatfield - CC BY-SA 3.0,
<https://commons.wikimedia.org/w/index.php?curid=4279886>

Fig. 6 – A sequential Stern-Gerlach experiment. Our own work. Created with Lucidchart.

Fig. 7 – A diagram of an experiment used to show when entanglements are broken. Our own work. Created with Lucidchart.

