Possible Tetraquark Explanation for the X(6200)

Joseph Bevelacqua

Funding: The author(s) received no specific funding for this work.

Potential competing interests: The author(s) declared that no potential competing interests exist.

Abstract

The recently proposed X(6200) structure is investigated using a first-order tetraquark mass formula. This mass relationship is based on weakly bound J/ψ plus J/ψ meson clusters and provides a reasonable prediction (within about 1.5%) of the derived X(6200) mass. The spin-parity assignments are also consistent with the derived values.

1.0 Introduction

Recently, the LHCb Collaboration¹ reported pronounced structures in the invariant mass spectrum of J/ψ pairs produced in proton-proton collisions at the Large Hadron Collider. Dong et al.² argue that the data near 6.2 GeV can be described in terms of J/ψ plus J/ψ clusters having $J^\pi = 0^+$ or 2^+ . The possible state is denoted as the X(6200) in Ref. 2.

These results suggest the possibility of a tetraquark structure composed of two weakly interacting meson clusters. In addition to a pair of J/ψ ($c\bar{c}$) clusters, Ref. 2 suggests the possibility that the data can also be represented by a pair of η_c ($c\bar{c}$) mesons.

In this paper, the first-order tetraquark mass formulas of Refs. 3 - 9 are applied to evaluate the possible configuration of the X(6200). A guide to the tetraquark configuration is provided in Ref. 2 that suggests possible $J/\psi + J/\psi$ and $\eta_c + \eta_c$ meson clusters. These configurations are investigated assuming a $c\bar{c}c\bar{c}$ quark structure using the first-order tetraquark mass formula³⁻⁹.

2.0 Model and Formulation

Zel'dovich and Sakharov^{10,11} proposed a semiempirical mass formula that provides a prediction of mesons and baryons in terms of effective quark masses. Within this formulation, quark wave functions are assumed to reside in their lowest 1S state. These mass formulas are used as the basis for deriving a first-order tetraquark mass formula. In particular, the model proposed in this paper assumes the tetraquark is partitioned into two clusters with the interaction between the clusters providing a minimal contribution to the tetraquark mass.

The meson mass (M) formula of Refs. 3 and 4 is:

$$M_m = \delta_m + m_1 + m_2 + b_m [m_0^2 / (m_1 m_2)] \sigma_1 \cdot \sigma_2 \quad (1)$$

where m_1 (m_2) are the mass of the first (second) quark comprising the meson, m_0 is the average mass of a first generation quark^{10,11}, and the σ_i ($i = 1$ and 2) are the spin vectors for the quarks incorporated into the meson. The parameters δ_m and b_m are $40 \text{ MeV}/c^2$ and $615 \text{ MeV}/c^2$, respectively¹¹.

The last term in Eq. 1 represents the spin-spin interaction of the quarks and $\sigma_1 \cdot \sigma_2$ is the scalar product of the quark spin vectors. $\sigma_1 \cdot \sigma_2$ has the value $-3/4$ and $+1/4$ for pseudoscalar and vector mesons, respectively¹¹.

In formulating the tetraquark mass formula, effective quark masses provided by Griffiths¹² are utilized. These effective masses for d, u, s, c, b, and t quarks are 340, 336, 486, 1550, 4730, and 177000 MeV/c^2 , respectively. These masses are utilized in Eq. 1.

These six quarks are arranged in three generations: $[d(-1/3), u(+2/3)]$, $[s(-1/3), c(+2/3)]$, and $[b(-1/3), t(+2/3)]$ ¹³. The three generations are specified by the square brackets and the quark charges are given within parentheses.

3.0 First-Order Mass Formula for the X(6200)

The spin of a tetraquark within the first order mass formula is determined by coupling the two meson clusters

$$J^\pi = J^\pi(1) \times L \times J^\pi \quad (2)$$

where the first-order mass formula assumes a minimally interacting $L=0$ configuration^{3,4} between the meson clusters. The J/ψ has a $J^\pi = 1^-$ assignment¹³ while the η_c meson has a 0^- value. Using the coupling structure of Eq. 2 suggests that the $J/\psi + J/\psi$ tetraquark could have $J^\pi = 0^+, 1^+$, and 2^+ assignments. However, the $\eta_c + \eta_c$ configuration only allows a $J^\pi = 0^+$ assignment. The $J^\pi = 0^+$ or 2^+ assignment for the $X(6200)$ ² is consistent with the $J/\psi + J/\psi$ tetraquark $J^\pi = 0^+$ and 2^+ assignment using Eq. 2. In a similar manner, the $\eta_c + \eta_c$ $J^\pi = 0^+$ configuration is consistent with Ref. 2. The $J^\pi = 1^+$ assignment from the $J/\psi + J/\psi$ first-order tetraquark model is not consistent with Ref. 2.

The first-order mass formula used in this paper partitions the tetraquark into two meson clusters. For the $J/\psi + J/\psi$ configuration both clusters are vector mesons and the tetraquark mass formula is assumed to have the form^{3,4}

$$M = M_{vm} + M_{vm} + \Phi \quad (3)$$

where Φ defines the interaction between the $J/\psi + J/\psi$ meson clusters. Within the scope of this mass formula, the meson-meson cluster interaction is assumed to be weak and sufficiently small to be ignored. Accordingly, Eq. 3 represents a quasimolecular four quark system whose basic character is a weakly bound

$J/\psi + J/\psi$ system.

The $\eta_c + \eta_c J^\pi = 0^+$ configuration involves only scalar mesons with a tetraquark mass formula given by Eq. 4

$$M = M_{sm} + M_{sm} + \Phi \quad (4)$$

and Φ defines the interaction between the $\eta_c + \eta_c$ meson clusters.

Ref. 2 noted two possible configurations for the X(6200) tetraquark. Section 3.1 addresses the $J/\psi + J/\psi$ meson cluster and Section 3.2 described the $\eta_c + \eta_c$ configuration.

3.1 $J/\psi + J/\psi$ Meson Clusters

In the $J/\psi + J/\psi$ configuration, the meson cluster mass formula is provided by Eq. 3. The resulting first-order mass formula is $6294.6 \text{ MeV}/c^2$. This value is about 1.5% larger than the nominal $6200 \text{ MeV}/c^2$ value suggested in Ref. 2. The $J^\pi = 0^+$ or 2^+ assignment derived from Eq. 2 is also consistent with the X(6200)² prediction.

3.2 $\eta_c + \eta_c$ Meson Clusters

In the $\eta_c + \eta_c$ configuration, both clusters are scalar mesons and the tetraquark mass is derived from Eq. 4. The resulting first-order mass formula leads to a mass of $6236.1 \text{ MeV}/c^2$. This value is about 0.6% larger than the nominal $6200 \text{ MeV}/c^2$ value suggested in Ref. 2. The $J^\pi = 0^+$ assignment for $\eta_c + \eta_c$ derived from Eq. 2 is also consistent with the X(6200)² prediction.

4.0 Results and Discussion

The angular momentum coupling from Eq. 2 and the first-order mass formula of Eqs. 1, 3, and 4 are used to construct the X(6200) state. This state is modeled as noted in Sections 3.1 and 3.2.

The first-order mass formula provides a reasonable representation of the X(6200) state in terms of either $J/\psi + J/\psi$ or $\eta_c + \eta_c$ configurations. Both configurations are consistent with the predictions of Ref. 2 in terms of the tetraquark mass and J^π assignments. However, the first order mass formula is not sufficiently accurate to select the correct meson cluster configuration. It is possible that the physical state is a combination of these configurations.

Although these results are encouraging, they are based on a first-order mass formula with a number of uncertainties including the assumed quark masses¹² and the magnitude of the meson-meson cluster interaction. However, the model does provide an initial description of the X(6200) mass and its derived $J^\pi = 0^+$ or 2^+ assignment² in terms of a tetraquark structure .

5.0 Conclusions

The first-order mass formula predicts two possible tetraquark configurations for the X(6200). Both weakly bound $J/\psi + J/\psi$ and $\eta_c + \eta_c$ meson clusters lead to predicted mass values that are within about 1.5% of the derived value². The predicted J^π assignments for both configurations are also consistent with the derived values².

References

- 1) LHCb Collaboration, Sci. Bull. 65, 1983 (2020).
- 2) X.-K. Dong, V. Baru, F.-K. Guo, C. Hanhart, and A. Nefediev, Phys. Rev. Lett. **126**, 132001 (2021).
- 3) J. J. Bevelacqua, First-Order Tetraquark Mass Formula, Physics Essays **29**, 198 (2016).
- 4) J. J. Bevelacqua, Description of the X(5568) and Proposed 750 GeV/c² State in Terms of a First-Order Tetraquark Mass Formula, Physics Essays **29**, 367 (2016).
- 5) J. J. Bevelacqua, Fusion of doubly heavy mesons into a tetraquark, Physics Essays **31**, 167 (2018).
- 6) J. J. Bevelacqua, Possible Tetraquark Explanation for the Proposed X(3872), Physics Essays **32**, 469 (2019).
- 7) J. J. Bevelacqua, Description of the X(6900) as a Four Charmed Quark State in Terms of a First-Order Tetraquark Mass Formula, QEIOS **KLXLKJ**, 1 (2020).
<https://doi.org/10.32388/KLXLKJ>.
- 8) J. J. Bevelacqua, Description of the X(2900) as an Open Flavor Tetraquark in Terms of a First-Order Mass Formula, QEIOS, **OVLMEB**, 1 (2020).
<https://doi.org/10.32388/OVLMEB>.
- 9) J. J. Bevelacqua, Possible Tetraquark Explanation for the Proposed Z_{cs}(3985)[−], QEIOS **GLTEU2**, 1 (2021). <https://doi.org/10.32388/GLTEU2>.
- 10) Ya. B. Zel'dovich and A. D. Sakharov, *Kvarkovaia struktura i massy sil'novzaimodeistvuyushchikh chastits*, Yad. Fiz. **4**, 395 (1966).
- 11) A. D. Sakharov, Mass formula for mesons and baryons, Sov. Phys. JETP **51**, 1059 (1980).
- 12) D. Griffiths, *Introduction to Elementary Particles*, 2nd ed., (Wiley-VCH, Weinheim, 2008).
- 13) Particle Data Group, Review of Particle Physics, Prog. Theor. Exp. Phys. **2020**, 083C01 (2020).