

v1: 11 July 2024

Research Article

New Approximate Symmetry Theorems and Comparisons with Exact Symmetries

Preprinted: 4 March 2024

Peer-approved: 11 July 2024

© The Author(s) 2024. This is an Open Access article under the CC BY 4.0 license.

Qeios, Vol. 6 (2024)
ISSN: 2632-3834

Mehmet Pakdemirli¹

1. Mechanical Engineering Department, Celal Bayar University, Turkey

Three new approximate symmetry theories are proposed. The approximate symmetries are contrasted with each other and with the exact symmetries. The theories are applied to nonlinear ordinary differential equations for which exact solutions are available. It is shown that from the symmetries, approximate solutions as well as exact solutions in some restricted cases can be retrieved. Depending on the specific approximate theory and the equations considered, the approximate symmetries may expand the Lie Algebra of the exact symmetries, may be a perturbed form of the exact symmetries, or may be a subalgebra of the exact symmetries. Exact and approximate solutions are retrieved using the symmetries.

Corresponding author: Mehmet Pakdemirli, pakdemirli@gmail.com

1. Introduction

Lie Group theory [1][2][3] is a systematized and unified approach in search of analytical solutions of differential equations. It is a generalized approach for finding solutions of especially nonlinear differential equations and has the capability of producing results obtained by other ad-hoc methods. The perturbation method [4] is another powerful technique employed in search of approximate symmetries for over a century. Attempts to combine these powerful techniques appeared in the literature. In the case of perturbed equations, depending on the specific equation, the exact symmetries may not be sufficient to extract enough solutions. To extend the Lie Algebra and to construct further solutions, many approximate symmetry theories were proposed.

There are three main theories of approximate symmetries and a number of variants of these methods. The first method (Method I) is due to Baikov et al. [5][6] in which the symmetry generator is expanded in a perturbation series without expanding the dependent variable. On the contrary, in the second method due to Fushchich and Shtelen [7] (Method II), the dependent variable is expanded in a perturbation series, and the equations form a coupled system when separated with respect to orders. The approximate symmetry is then defined to be the exact symmetry of these coupled systems. In this method, since the number of dependent variables increases, the algebra for determining symmetries becomes rather involved. By assuming a linear unperturbed part and a nonlinear perturbed part for the differential equations, the hierarchical equations appearing in a separated block can be viewed as a linear non-homogeneous equation with a known function for the non-homogeneous part. This assumption drastically reduces the algebra, and the approximate symmetries of the nonlinear perturbed equation correspond to the exact symmetries of the linear non-homogeneous equation [8][9] (Method III). The three methods were contrasted with each other, and the advantages and disadvantages were outlined by applying the methods to the potential Burgers equation [8], creeping flow equations of a second-grade fluid [8] and an ordinary differential equation with quadratic nonlinearity [9]. A more theoretical basis for the comparisons of Method I and Method II was later presented [10].

Many papers have appeared in the literature applying the three methods to differential equations arising from mathematical physics. While a complete list of all work on the applications of the symmetry methods is beyond the scope of this study, a partial list will be given for the applications: Method I is applied in references [11][12][13][14][15][16][17][18][19], Method II in references [19][20][21][22][23][24][25][26][27][28][29][30][31][32] and Method III in references [33][34][35]. A Matlab package [36] was

developed to symbolically compute approximate symmetries for all three methods. Noetherian symmetries are another alternative to the conventional Lie Group symmetries, which involve Lagrangians. Approximate Noether symmetries were also calculated for mathematical physics models [37][38][39][40][41][42]. Exterior calculus is the other alternative to the classical Lie Group methods for calculating symmetries. The pioneering work on the topic is due to Harrison and Estabrook [43] and was later employed by others [44][45][46][47]. The approximate symmetry version of the exterior calculus approach was also presented [48][49]. The Approximate Homotopy Symmetry method is another approach developed in search of approximate symmetries [50][51][52].

In this work, three new approximate symmetry definitions are given for the first time. The exact symmetries and the approximate symmetries by the new three methods are contrasted with each other for sample ordinary differential equations whose exact solutions are known. Exact and approximate group invariant solutions are derived using the symmetries of each method. The new methods may extend the Lie Algebra, may be perturbed expansions of the exact symmetries, or may be a subgroup of the exact symmetries depending on the method used and the specific equation considered. The approximate symmetries are capable of retrieving approximate solutions as well as exact solutions.

2. Approximate Symmetry Theories

Three new definitions for approximate symmetries will be given in this section for the first time. The definitions have some differences from each other, which leads to different symmetry generators. To distinguish them from the Approximate Symmetry Theorems I-II and III discussed in the introduction, the new ones are numbered as IV-V and VI.

Approximate Symmetry Definition IV

For the k 'th order perturbed nonlinear ordinary differential equation

$$F(x, y, y', y'', \dots, y^{(k)}, \varepsilon) = 0 \quad (2.1)$$

with ε being the perturbation parameter and the Lie Group transformation parameter, the first order approximate symmetry corresponds to

$$F|_{\varepsilon=0} + \varepsilon X F|_{\varepsilon=0} = 0 \quad (2.2)$$

where

$$X = \xi(x, y) \frac{\partial}{\partial x} + \eta(x, y) \frac{\partial}{\partial y} + \mu \frac{\partial}{\partial \varepsilon} + \eta^1 \frac{\partial}{\partial y_1} + \dots + \eta^k \frac{\partial}{\partial y_k} + \mu \frac{\partial}{\partial \varepsilon} \quad (2.3)$$

is the approximate symmetry generator extended to k 'th order with the group transformations

$$\begin{aligned} x^* &= x + \varepsilon \xi(x, y, \varepsilon) \\ y^* &= y + \varepsilon \eta(x, y, \varepsilon) \\ y_1^* &= y_1 + \varepsilon \eta^1(x, y, y_1, \varepsilon) \\ &\vdots \\ y_k^* &= y_k + \varepsilon \eta^k(x, y, y_1, \dots, y_k, \varepsilon) \\ \mu^* &= \varepsilon \mu \end{aligned} \quad (2.4)$$

where

$$y_k = y^{(k)}, \eta^k = \frac{D\eta^{k-1}}{Dx} - y_k \frac{D\xi}{Dx}, \frac{D}{Dx} = \frac{\partial}{\partial x} + y_1 \frac{\partial}{\partial y} + y_2 \frac{\partial}{\partial y_1} + \dots + y_{k+1} \frac{\partial}{\partial y_k} \quad \square \quad (2.5)$$

Note that in determining the approximate symmetry generator, the whole block of (2.2) is used. In the case of exact symmetries, equation (2.2) separates into two equations, and the Lie Group transformation parameter is different from the perturbation parameter.

A slightly different definition is suggested below as the Symmetry Definition V.

Approximate Symmetry Definition V

For the k 'th order perturbed nonlinear ordinary differential equation

$$F(x, y, y', y'', \dots, y^{(k)}, \varepsilon) = 0 \quad (2.6)$$

with ε being the perturbation parameter and the Lie Group transformation parameter, the first order approximate symmetry corresponds to

$$X F|_{\varepsilon=0} = 0 \text{ when } F|_{\varepsilon=0} = 0 \quad (2.7)$$

where

$$X = \xi(x, y) \frac{\partial}{\partial x} + \eta(x, y) \frac{\partial}{\partial y} + \mu \frac{\partial}{\partial \varepsilon} + \eta^1 \frac{\partial}{\partial y_1} + \dots + \eta^k \frac{\partial}{\partial y_k} + \mu \frac{\partial}{\partial \varepsilon} \quad (2.8)$$

is the approximate symmetry generator extended to k 'th order with the group transformations

$$\begin{aligned} x^* &= x + \varepsilon \xi(x, y) \\ y^* &= y + \varepsilon \eta(x, y) \\ y_1^* &= y_1 + \varepsilon \eta^1(x, y, y_1) \\ &\vdots \\ y_k^* &= y_k + \varepsilon \eta^k(x, y, y_1, \dots, y_k) \\ \mu^* &= \varepsilon \mu \end{aligned} \quad (2.9)$$

where

$$y_k = y^{(k)}, \eta^k = \frac{D\eta^{k-1}}{Dx} - y_k \frac{D\xi}{Dx}, \frac{D}{Dx} = \frac{\partial}{\partial x} + y_1 \frac{\partial}{\partial y} + y_2 \frac{\partial}{\partial y_1} + \dots + y_{k+1} \frac{\partial}{\partial y_k} \quad (2.10)$$

In the above version, the block, i.e., Eq. (2.2), is separated into two parts. It is still different from the exact symmetry definition, since the Lie Group transformation parameter is different from the perturbation parameter in the exact symmetry case. Also, in the exact case $F = 0$, whereas in this definition, the unperturbed equation satisfies the condition $F|_{\varepsilon=0} = 0$ which is merely an approximation of the original equation, namely the unperturbed equation itself. Note also that the infinitesimals $\xi(x, y)$ and $\eta(x, y)$ do not contain the perturbation parameter as an argument, while this is not the case for Approximate Symmetry Method IV.

A variant of the fourth definition may also be proposed where the Lie Group parameter is not the perturbation parameter.

Approximate Symmetry Definition VI

For the k 'th order perturbed nonlinear ordinary differential equation

$$F(x, y, y', y'', \dots, y^{(k)}, \varepsilon) = 0 \quad (2.11)$$

with ε being the perturbation parameter and α being the Lie Group transformation parameter, the first order approximate symmetry corresponds to

$$F|_{\alpha=0} + \alpha X F|_{\alpha=0} = 0 \quad (2.12)$$

where

$$X = \xi(x, y) \frac{\partial}{\partial x} + \eta(x, y) \frac{\partial}{\partial y} + \eta^1 \frac{\partial}{\partial y_1} + \dots + \eta^k \frac{\partial}{\partial y_k} \quad (2.13)$$

is the approximate symmetry generator extended to k 'th order with the group transformations

$$\begin{aligned} x^* &= x + \alpha \xi(x, y, \varepsilon) \\ y^* &= y + \alpha \eta(x, y, \varepsilon) \\ y_1^* &= y_1 + \alpha \eta^1(x, y, y_1, \varepsilon) \\ &\vdots \\ y_k^* &= y_k + \alpha \eta^k(x, y, y_1, \dots, y_k, \varepsilon) \end{aligned} \quad (2.14)$$

where

$$y_k = y^{(k)}, \eta^k = \frac{D\eta^{k-1}}{Dx} - y_k \frac{D\xi}{Dx}, \frac{D}{Dx} = \frac{\partial}{\partial x} + y_1 \frac{\partial}{\partial y} + y_2 \frac{\partial}{\partial y_1} + \dots + y_{k+1} \frac{\partial}{\partial y_k} \quad (2.15)$$

If the two terms in (2.12) are separated, then one obtains the exact symmetries. The idea here is not to separate the block in search of approximate symmetries. This definition indeed is not an approximate symmetry definition in the sense that it does not extend the Lie Algebra of the exact symmetries, rather produces a subgroup of the exact symmetries. It is included for comparison reasons and for outlining the importance of selecting the perturbation parameter as the Lie Group parameter as was done in definitions V and VI.

3. Approximate Symmetry Calculations

For a number of ordinary differential equations, symmetries corresponding to the three methods are calculated together with the exact symmetries (Table 1).

Equation	Exact Symmetry	Approximate Symmetry IV	Approximate Symmetry V	Approximate Symmetry VI
$y' + \varepsilon y = 0$	Unsolvable $\begin{aligned} &\eta_x \\ &+ \varepsilon(\eta - y\eta_y \\ &+ y\xi_x) - \varepsilon^2\xi_y y^2 \\ &= 0 \end{aligned}$	$\begin{aligned} \xi &= -\frac{\mu}{2}x^2 + a_1x \\ &+ a_2 \\ \eta &= \left(-\mu x + a_1 \right. \\ &\left. - \frac{1}{\varepsilon}\right)y + b \end{aligned}$	$\begin{aligned} \xi &= \xi(x, y) \\ \eta &= -\mu xy + a(y) \end{aligned}$	$\begin{aligned} \xi &= a + be^{-\varepsilon x} \\ \eta &= -(\varepsilon be^{-\varepsilon x} \\ &+ \frac{1}{\alpha})y + ce^{-\varepsilon x} \end{aligned}$
$y' + e^{\varepsilon y} = 0$	Unsolvable $\begin{aligned} &\eta_x \\ &- e^{\varepsilon y}(\eta_y - \xi_x \\ &- \varepsilon\eta) - \xi_y e^{2\varepsilon y} \\ &= 0 \end{aligned}$	$\begin{aligned} \xi &= -\frac{\mu}{2}x^2 + a_1x \\ &+ a_2 \\ \eta &= \left(-\mu x + a_1 \right. \\ &\left. - \frac{1}{\varepsilon}\right)y \\ &- \frac{1}{\varepsilon}x + b_1 \end{aligned}$	$\begin{aligned} \xi &= \xi(x, y) \\ \eta &= -\xi + \mu \frac{y^2}{2} \\ &+ a(x + y) \end{aligned}$	$\begin{aligned} \xi &= \frac{1}{\alpha}x + b \\ \eta &= -\frac{1}{\varepsilon\alpha} \end{aligned}$
$y'' + \varepsilon y'^2 = 0$	$\begin{aligned} \xi &= (ax + b)e^{\varepsilon y} \\ &+ cx^2 \\ &+ dx + e \\ \eta &= \left(fe^{-\varepsilon y} + \frac{c}{\varepsilon}\right)x \\ &+ ge^{-\varepsilon y} + h \\ &+ \frac{a}{\varepsilon}e^{\varepsilon y} \end{aligned}$	$\begin{aligned} \xi &= ax + b \\ \eta &= \left(2a - \frac{1}{\varepsilon}\right)y \\ &+ cx + d \end{aligned}$	$\begin{aligned} \xi &= (a_2x + a_3)y \\ &+ c_1x^2 + b_1x + b_2 \\ \eta &= (2a_2 - \mu)\frac{y^2}{2} \\ &+ (c_1x + c_2)y + d_1x \\ &+ d_2 \end{aligned}$	$\begin{aligned} \xi &= \frac{1}{2\alpha}x + a \\ \eta &= b \end{aligned}$
$y'' - 2\varepsilon yy' = 0$	$\begin{aligned} \xi &= ax + b \\ \eta &= -ay \end{aligned}$	$\begin{aligned} \xi &= a_2x + a_3 \\ \eta &= \left(2a_2 - \frac{1}{\varepsilon}\right)y \\ &+ b_1x + b_2 \end{aligned}$	$\begin{aligned} \xi &= \left(\frac{\mu}{3}x^2 + a_2x \right. \\ &\left. + a_3\right)y \\ &+ c_1x^2 + b_1x + b_2 \\ \eta &= \left(\frac{2\mu}{3}x + a_2\right)y^2 \\ &+ (c_1x + c_2)y \\ &+ d_1x + d_2 \end{aligned}$	$\begin{aligned} \xi &= ax + b \\ \eta &= -ay \end{aligned}$
$y'' - y + \varepsilon y^2 = 0$	$\begin{aligned} \xi &= a \\ \eta &= 0 \end{aligned}$	$\begin{aligned} \xi &= a \\ \eta &= -\frac{1}{\varepsilon}y + b_1e^x \\ &+ b_2e^{-x} \end{aligned}$	$\begin{aligned} \xi &= (a_1e^x + a_2e^{-x})y \\ &+ b_1 + b_2e^{2x} \\ &+ b_3e^{-2x} \\ \eta &= (a_1e^x - a_2e^{-x})y^2 \\ &+ (c_1 + b_2e^{2x} \\ &- b_3e^{-2x})y \\ &+ d_1e^x + d_2e^{-x} \end{aligned}$	$\begin{aligned} \xi &= 0 \\ \eta &= 0 \end{aligned}$

Equation	Exact Symmetry	Approximate Symmetry IV	Approximate Symmetry V	Approximate Symmetry VI
$y''' = \varepsilon f(y', y'')$	$\xi = a$ $\eta = b$	$\xi = a_1 x + a_2$ $\eta = \left(3a_1 - \frac{1}{\varepsilon}\right) y$ $+ b_1 x^2 + b_2 x + b_3$	$\xi = a_1 x^2 + a_2 x + a_3$ $\eta = (2a_1 x + a_2 + c) y$ $+ b_1 x^2 + b_2 x + b_3$	$\xi = 0$ $\eta = 0$

Table 1. Exact and Approximate Symmetries

From the symmetries, for the specific problems considered, some conclusions can be given:

For first-order equations:

- In the case of exact symmetries, usually the determining equation for the infinitesimals cannot be separated and remains unsolvable, unless some further simplifying assumptions are made.
- On the contrary, the infinitesimals are solvable for the approximate symmetries.
- Among the symmetries, the richest symmetry corresponds to the approximate symmetry V case for first-order equations.

For the higher-order equations:

- For the equation $y'' + \varepsilon y'^2 = 0$, while the exact and approximate symmetry V possess 8-parameter Lie Group transformations, the other symmetries possess fewer parameters.
- For the equation $y'' + \varepsilon y'^2 = 0$, if the exact symmetry is expanded in a Taylor series up to $O(\varepsilon)$, the approximate symmetry V result can be retrieved.
- For the last 3 equations, approximate symmetries IV and V are richer than the exact symmetries. For the equation $y'' - y + \varepsilon y^2 = 0$, while the exact symmetries are one-parameter, the approximate symmetry IV contains 3 parameters, and the approximate symmetry V contains 8-parameter Lie Group transformations.
- As a general rule, approximate symmetry VI is a subalgebra of exact symmetries if not equal.
- As a general rule, approximate symmetry V produces the richest symmetries among the approximate ones.

4. Solutions

Using the symmetries, group-invariant solutions are constructed for the four problems and listed in Table 2. In the table, the exact and one-correction-term approximate solutions of the problem are given first, and the specific symmetries to retrieve the results are given. The equation to be solved is

$$\frac{dx}{\xi(x, y)} = \frac{dy}{\eta(x, y)}. \quad (4.1)$$

Substituting the outcome into the original equation to satisfy it and then applying the initial conditions, the approximate and exact solutions are obtained.

Equation	Exact and Approximate Solutions	Exact Symmetry	Approximate Symmetry IV	Approximate Symmetry V	Approximate Symmetry VI
$y' + \varepsilon y = 0$ $y(0) = 1$	$y_e = e^{-\varepsilon x}$	Retrievable	$\xi = a_2$ $\eta = -\frac{1}{\varepsilon}y$	$\xi = b$ $\eta = y$	$\xi = be^{-\varepsilon x}$ $\eta = -\varepsilon be^{-\varepsilon x}y$
	$y_a = 1 - \varepsilon x$	Not directly retrievable	$\xi = a_2$ $\eta = b$	$\xi = 1$ $\eta = a$	Not directly retrievable
$y' + e^\varepsilon y = 0$ $y(0) = 0$	$y_e = -\frac{1}{\varepsilon} \ln(1 + \varepsilon x)$	Not directly retrievable	Not directly retrievable	Not directly retrievable	$\xi = \frac{1}{\alpha}x + b$ $\eta = -\frac{1}{\varepsilon\alpha}$
	$y_a = -x + \varepsilon \frac{x^2}{2}$	Not directly retrievable	$\xi = a_2$ $\eta = -\frac{1}{\varepsilon}x + b_1$	Not directly retrievable	Not directly retrievable
$y'' + \varepsilon y'^2 = 0$ $y(0) = 0$ $y'(0) = 1$	$y_e = \frac{1}{\varepsilon} \ln(1 + \varepsilon x)$	$\xi = dx + e$ $\eta = h$	$\xi = ax + b$ $\eta = d$	$\xi = b_1x + b_2$ $\eta = d_2$	$\xi = \frac{1}{2\alpha}x + a$ $\eta = b$
	$y_a = x - \varepsilon \frac{x^2}{2}$	$\xi = e$ $\eta = \frac{c}{\varepsilon}x + h$	$\xi = b$ $\eta = cx + d$	$\xi = b_2$ $\eta = d_1x + d_2$	Not directly retrievable
$y'' - 2\varepsilon yy' = 0$ $y(0) = 1$ $y'(0) = \varepsilon$	$y_e = \frac{1}{1 - \varepsilon x}$	$\xi = ax + b$ $\eta = -ay$	$\xi = a_2x + a_3$ $\eta = \left(2a_2 - \frac{1}{\varepsilon}\right)y$	$\xi = b_1x + b_2$ $\eta = c_2y$	$\xi = ax + b$ $\eta = -ay$
	$y_e = 1 + \varepsilon x$	Not directly retrievable	$\xi = a_3$ $\eta = b_2$	$\xi = b_2$ $\eta = d_2$	Not directly retrievable

Table 2. Group Invariant Solutions

Regarding the retrieval of solutions, approximate symmetries IV and V perform better than approximate symmetry VI in most cases. Approximate symmetry VI cannot produce approximate solutions for all the problems considered since it produces a subgroup of the exact symmetries. In most cases, the approximate symmetries also lead to the exact solutions. This is because the dependent variable is not expanded in a perturbation series, a feature observed in Approximate Symmetry I theory due to Baikov et. al. [5][6] also, which can be questioned from the perturbation theory point of view [8]. In contrast to this similarity, the main difference between the mentioned Method I [5][6] and the approximate symmetry theories presented here is that the generator is expanded in a perturbation series in the former case, while it is not expanded in a series in the ones presented here.

5. Concluding Remarks

Based on this study and the previous work on approximate symmetry theories [5][6][7][8][9][10], the following conclusions can be made

- If the goal is to produce only the approximate solutions, Method II [7] and Method III with less algebra [8] are recommended, since those methods are more consistent with perturbation theory and directly lead to the approximate solutions.
- If the goal is to produce both the approximate and exact solutions, Method I [5][6] and Methods IV and V presented in this study can be employed.
- Among the new three approximate methods, Method V is recommended for second and higher order equations most, since it leads to richer symmetries.
- For first-order differential equations, however, Method IV leads to simpler and solvable symmetry infinitesimals than those of exact symmetry and Method V cases.
- Method VI corresponds to the subgroup of the exact symmetries which leads to the group invariant solutions.
- The work can be extended directly to include partial differential equations. A comparison of the symmetries and solutions for partial differential equations is a further research topic in the future.

Statements and Declarations

Data Availability

Data sharing is not applicable to this article as no new data were created or analyzed in this study. All results are derived from the mathematical analysis presented within the text.

Author Contributions

MP: Conceptualization, Methodology, Formal analysis, Investigation, Writing – original draft, Writing – review & editing.

References

1. [△]Bluman GW, Kumei S (1989). *Symmetries and Differential Equations*. New York: Springer Verlag.
2. [△]Stephani H (1989). *Differential Equations: Their Solution Using Symmetries*. New York: Cambridge University Press.
3. [△]Ibragimov NH (1994). *CRC Handbook of Lie Group Analysis of Differential Equations, Volume 1*. Boca Raton: CRC Press.
4. [△]Nayfeh AH (1981). *Introduction to Perturbation Techniques*. New York: John Wiley and Sons.
5. ^{a, b, c, d, e}Baikov VA, Gazizov RK, Ibragimov NH (1988). "Approximate symmetries." *Matematicheskii Sbornik*. **136**:435–450. doi:10.1070/SM1989v064n02ABEH003388.
6. ^{a, b, c, d, e}Baikov VA, Gazizov RK, Ibragimov NH (1996). "Approximate transformation groups and deformations of symmetry Lie algebras." In: Ibragimov NH, editor. *CRC Handbook of Lie Group Analysis of Differential Equations*. Vol 3. Boca Raton, Florida: CRC Press. Chapter 2.
7. ^{a, b, c}Fushchich WI, Shtelen WH (1989). "On approximate symmetry and approximate solutions of the non-linear wave equation with a small parameter." *Journal of Physics A: Mathematical and General*. **22**:887–890. doi:10.1088/0305-4470/22/18/007.
8. ^{a, b, c, d, e}Pakdemirli M, Yürüsoy M, Dolapçı İT (2004). "Comparison of approximate symmetry methods for differential equations." *Acta Applicandae Mathematica*. **80**(3):243–271. doi:10.1023/B:ACAP000018792.87732.25.
9. ^{a, b, c}Pakdemirli M, Yürüsoy M (2000). "On approximate symmetries of a wave equation with quadratic non-linearity." *Mathematical and Computational Applications*. **5**(3):179–184. doi:10.3390/mca5020179.
10. ^{a, b}Wiltshire R (2006). "Two approaches to the calculation of approximate symmetry exemplified using a system of advection–diffusion equations." *Journal of Computational and Applied Mathematics*. **197**:287–301. doi:10.1016/j.cam.2005.11.003.
11. [△]Nadjafikhah M, Mokhtary A (2013). "Approximate Hamiltonian symmetry groups and recursion operators for perturbed evaluation equations." *Advances in Mathematical Physics*. doi:10.1155/2013/56863

12. ^ΔGrebenev VN, Oberlack M (2007). "Approximate Lie Symmetries of the Navier-Stokes equations." *Journal of Nonlinear Mathematical Physics*. 14(2):157–163. doi:[10.2991/jnmp.2007.14.2.1](https://doi.org/10.2991/jnmp.2007.14.2.1).
13. ^ΔKara AH, Mahomed FM, Qadir A (2008). "Approximate symmetries and conservation laws of the geodesic equations for the Schwarzschild metric." *Nonlinear Dynamics*. 51:183–188. doi:[10.1007/s11071-007-9201-x](https://doi.org/10.1007/s11071-007-9201-x).
14. ^ΔNadjafikhah M, Kabi-Nejad P (2013). "Approximate Symmetries of the Harry Dym equation." *ISRN Mathematical Physics*. doi:[10.1155/2013/109170](https://doi.org/10.1155/2013/109170).
15. ^ΔRahimian M, Tomanian M, Nadjafikhah M (2017). "Approximate Symmetry and exact solutions of the singularly perturbed Boussinesq equation." *Communications in Nonlinear Science and Numerical Simulations*. 53:1–9. doi:[10.1016/j.cnsns.2017.04.033](https://doi.org/10.1016/j.cnsns.2017.04.033).
16. ^ΔBokhari AH, Kara AH, Zaman FD (2007). "Exact solutions of some general nonlinear wave equations in elasticity." *Nonlinear Dynamics*. 48:49–54. doi:[10.1007/s11071-006-9050-z](https://doi.org/10.1007/s11071-006-9050-z).
17. ^ΔAhmed WA, Zaman FD, Saleh K (2017). "Invariant solutions for a class of perturbed nonlinear wave equations." *Mathematics*. 5:59. doi:[10.3390/math5040059](https://doi.org/10.3390/math5040059).
18. ^ΔHussain I, Mahomed FM, Qadir A (2007). "Second order approximate symmetries of the geodesic equations for the Reissner-Nordstrom metric and re-scaling of energy of a test particle." *Symmetry, Integrability and Geometry: Methods and Applications*. 3:115. doi:[10.3842/SIGMA.2007.115](https://doi.org/10.3842/SIGMA.2007.115).
19. ^a ^b Mahdavi A, Nadjafikhah M (2015). "Two approaches to the calculation of approximate symmetry of Ostrovsky equation with small parameter." *Mathematical Physics and Analytical Geometry*. 18:3. doi:[10.1007/s11040-015-9170-0](https://doi.org/10.1007/s11040-015-9170-0).
20. ^ΔEuler N, Shulga MW, Steeb WH (1992). "Approximate symmetries and approximate solutions for a multi-dimensional Landau-Ginzburg equation." *Journal of Physics A: Mathematical and General*. 25:1095–1103. doi:[10.1088/0305-4470/25/18/002](https://doi.org/10.1088/0305-4470/25/18/002).
21. ^ΔEuler M, Euler N, Köhler A (1994). "On the construction of approximate solutions for a multi-dimensional nonlinear heat equation." *Journal of Physics A: Mathematical and General*. 27:2083–2092. doi:[10.1088/0305-4470/27/6/031](https://doi.org/10.1088/0305-4470/27/6/031).
22. ^ΔEuler N, Euler M (1994). "Symmetry properties of the approximations of multidimensional generalized Van der Pol equations." *Nonlinear Mathematical Physics*. 1:41–59. doi:[10.2991/jnmp.1994.1.1.3](https://doi.org/10.2991/jnmp.1994.1.1.3).
23. ^ΔSalvo RD, Gorgone M, Oliveri F (2018). "A consistent approach to approximate Lie symmetries of differential equations." *Nonlinear Dynamics*. 91:371–386. doi:[10.1007/s11071-017-3875-5](https://doi.org/10.1007/s11071-017-3875-5).
24. ^ΔZhang ZY, Zhang WM, Chen YF (2016). "A new method to find series solutions of a nonlinear wave equation." *Applied Mathematics Letters*. 57:20–24. doi:[10.1016/j.aml.2015.12.017](https://doi.org/10.1016/j.aml.2015.12.017).
25. ^ΔDiatla B, Soh CW, Khaliq CM (2008). "Approximate symmetries and solutions of the hyperbolic heat equation." *Applied Mathematics and Computation*. 205:263–272. doi:[10.1016/j.amc.2008.06.060](https://doi.org/10.1016/j.amc.2008.06.060).
26. ^ΔJiao X, Yao R, Zhang S, Lov SY (2009). "Approximate symmetry reduction approach: Infinite series reductions to the KdV-Burgers equation." *Zeitschrift für Naturforschung*. 64a:676–684. doi:[10.1515/zna-2009-1102](https://doi.org/10.1515/zna-2009-1102).
27. ^ΔZhang ZY, Yong XL, Chen YF (2012). "Classification and approximate solutions to perturbed diffusion-convection equations." *Applied Mathematics and Computation*. 219:1120–1124. doi:[10.1016/j.amc.2012.07.019](https://doi.org/10.1016/j.amc.2012.07.019).
28. ^ΔIbragimov R, Jefferson G, Carminati J (2013). "Explicit invariant solutions associated with nonlinear atmospheric flows in a thin rotating spherical shell with and without west-to-east jets perturbations." *Annals of Mathematical Physics*. 3:375–391. doi:[10.1007/s13324-013-0062-9](https://doi.org/10.1007/s13324-013-0062-9).
29. ^ΔIbragimov R, Jefferson G, Carminati J (2013). "Invariant and approximately invariant solutions of nonlinear internal gravity waves forming a column of stratified fluid affected by the Earth rotation." *International Journal of Nonlinear Mechanics*. 51:28–44. doi:[10.1016/j.ijnonlinmec.2012.12.001](https://doi.org/10.1016/j.ijnonlinmec.2012.12.001).
30. ^ΔGrebenev VN, Oberlack M, Grishkov AN (2008). "Lie Algebra methods for the applications to the statistical theory of turbulence." *Journal of Nonlinear Mathematical Physics*. 15(2):227–251. doi:[10.2991/jnmp.2008.15.2.9](https://doi.org/10.2991/jnmp.2008.15.2.9).
31. ^ΔJefferson GF (2013). "On the second order approximate symmetry classification and optimal systems of subalgebras for a forced Korteweg de Vries equation." *Communications in Nonlinear Science and Numerical Simulation*. 18:2340–2358. doi:[10.1016/j.cnsns.2012.12.022](https://doi.org/10.1016/j.cnsns.2012.12.022).
32. ^ΔRuggieri M, Speciale MP (2019). "Optimal system and new approximate solutions of a generalized Ames's equation." *Symmetry*. 11(10):1230. doi:[10.3390/sym11101230](https://doi.org/10.3390/sym11101230).
33. ^ΔDolapci IT, Pakdemirli M (2004). "Approximate symmetries of creeping flow equations of a second-grade fluid." *International Journal of Non-Linear Mechanics*. 39:1603–1619. doi:[10.1016/j.ijnonlinmec.2004.05.001](https://doi.org/10.1016/j.ijnonlinmec.2004.05.001).

34. ^ΔPakdemirli M, Şahin AZ (2005). "Approximate symmetries of hyperbolic heat conduction equation with temperature dependent thermal properties." *Mathematical and Computational Applications*. 10 (1):139–145. doi:[10.3390/mca10010139](https://doi.org/10.3390/mca10010139).
35. ^ΔNadjafikhah M, Mokhtary A (2014). "Symmetry analysis of Black-Sholes equation for small values of volatility and rate of return." *Journal of Interpolation and Approximation in Scientific Computing*. 1–10. doi:[10.105899/2014/jiasc-00054](https://doi.org/10.105899/2014/jiasc-00054).
36. ^ΔJeffersen GF, Carminati J (2013). "ASP: Automated symbolic computation of approximate symmetries of differential equations." *Computer Physics Communications*. 184(3):1045–1063. doi:[10.1016/j.cpc.2012.11.012](https://doi.org/10.1016/j.cpc.2012.11.012).
37. ^ΔJamal S, Mnguni N (2018). "Approximate conditions admitted by classes of the Lagrangian $L=1/2 (-u \wedge (\Box^2 + u^2) + e^{\Lambda i} G_i(u, u^{\Lambda \Box}, u^{\Lambda \Box \Box})$." *Applied Mathematics and Computation*. 335:65–74. doi:[10.1016/j.amc.2018.04.020](https://doi.org/10.1016/j.amc.2018.04.020).
38. ^ΔJamal S (2019). "Approximate conservation laws of nonvariational differential equations." *Mathematics*. 7:574. doi:[10.3390/math7070574](https://doi.org/10.3390/math7070574).
39. ^ΔNaeem I, Mahomed FM (2010). "Approximate first integrals for a system of two coupled Van der Pol oscillators with linear diffusive coupling." *Mathematical and Computational Applications*. 15(4):720–731. doi:[10.3390/mca15040720](https://doi.org/10.3390/mca15040720).
40. ^ΔHejazi SR, Husseinpour S, Lashkarian E (2019). "Approximate symmetries, conservation laws and numerical solutions for a class of perturbed linear wave type system." *Questiones Mathematicae*. 42(10):1393–1409. doi:[10.2989/16073606.2018.1538062](https://doi.org/10.2989/16073606.2018.1538062).
41. ^ΔJamal S (2018). "n'th order approximate Lagrangians induced by perturbative geometries." *Mathematical Physics and Analytical Geometry*. 21:25. doi:[10.1007/s11040-018-9283-3](https://doi.org/10.1007/s11040-018-9283-3).
42. ^ΔJamal S (2019). "Perturbative manifolds and the Noether generators of n'th order Poisson equations." *Journal of Differential Equations*. 266:4018–4026. doi:[10.1016/j.jde.2018.09.025](https://doi.org/10.1016/j.jde.2018.09.025).
43. ^ΔHarrison BK, Estabrook FB (1971). "Geometric approach to invariance groups and solution of partial differential systems." *Journal of Mathematical Physics*. 12(4):653. doi:[10.1063/1.1665631](https://doi.org/10.1063/1.1665631).
44. ^ΔŞuhubi ES (1991). "Isovector fields and similarity solutions for general balance equations." *International Journal of Engineering Science*. 29(1):133–150. doi:[10.1016/0020-7225\(91\)90083-F](https://doi.org/10.1016/0020-7225(91)90083-F).
45. ^ΔPakdemirli M, Suhubi ES (1992). "Similarity solutions of boundary layer equations for second order fluids." *International Journal of Engineering Science*. 30(5):611–629. doi:[10.1016/0020-7225\(92\)90006-3](https://doi.org/10.1016/0020-7225(92)90006-3).
46. ^ΔPakdemirli M, Yürüşoy M (1999). "Equivalence transformations applied to exterior calculus approach for finding symmetries: An example of non-Newtonian fluid flow." *International Journal of Engineering Science*. 37:25–32. doi:[10.1016/S0020-7225\(98\)00028-7](https://doi.org/10.1016/S0020-7225(98)00028-7).
47. ^ΔŞuhubi ES (1991). "Group properties and similarity solutions for a quasi-linear wave equation in the plane." *International Journal of Non-Linear Mechanics*. 26(5):567–584. doi:[10.1016/0020-7462\(91\)90010-Q](https://doi.org/10.1016/0020-7462(91)90010-Q).
48. ^ΔDavison AH, Kara AH (2008). "Symmetries and differential forms." *Journal of Nonlinear Mathematical Physics*. 15:36–43. doi:[10.2991/jnmp.2008.15.s1.3](https://doi.org/10.2991/jnmp.2008.15.s1.3).
49. ^ΔPittman ER, Schmidt EM, Ramsey SD (2000). "Symmetries of the P3 approximation to the Boltzmann neutron transport equation." *Annals of Nuclear Energy*. 144. doi:[10.1016/j.anucene.2020.107502](https://doi.org/10.1016/j.anucene.2020.107502).
50. ^ΔJiao X, Zheng Y, Wu B (2012). "Approximate Homotopy symmetry and infinite series solutions to the perturbed mKdV equation." *Applied Mathematics and Computation*. 218(17):8486–8491. doi:[10.1016/j.amc.2012.02.008](https://doi.org/10.1016/j.amc.2012.02.008).
51. ^ΔJiao XY, Gao Y, Lov SY (2009). "Approximate Homotopy symmetry method: Homotopy series solutions to the sixth order Boussinesq equation." *Science in China Series G: Physics, Mechanics and Astronomy*. 52(8):1169–1178. doi:[10.1007/s11433-009-0181-3](https://doi.org/10.1007/s11433-009-0181-3).
52. ^ΔZhang Z, Chen Y (2011). "A comparative study of approximate symmetry and approximate homotopy symmetry to a class of perturbed nonlinear wave equations." *Nonlinear Analysis: Theory, Methods & Applications*. 74:4300–4318. doi:[10.1016/j.na.2011.03.005](https://doi.org/10.1016/j.na.2011.03.005).

Declarations

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.