

[Open Peer Review on Qeios](#)

# Obesity and life events: the hypothesis of psychological phenotypes

Amelia Rizzo<sup>1</sup>

<sup>1</sup> University of Messina

**Funding:** No specific funding was received for this work.

**Potential competing interests:** No potential competing interests to declare.

## Abstract

Psychological studies on obesity have focused on finding common traits in the population of people with obesity, such as personality traits. In contrast, in the field of medicine, it is pointed out that individuals with obesity can be very different from each other, e.g. in terms of their metabolic profile, which gives rise to different phenotypes.

This article proposes a clinical and theoretical observation of the role of life events in the development of obesity. Obesity patients indeed report differentiated life and weight control histories. The proposal is to distinguish these types of patients into two psychological phenotypes at initial assessment, as surgical and post-surgical outcomes may differ. While from a mental health point of view, identifying these elements may be an indicator to send the patient to effective and targeted psychological treatments.

**Keywords:** Obesity, Life events, Trauma, Phenotypes, Psychotherapy.

## Introduction

The World Health Organization (WHO) defines "obesity" as a condition characterized by an excessive presence of adipose tissue in the human body which induces a significant increase in risk for the state of health. The simple definition of obesity as an imbalance between the energy introduced into the body and the energy expended. This explanation actually tends to ignore the complexity of the pathology. Different factors interacting with each other are responsible for weight gain, so the etiology of obesity must be sought individually on the basis of complex genetic, environmental and psychological factors interactions.

Obesity is widely recognized as a risk factor for multiple diseases, such as hypertension, dyslipidemia, cardiovascular disease, type 2 diabetes, arthritis, some cancers and depression, and is associated with reduced life expectancy and quality of life, as well as with lower general mental health. However, the underlying causes of obesity are poorly understood, even among health professionals. Two surveys found that a significant number of primary care professionals

believed that the three most important causes of obesity were physical inactivity, overeating, and high-fat dieting (Farooqi & O'rahilly, 2007; Hill, Williams & Fruhbeck, 2009; Smith & Smith, 2016).

However, the 2007 Foresight report highlighted over 100 different biological, psychological, environmental and social factors that potentially contribute to obesity. These factors range from genetic predisposition or neurochemical imbalances to emotional aspects, such as boredom or comfort eating, and complex social pressures that can begin in childhood with the need to finish a plate of food, and continue throughout life and can make healthy choices difficult (McPherson, Marsh & Brown, 2007).

Regardless of the underlying cause, once weight gain occurs, the body resists any attempts at weight loss. A normal physiological response to weight gain would be a lowering of the metabolic rate and an increase in hunger to maintain the new status quo. Therefore, even when an individual is successful in losing weight, weight gain will occur in most people and lifelong treatment is required to maintain the new normal weight. For this reason, obesity requires ongoing management similar to that of many other long-term chronic conditions, but is accompanied by an unprecedented degree of stigmatization (Puhl & Heuer, 2010).

In our view, the first step in combating this stigma is to speak correctly, both among healthcare professionals, when confronting the general public, and in clinical encounters between a healthcare professional and a person with obesity, putting in light of the possible psychological factors involved (Di Pauli, 2016).

## The myth of obesity psychological patterns

The first formal attempts to establish a relationship between obesity and psychological aspects date back to the original psychoanalytic conceptualizations of Hilde Bruch (1948), up to the more recent relational-systemic and cognitive-behavioral perspectives. The aim of the Authors was to formulate psychogenetic hypotheses of obesity, theoretical models that could explain and predict the onset of obesity starting from previous psychological, psychosocial and dysfunctional family conditions (Slochower, 1987).

However, scientific reflection on the links between obesity and psychological factors has evolved considerably over the years, especially from an empirical and methodological point of view. The empirical studies of the first generation, in the wake of the psychogenetic considerations of obesity, tried to find similarities in the psychological, psychosocial and psychopathological profiles of people affected by obesity, comparing them with normal weight individuals. The main objective of these researches was precisely to explain the etiology of obesity by analyzing the causal contribution of psychological factors. However, these early empirical studies failed to demonstrate psychological homogeneity among obese people (see "The psychogenic theory of obesity" Burchinal and Eppright, 1959). The authors concluded that there is no personality, psychopathology or psychological profile that could explain the development of obesity (Friedman and Brownell, 1995).

To date, despite the fact that research has grown and hundreds of studies have been carried out, there is no sufficiently

valid and reliable empirical evidence that clearly demonstrates the existence of a specific psychological and psychopathological configuration among people with obesity (Vaidya, 2006).

In parallel, and subsequently to the etiological studies, some researchers changed their perspective and began to consider the psychological and psychopathological consequences of obesity rather than the causes. The idea spread that personality disorders or psychopathological disorders were more likely the results of being affected by obesity than the causes (Bocchieri *et al.*, 2002). However, the cross-sectional comparisons that were performed between groups of obese individuals and groups of individuals of normal weight produced inconsistent results with respect to the psychopathological consequences of obesity. In fact, some studies showed that people with obesity suffered more than people of normal weight, while others found that obesity had an apparent protective effect against psychological distress. The extreme diversity of the results was largely due to the differences that existed between the samples of individuals recruited in the researches and also between the weak methodological approaches used (Friedman and Brownell, 1995). The inconsistency of the results also reflected an inconsistent phenomenon. Indeed, the effects of being obese vary between people and, therefore, obesity can create serious psychological, emotional and psychosocial problems in some individuals, minor distress in others and possibly no disturbance in still others. In general, first generation studies have collectively suggested that obesity per se is not directly associated with psychological distress but that this association is mediated and moderated by other conditions. Research on the psychological and psychosocial correlates of obesity has therefore progressively entered the second generation and researchers have begun to study the factors which, in the heterogeneous population of people with obesity, seem to increase the risk of developing a psychopathology (Friedman and Brownell, 1995). The best evidence was gathered in some studies conducted on large representative samples of the population, mainly American, and suggested that obese women, but not obese men, have higher levels of depression and suicidal ideation than their normal weight counterparts (Istvan *et al.*, 1992; Carpenter *et al.*, 2000). Being a woman, as suggested by the studies cited above, is one such risk factor. Even severe obesity (BMI > 40) and comorbidity with binge eating disorder (BED) are consistently associated with an increased risk of psychopathology. Research has also documented a long series of psychological and physical vulnerabilities associated with obesity, such as low self-esteem and negative self-perceptions, body image disturbances (Stunkard and Wadden, 1992), sexual problems (Assimakopoulos *et al.*, 2006; Kolotkin *et al.*, 2006), eating disorders (Davin and Taylor, 2009), less social contact (Bocchieri *et al.*, 2002) and early trauma.

## Obesity Metabolic Phenotypes

On the contrary, the focus in medicine has long been on individual differences. Obesity has been studied biologically to identify inter-individual differences in order to provide the patient with the best treatment and increase the chance of a more favourable outcome.

Some nutrition experts have proposed to classify the patient suffering from obesity into phenotypes (Blundell & Cooling, 2000). In particular, four phenotypes of obesity have been described (De Lorenzo *et al.*, 2016):

1. normal weight obese (NWO);
2. normal weight metabolically obese (MONW);
3. metabolically healthy obese (MHO);
4. metabolically unhealthy obese (MUO).

The NWO type presents: normal BMI, high percentage of fat with levels typical of obese subjects and associated with a high level of vascular inflammation which greatly increases the risk of cardiovascular disease. A higher prevalence is found with increasing age and is more frequent among women than men. Arguably, all women older than 55 years with a BMI < 25 kg/m are considered NWO. They have elevated blood pressure, elevated fasting glucose, and a worse lipid profile than healthy normal-weight subjects. They appear to be sarcopenic especially in the lower limbs. They do not show metabolic syndrome and have a high level of oxidative stress. Finally, they have a low basal metabolic rate.

The MONW type has normal BMI, normal weight, normal BMR, but demonstrate metabolic characteristics more prone to the development of metabolic syndrome. They have characteristics typical of obese people: high visceral fat, high fat mass, low lean mass, low insulin sensitivity, high concentration of triglycerides, fatty liver disease and chronic degenerative diseases. It is a more frequent condition in men and when it concerns women it is typical of subjects with PCOS. Often these are smokers, hypertensives, who perform little physical activity, with high levels of CRP, TNF- $\alpha$  and IL-6 and low levels of HDL and adiponectin.

The MHO type demonstrates: an unequivocally obese phenotype, but without any metabolic abnormality. In fact, they have an excess of adipose tissue and a metabolic profile characterized by high insulin sensitivity, an optimal lipid profile and no signs of hypertension. They have a normal BMR and the inflammatory profile is also normal. This condition is often seen in healthy postmenopausal women. Making a comparison with healthy subjects of normal weight, although the parameters are all in the normal range, there are still slightly higher levels of fasting glucose, triglycerides, CRP, LDL and systolic blood pressure and a lower fiber intake and less physical activity. In any case, these subjects demonstrate an increased risk of contracting cardiovascular disease within fifteen years of the diagnosis of MHO and of metabolic syndrome at ten years. However, it has been observed that they are subjects with hepatic steatosis and elevated hs-CRP levels.

The MUO type is characterized by high BMI, high percentage of fat mass, high concentration of visceral adipose tissue, low BMR, high incidence of metabolic syndrome, type 2 diabetes mellitus, atherosclerosis and cardiovascular disease. They have high insulin resistance, an altered lipid profile, fatty liver disease, and elevated levels of inflammation.

## Individual differences in Obesity history

According to the NIH guidelines Patient candidates for bariatric surgery can be subjects with a BMI > 40 kg/m<sup>2</sup>, with a history of repeated failures in weight control treatments or subjects with a BMI >35 kg/m<sup>2</sup>, but with comorbidities (Yermilov *et al.*, 2009).

As is known, the intervention includes a pre-surgical psychological assessment phase useful for promptly identifying any major psychiatric disorders in progress, including, first and foremost, the presence of eating disorders. However, the evaluation, as recommended in the guidelines of the Italian Society of Surgery of Obesity (SICOB), also includes a phase of medical history collection of the history of development of weight problems (Busetto *et al.*, 2011).

In clinical practice we were able to observe that the subjects responded very differently to the question "when did the weight problems start?". In fact:

1. a first group of subjects reports being affected by obesity and therefore on a "constantly diet" condition;
2. a second group, on the other hand, was characterized by having been of normal weight up to a precise moment and by having begun to accumulate weight "after a stressful life event".

There are thus two possible types of individual affected by obesity:

- The individual affected by obesity from birth has a history of failures in weight control, probably already started in childhood, but also a body image - for example the representation of oneself in space - quite stable (Micanti *et al.*, 2016).
- The individual suffering from obesity following a traumatic life event, on the other hand, had experience of the normal weight self and could have used food as a coping strategy for stress, until body image dissatisfaction, depressive symptoms, and low self-esteem (Dalle Grave, 2007).

In more than 220 clinical interviews using the Paykel Life Event Scale (1997), the stressors that triggered the weight gain were identified as:

Most reported Life events among people affected by obesity
Maternal / paternal mourning
Brotherly mourning
One or more abortions
Loss of a child
Pregnancy / postpartum
Marriage
Job loss
Divorce / Separation
Broken bone / accidents / health problems with ospedalization
Moving to another city

Many patients suffering from obesity report that the difficulties in weight control began following a life experience experienced as traumatic. However, to our knowledge, there are no studies that have explored the possible presence of

psychological subtypes among obese patients based on weight history. Most of the studies compared obese subjects with subjects of normal weight (Abilés *et al.*, 2010), or subjects with obesity with and without Binge Eating Disorder (BED) (de Zwaan *et al.*, 2003; Rieger *et al.*, 2005). These subjects could therefore also have a different motivation for the request for intervention and different paths of adherence and compliance after the weight reduction intervention eg. sleeve gastrectomy.

Given these premises, it would be important to understand whether there may be different psychological phenotypes based on the individual history of subjects affected by obesity. The results support the existence of two profiles, which when compared are worse, from a psychological point of view, when subjects develop a weight pathology after a life event.

## Obesity, life stressors and emotional abuse

Studies have shown that experiencing emotional or behavioral abuse in childhood appears to increase the risk of obesity in adulthood. In particular Hemmingsson, Johansson & Reynisdottir, S. (2014) conducted a meta-analysis of 23 studies that included a total of 112,000 participants. The study found that childhood physical abuse increased the risk of obesity by 28 percent, emotional abuse by 36 percent, sexual abuse by 31 percent, and unspecified abuse by 45 percent. Overall, those who have experienced severe abuse have a 50 percent risk of obesity compared with 13 percent for those who have experienced moderate abuse.

Again Hemmingsson (2014) hypothesized that stressful life events leave traces that can manifest themselves late in life through physical ailments. Negative events would lead to greater stress on a physiological level, with consequent negative thoughts and emotions, in association with a possible increase in inflammation in various parts of the body, a lowering of the immune system and metabolism associated with obesity.

These life events, therefore, would have a negative impact on the regulation of appetite, metabolism and eating behaviour up to modifying a series of habits which in the long run lead to the onset of obesity.

Several authors have hypothesized that emotional eating may be a coping response to a stressful event. Going into more detail, it would not be so much the event itself that causes the behavioural response, but its negative interpretation towards the emotional response. The concept of interoceptive awareness in fact includes both the aspect of lack of clarity with respect to the emotion felt, but also the non-acceptance of the emotion felt.

Recent studies have demonstrated a strong association between interoceptive awareness and body image. Researchers have found that people who can sustain attention to their body's internal cues tended to report higher levels of positive body image (Todd *et al.* 2019). People who trust their body's internal signals, such as heart rate or feeling uncomfortable or hungry, are more likely to have a positive view of their body and are less concerned about being overweight (Szulc, 2019).

## Surgical and Treatment outcomes

It is important to underline how the emotional aspects have a fundamental "weight" in the success of the intervention and maintenance of weight loss; in fact, the surgical action alone fails to modify in the long term those dysfunctional attitudes that have been the cause of the problematic relationship with food. The literature tells us that in some cases bariatric surgery has proved unsuccessful in terms of weight loss (Van Hout *et al.*, 2006). Typically patients show a 50-60% reduction two years after surgery but, unfortunately, about 20% of them fail to lose a significant percentage of weight. This failure is very often attributable to psychological factors (Busetto *et al.*, 2005). Therefore, despite the benefits of bariatric surgery, psychological and behavioural factors, likely, they play an influential role in postoperative outcome.

At the same time, with regard to the emotional consequences of a state of obesity, a study conducted in America at the UCSD Medical Center (Russo, 2017) evaluated how surgery can, in some way, influence psychological and emotional factors of patients, such as eating behaviour, mood, anxiety, and quality of life. The results showed that, after the intervention, people tended not to overeat anymore in response to negative emotions, in parallel with a decrease in depressive and anxious symptoms. Quality of life improved even though, being a pilot study, the results were based as close as 6 months postoperatively.

It is clear, therefore, that the psychological component is associated with a constellation of physical, psychic and behavioural variables which, if on the one hand allow the identification of common elements in the obesity "syndrome", on the other leave ample opportunity for considering the person can be explained and treated only if we take into consideration individual history and inner experience.

## Conclusions and psychotherapeutic indications

The possibility of identifying a potential psychological phenotype during the clinical pre-surgical psychological interview, provides the clinician a tool to identify the most indicated intervention and support strategies, focused on increasing post-operative success. These observations could help to reduce stigmatization that exist about people with obesity, the incidence of life events shows that obesity is caused by many factors and not only by overeating or a sedentary lifestyle. The holistic approach is highly recommended and should be adopted in the psychological treatment and prevention of obesity, with a particular attention to the individual's development and to the psychological and emotional aspects connected to patients' history.

The present findings add further empirical data to the literature on the role of life events in the development of obesity, a population often neglected and stigmatized. Individuals with obesity have different histories with different psychological and psychotherapeutic needs.

Therapists and mental health professionals have at their disposal evidence-based treatments such as CBT-OB therapy (Dalle Grave, Sartirana & Calugi, 2020). Considering these observations, however, some reflections can be raised. With regards to the population of people suffering from obesity since ever, the results suggest that they might benefit from

cognitive-behavioral therapies focusing on emotional regulation such as Marsha Linehan's DBT, especially for "suffering tolerance" and "crisis survival" skills. With regards people with obesity starting from life events, who have different characteristics and specific needs, a programme focused on trauma healing, such as EMDR, might be more indicated (Shapiro, 2014; Cairns, 2018). Awareness of individual differences can point to increasingly targeted pre- and post-surgical prevention and may decrease prejudice and care disparity in psychological treatments.

## Acknowledgements

I would like to thank my husband, Dr. Aldo Sitibondo, as physician specialized in Internal Medicine at the University Hospital of Messina, for the critical review of the cited aspects of clinical medicine related to metabolic syndrome and obesity.

## Funding

This study has received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

## Conflicts of Interest

The author declares that there is no financial, general, and institutional conflict of interest regarding the publication of this article.

## Note

The Interview for Recent Life Events - IRLE by Paykel and co-workers (1997), takes into consideration the last six months and divides the events into 10 categories (work, education, economic problems, health, bereavement, emigration, sentimental life, legal problems, family relationships and marital area) and of each event requires to assess (on a scale from 1 to 5) the independence from the illness and the objectively negative impact; the assessment is carried out by means of a semi-structured interview and subsequently validated by means of a score freely attributed by the patient. The items of the Paykel scale with their normative value. In the Italian version, edited by Fava and Osti (1982), the items were increased to 63.

## References

- Abilés, V., Rodríguez-Ruiz, S., Abilés, J., Mellado, C., García, A., De La Cruz, A. P., & Fernández-Santaella,
- Assimakopoulos, K., Panayiotopoulos, S., Iconomou, G., Karaivazoglou, K., Matzaroglou, C., Vagenas, K., &



- Kalfarentzos, F. (2006). Assessing sexual function in obese women preparing for bariatric surgery. *Obesity surgery*, 16(8), 1087-1091.
- Blundell, J. E., & Cooling, J. (2000). Routes to obesity: phenotypes, food choices and activity. *British journal of nutrition*, 83(S1), S33-S38.
  - Bocchieri, L. E., Meana, M., & Fisher, B. L. (2002). A review of psychosocial outcomes of surgery for morbid obesity. *Journal of psychosomatic research*, 52(3), 155-165.
  - Bocchieri, L. E., Meana, M., & Fisher, B. L. (2002). A review of psychosocial outcomes of surgery for morbid obesity. *Journal of psychosomatic research*, 52(3), 155-165.
  - Bruch, H. (1948). Psychological aspects of obesity. *Bulletin of The New York academy of medicine*, 24(2), 73.
  - Slochower, J. (1987). The psychodynamics of obesity: A review. *Psychoanalytic Psychology*, 4(2), 145.
  - Burchinal, L. G., & Eppright, E. S. (1959). Test of the psychogenic theory of obesity for a sample of rural girls. *The American Journal of Clinical Nutrition*, 7(3), 288-294.
  - Busetto, L., Segato, G., De Luca, M., De Marchi, F., Foletto, M., Vianello, M., ... & Enzi, G. (2005). Weight loss and postoperative complications in morbidly obese patients with binge eating disorder treated by laparoscopic adjustable gastric banding. *Obesity Surgery*, 15(2), 195-201.
  - Bussetto, L., Cerbone, M. R., Lippi, C., Micanti, F., & Sampietro, S. (2011). Suggerimenti per la valutazione psicologica psichiatrica del paziente obeso candidato alla chirurgia bariatrica. *SIC OB*.
  - Cairns, S. (2018). Using EMDR to Treat Obesity.
  - Carpenter, K. M., Hasin, D. S., Allison, D. B., & Faith, M. S. (2000). Relationships between obesity and DSM-IV major depressive disorder, suicide ideation, and suicide attempts: results from a general population study. *American journal of public health*, 90(2), 251.
  - Dalle Grave, R., Cuzzolaro, M., Calugi, S., Tomasi, F., Temperilli, F., Marchesini, G., & QUOVADIS Study Group. (2007). The effect of obesity management on body image in patients seeking treatment at medical centers. *Obesity*, 15(9), 2320-2327.
  - Dalle Grave, R., Sartirana, M., & Calugi, S. (2020). Personalized cognitive-behavioural therapy for obesity (CBT-OB): theory, strategies and procedures. *BioPsychoSocial Medicine*, 14(1), 1-8.
  - Davin, S. A., & Taylor, N. M. (2009). Comprehensive review of obesity and psychological considerations for treatment. *Psychology, health & medicine*, 14(6), 716-725.
  - De Lorenzo, A., Soldati, L., Sarlo, F., Calvani, M., Di Lorenzo, N., & Di Renzo, L. (2016). New obesity classification criteria as a tool for bariatric surgery indication. *World journal of gastroenterology*, 22(2), 681.
  - Di Pauli, D. (2016). Where's Icarus? Stigma based on weight that no one wants to see. A candid answer to "A candid discussion of obesity". *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*, 21(3), 349-351.
  - Farooqi, I. S., & O'rahilly, S. (2007). Genetic factors in human obesity. *Obesity reviews*, 8, 37.
  - Friedman, M. A., & Brownell, K. D. (1995). Psychological correlates of obesity: moving to the next research generation. *Psychological bulletin*, 117(1), 3.
  - Hemmingsson, E. (2014). A new model of the role of psychological and emotional distress in promoting obesity: conceptual review with implications for treatment and prevention. *Obesity Reviews*, 15(9), 769-779.

- Hemmingsson, E., Johansson, K., & Reynisdottir, S. (2014). Effects of childhood abuse on adult obesity: a systematic review and meta-analysis. *Obesity Reviews*, 15(11), 882-893.
- Hill, A., Williams, G., & Fruhbeck, G. (2009). Social and psychological factors in obesity. *Obesity: Science to practice*, 347-366.
- Istvan, J., Zavela, K., & Weidner, G. (1992). Body weight and psychological distress in NHANES I. *International Journal of Obesity and Related Metabolic Disorders: Journal of the International Association for the Study of Obesity*, 16(12), 999-1003.
- Kolotkin, R. L., Binks, M., Crosby, R. D., Østbye, T., Gress, R. E., & Adams, T. D. (2006). Obesity and sexual quality of life. *Obesity*, 14(3), 472-479.
- Linehan, M. (2014). *DBT. Skills training manual*: Guilford Publications.
- M. C. (2010). Psychological characteristics of morbidly obese candidates for bariatric surgery. *Obesity surgery*, 20(2), 161-167.
- McPherson, K., Marsh, T., & Brown, M. (2007). Foresight report on obesity. *The Lancet*, 370(9601), 1755.
- Micanti, F., Iasevoli, F., Cucciniello, C., Costabile, R., Loiarro, G., Pecoraro, G., ... & Galletta, D. (2017). The relationship between emotional regulation and eating behaviour: a multidimensional analysis of obesity psychopathology. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*, 22(1), 105-115.
- Paykel, E. S. (1997). The interview for recent life events. *Psychological medicine*, 27(2), 301-310.
- Puhl, R. M., & Heuer, C. A. (2010). Obesity stigma: important considerations for public health. *American journal of public health*, 100(6), 1019-1028.
- Pull, C. B. (2010). Current psychological assessment practices in obesity surgery programs: what to assess and why. *Current opinion in psychiatry*, 23(1), 30-36.
- Rieger, E., Wilfley, D. E., Stein, R. I., Marino, V., & Crow, S. J. (2005). A comparison of quality of life in obese individuals with and without binge eating disorder. *International Journal of Eating Disorders*, 37(3), 234-240.
- Russo, D. (2017). *The Relationship Between Psychosocial Factors and Weight Loss Maintenance in a Sample of Bariatric Surgery Patients* (Doctoral dissertation, Alliant International University).
- Shapiro, F. (2014). The role of eye movement desensitization and reprocessing (EMDR) therapy in medicine: addressing the psychological and physical symptoms stemming from adverse life experiences. *The Permanente Journal*, 18(1), 71.
- Smith, K. B., & Smith, M. S. (2016). Obesity statistics. *Primary care: clinics in office practice*, 43(1), 121-135.
- Stunkard, A. J., & Wadden, T. A. (1992). Psychological aspects of severe obesity. *The American journal of clinical nutrition*, 55(2), 524S-532S.
- Szulc, J. S. (2019). *Adapting an Integrated Model of Body Appreciation in Women: The Role of Interoceptive Awareness* (Doctoral dissertation).
- Todd, J., Aspell, J. E., Barron, D., & Swami, V. (2019). Multiple dimensions of interoceptive awareness are associated with facets of body image in British adults. *Body Image*, 29, 6-16.
- Vaidya, V. (2006). Psychosocial aspects of obesity. *Health and Treatment strategies in obesity*, 27, 73-85.
- World Health Organization. (2000). *Obesity: preventing and managing the global epidemic*.

- Yermilov, I., McGory, M. L., Shekelle, P. W., Ko, C. Y., & Maggard, M. A. (2009). Appropriateness criteria for bariatric surgery: beyond the NIH guidelines. *Obesity*, 17(8), 1521-1527.
- de Zwaan, M., Mitchell, J. E., Howell, L. M., Monson, N., Swan-Kremeier, L., Crosby, R. D., & Seim, H. C. (2003). Characteristics of morbidly obese patients before gastric bypass surgery. *Comprehensive Psychiatry*, 44(5), 428-434.
- van Hout, G., Boekestein, P., Fortuin, F. A., Pelle, A. J., & van Heck, G. L. (2006). Psychosocial functioning following bariatric surgery. *Obesity Surgery*, 16(6), 787-794.