

Open Peer Review on Qeios

Stage IVA Oral Cavity Mucoepidermoid Carcinoma AJCC v6 and v7

National Cancer Institute

Source

National Cancer Institute. <u>Stage IVA Oral Cavity Mucoepidermoid Carcinoma AJCC v6 and v7.</u> NCI Thesaurus. Code C5868.

Stage IVA includes: (T4a, N0, M0); (T4a, N1, M0); (T1, N2, M0); (T2, N2, M0); (T3, N2, M0); (T4a, N2, M0). T4a: Oral cavity cancer with moderately advanced local disease. Tumor invades adjacent structures only (e.g., through cortical bone [mandible or maxilla] into deep [extrinsic] muscle of tongue [genioglossus, hyoglossus, palatoglossus, and styloglossus], maxillary sinus, skin of face). T1: Tumor 2 cm or less in greatest dimension. T2: Tumor more than 2 cm but not more than 4 cm in greatest dimension. T3: Tumor more than 4 cm in greatest dimension. N0: No regional lymph node metastasis. N1: Metastasis in a single ipsilateral lymph node, 3 cm or less in greatest dimension. N2: Metastasis in a single ipsilateral lymph node, more than 3 cm but not more than 6 cm in greatest dimension; or in multiple ipsilateral lymph nodes, none more than 6 cm in greatest dimension; or in bilateral or contralateral lymph nodes, none more than 6 cm in greatest dimension. M0: No distant metastasis. (AJCC 7th ed.)

Qeios ID: NTZJGK · https://doi.org/10.32388/NTZJGK