

## Research Article

# Interventions to increase personal protective behaviours to limit the spread of respiratory viruses: A rapid evidence review and meta-analysis

Olga Perski<sup>1</sup>, Dorothy Szinay<sup>2</sup>, Elizabeth Corker<sup>1</sup>, Lion Shahab<sup>1</sup>, Robert West<sup>1</sup>, Susan Michie<sup>1</sup>

1. University College London, University of London, United Kingdom; 2. University of East Anglia, United Kingdom

**Purpose:** Increasing personal protective behaviours is critical for stopping the spread of respiratory viruses, including SARS-CoV-2: we need evidence to inform how to achieve this. We aimed to synthesise evidence on interventions to increase six personal protective behaviours (e.g. hand hygiene, face mask use) to limit the spread of respiratory viruses.

**Methods:** We used best practice for rapid evidence reviews. We searched Ovid MEDLINE and Scopus. Studies conducted in adults or children with active or passive comparators were included. We extracted data from published intervention descriptions on study design, intervention content, delivery mode, population, setting, mechanism(s) of action, acceptability, practicability, effectiveness, affordability, spill-over effects and equity impact. Study quality was assessed with Cochrane's risk of bias tool. A narrative synthesis and random-effects meta-analyses were conducted.

**Results:** We identified 39 studies conducted across 15 countries. Interventions targeted hand hygiene (n=30) and/or face mask use (n=12) and used two- or three-arm study designs with passive comparators. Interventions were typically delivered face-to-face and included a median of three behaviour change techniques. The quality of included studies was low. Interventions to increase hand hygiene (k=6) had a medium, positive effect ( $d=0.62$ , 95% CI=0.43–0.80,  $p<.001$ ,  $I^2=81.2\%$ ). Interventions targeting face mask use (k=4) had mixed results, with an imprecise pooled estimate (OR=4.14, 95% CI=1.24–13.79,  $p<.001$ ,  $I^2=89.67\%$ ). Between-study heterogeneity was high.

**Conclusions:** We found low-quality evidence for positive effects of hand hygiene interventions, with unclear results for face mask use interventions. There was a lack of evidence for interventions targeting most behaviours of interest within this review.

## Introduction

Respiratory viruses such as influenza, respiratory syncytial virus, parainfluenza, rhinovirus, coronavirus (including SARS-CoV-2) and adenovirus enter the body through the eyes, nose and mouth (the 'T-Zone') (Killingley & Nguyen-Van-Tam, 2013b; West et al., 2020). Changing human behaviour is critical for stopping the spread of respiratory viruses in general and the SARS-CoV-2 virus in particular, and for supporting the easing of financially and psychologically costly physical distancing measures during viral epidemics (Ferguson et al., 2006; Michie, Rubin, et al., 2020; Michie, West, & Amlôt, 2020; West et al., n.d.). Personal protective behaviours, including hand washing, disinfecting fomites such as clothes or furniture, and face mask wearing, are advocated for limiting the spread of SARS-CoV-2 (Lunn et al., 2020; World Health Organisation, 2019). Simply advising people to adopt these behaviours has been found to be insufficient, just as has explaining what to do and why these behaviours are necessary (Bish & Michie, 2010). Directly relevant evidence on interventions to promote adherence to personal protective behaviours in community-dwelling children and adults is sparse but there is an urgent need to identify and synthesise what evidence does exist. Policymakers need evidence to inform the development of public health guidance and decide which interventions to prioritise. We adopted best practice for rapid evidence reviews to evaluate the acceptability, practicability, effectiveness, affordability, spill-over effects (i.e. unintended consequences) and equity impact (the 'APEASE' criteria (Michie et al., 2014)) of interventions to increase personal protective behaviours to limit the spread of respiratory viruses.

During pandemics of respiratory viruses, multipronged approaches involving both pharmacological (e.g. vaccination) and behavioural measures (e.g. hand washing, physical distancing) are required to bring the reproductive number below 1 (Ferguson et al., 2006; Michie, Rubin, et al., 2020; Michie, West, & Amlôt, 2020; Michie, West, Amlôt, et al., 2020). Vaccination of populations will take months, even years, to roll out, especially in low- and middle-income countries. Hence, physical distancing and other behavioural measures will be required, possibly permanently. Population-wide restrictions are costly from financial, social and psychological perspectives: the world economy has been projected to shrink by approximately 4.9% in 2020 (International Monetary Fund, 2020), with an additional 88 million people globally being pushed into extreme poverty (i.e. living on less than \$1.90/day) (Blake & Wadhwa, 2020), and prolonged periods of social isolation are associated with increases in domestic violence (SafeLives, 2020) and negative mental health effects, such as post-traumatic stress disorder, confusion and anger (Brooks et al., 2020). Less costly, yet highly effective (Warren-Gash et al., 2012) personal protective behaviours are thus important for supporting the easing of lockdown measures to ensure long-term suppression of viral transmission and preparedness for new viral waves and future pandemics (Michie, West, & Amlôt, 2020; Michie, West, Amlôt, et al., 2020; West et al., n.d.). To successfully block the

spread of respiratory viruses including (but not limited to) SARS-CoV-2 – which are transmitted via droplets, aerosols and direct physical contact (Killingley & Nguyen-Van-Tam, 2013a; West et al., 2020) – several personal protective behaviours must be adopted across the population (see Figure 1). Although systematic reviews of interventions to change hand hygiene in healthcare professionals are available (Edwards et al., 2012; Huis et al., 2012; Luangasanatip et al., 2015; Mbakaya et al., 2017; Olena Doronina et al., 2017), generalisability to community settings is limited. There also appears to be little evidence about interventions to change behaviours such as not touching the T-Zone (eyes, nose and mouth), which would have a significant effect if adopted (Kwok et al., 2015) and carry little or no costs to people or society. If adopted at scale across the population including disadvantaged communities, such interventions have the potential to reduce health inequalities. Here, we aimed to conduct a rapid evidence review to evaluate the acceptability, practicability, effectiveness, affordability, spill-over effects and equity of interventions to increase personal protective behaviours that limit the spread of respiratory viruses.




Figure 1. Map of personal protective behaviours relevant for blocking transmission of respiratory viruses, including SARS-CoV-2: hand washing and use of hand sanitizers; avoiding touching the 'T-Zone'; catching droplets in tissues and discarding these; face mask use; disinfecting surfaces; and maintaining physical distancing. Reproduced with permission from the authors (West & Michie, 2020).

## Methods

### Study design

The study protocol was pre-registered on the Open Science Framework (<https://osf.io/7cphy>). During ongoing pandemics, the World Health Organization recommends the use of rapid evidence reviews for swift knowledge generation (World Health Organisation, 2017). We adopted acknowledged best practice for rapid evidence reviews, which involved completing the review in a timely fashion, limiting the search to main databases and the published literature, and having one reviewer extract data and another verify (Haby et al., 2016; Tricco et al., 2015).

### Criteria for considering studies for this review

#### Population

We included studies that recruited as participants community-dwelling children or adults (as opposed to qualified or trainee healthcare professionals in hospital or care home settings) across any type of study setting (e.g. schools, primary care).

#### Intervention

We included reports of evaluations of any type of intervention (e.g. mass media, face-to-face, technology-mediated) designed to change at least one of six personal protective behaviours to block transmission of respiratory viruses, such as influenza, respiratory syncytial virus, parainfluenza, rhinovirus, coronavirus or adenovirus, which have a shared route of transmission (i.e. droplets, aerosols, direct physical contact) (Killingley & Nguyen-Van-Tam, 2013a). Although the relative importance of different personal protective behaviours depends on properties of the specific respiratory virus in addition to the clinical and/or environmental context – for example, fomite transmission may be more pronounced for respiratory syncytial virus compared with coronaviruses (Boone & Gerba, 2007) – at the time of planning this rapid review, little was known about the properties of SARS-CoV-2, and we therefore opted for an inclusive scope.

#### Comparison

We included studies with an active or passive (e.g. wait-list control, baseline) comparator.

#### Outcomes

We included studies that reported as outcome either the acceptability, practicability, effectiveness, affordability, spill-over effects and/or equity of interventions, provided that these were measured at the individual level via self-report or direct observation.

#### Study designs

We included primary research studies that used experimental (e.g. randomised controlled trial) or quasi-experimental (e.g. pre- and post-test) study designs, with individuals or clusters as the unit of randomisation, providing that they were conducted under free-living (as opposed to laboratory) conditions. In line with rapid review guidelines, we only included studies that were published in peer-reviewed journals and written in English (Tricco et al., 2015; World Health Organisation, 2017).

#### *Search methods for identification of studies*

##### Electronic searches

We searched Ovid MEDLINE and Scopus. Search terms for each behaviour (e.g. 'hand hygiene', 'hand washing', 'face mask') were piloted and refined to achieve balance between sensitivity and specificity (see Supplementary file 1).

##### Searching for other sources

Expertise within the review team and consultation with topic experts was used to identify additional articles of interest. We had specified the use of reference chaining in the review protocol; however, given the rapid focus of the review and large number of identified studies, reference chaining was not performed.

#### *Data collection and analysis*

##### Selection of studies

Two reviewers (OP, DS) independently screened i) titles and abstracts and ii) full texts against the inclusion criteria. Discrepancies were discussed and resolved through consulting with a third reviewer (EC) if necessary.

##### Data extraction and management

-

A data extraction form was developed on the basis of the Behaviour Change Intervention Ontology (

[www.humanbehaviourchange.org](http://www.humanbehaviourchange.org)

) and Cochrane's PICO ontology (

<https://linkeddata.cochrane.org/pico-ontology>

). Ontologies are classification systems which enable researchers to specify entities (e.g. behaviours, interventions) and their inter-relationships. The use of ontologies in systematic reviews can help ensure that a comprehensive set of entities are considered and defined in standardised ways, thus facilitating systematic knowledge synthesis (Norris et al., 2019). We extracted data from relevant sections of published articles and available supplementary materials on study design, intervention content (i.e. behaviour change techniques (BCTs), coded against the BCT Taxonomy v1 (Michie et al., 2013)), mode of delivery, population, setting, and mechanism(s) of action (Carey et al., 2019; Moore & Evans, 2017). As a validated taxonomy of mechanisms of action is, to our knowledge, not yet available, authors' own definitions of mechanisms of action were extracted if they explicitly discussed how the selected intervention components/BCTs were expected to influence the target behaviour. Due to limited resources, we did not contact study authors for more detail on intervention descriptions. As an intervention may be effective but have negative spill-over effects to other behaviours, or be impracticable and/or

unacceptable to key stakeholders, we also deemed it important to extract data from relevant sections of published articles and available supplementary materials pertaining to the APEASE criteria (see Table 1) (Michie et al., 2014). Although criteria such as affordability or practicability are arguably closely tied to the context in which the intervention was/will be implemented, APEASE intends to capture higher-order criteria against which to rate interventions, broadly applicable irrespective of the specific context. Therefore, authors' own descriptions of, for example, acceptability or spill-over effects (as opposed to reviewers' ratings) were extracted. Spill-over effects were broadly defined as any unintended consequences (positive or negative) reported by the authors, including but not limited to other behaviours that were changed by the intervention that it was not designed to target (e.g. teachers' improved hand hygiene positively or negatively influencing children's hand hygiene). Data were extracted by one reviewer (OP or DS). In the review protocol, we had specified that extracted data would be verified by a second reviewer to assess accuracy and completeness. However, given the large number of identified studies, a second reviewer (EC) verified 10% of studies.

Table 1. APEASE criteria for evaluating intervention approaches or components (Michie et al., 2014).

Criterion	To what extent ...
Acceptability	...is the intervention judged to be acceptable by all key stakeholders
Practicability	...can the intervention be delivered as intended at the scale intended and in the context intended
Effectiveness	...will the intervention deliver the desired outcome in the target population
Affordability	...can the intervention be afforded within an acceptable budget
Spill-over effects	...is the intervention likely to have additional negative or positive consequences
Equity	...is the intervention likely to increase or decrease inequalities in society

#### Quality appraisal

The methodological rigour of included evaluation reports was assessed by one reviewer (OP or DS) using Cochrane's risk of bias tool (The Cochrane Collaboration, 2011). A second reviewer (EC) verified 10% of studies.

#### Stakeholder involvement

We solicited input from key stakeholders, including patient and public representatives recruited via panels convened by Public Health England (n = 282) and the University of East Anglia (n = 3), and UK policymakers and academic researchers contacted via a mailing list on the research objectives, target behaviours and outcomes assessed. Feedback from patient and public representatives (n = 20) was incorporated into the review protocol; we did not receive any suggestions for improvement or clarification from the policymakers and academic researchers. The rapid review results will be disseminated to stakeholders via an infographic.

#### Data synthesis

A narrative (descriptive) synthesis was conducted for each of the personal protective behaviours. We had specified in the review protocol that meta-analyses would be conducted if practicable and appropriate (i.e. >5 studies with homogeneous study designs and outcome variables). After inspection of study designs and outcome variables, however, we deemed it useful to conduct a meta-analysis with k = 4 studies. Random-effects meta-analyses to estimate a pooled odds ratio (OR) or standardised mean difference (d) were conducted in RStudio v.1.2.5033 with the *metafor* package (Viechtbauer, 2010). Cohen's conventions for small (d = 0.2), medium (d = 0.5) and large (d = 0.8) effects were used in the interpretation of the results (Cohen, 1988). In studies with more than two arms (e.g. three-arm RCTs), we compared the 'most active' (i.e. the arm with the greatest number of intervention components) and the 'most passive' arms (i.e. the arm with the lowest number of intervention components, typically labelled the 'control' arm by study authors). Where studies reported more than one hand hygiene outcome (e.g. hand washing and hand sanitizer use), only the first reported outcome was included in the meta-analysis, so as not to violate the assumption of independence (Cheung, 2019). Where studies did not report sufficient detail to calculate effect sizes, authors' own description/interpretation of results were grouped into 'positive' effects (i.e. a significant difference between intervention and control groups, favouring the intervention group, was detected), 'no difference' (i.e. a significant difference between groups was not detected), 'negative' effects (i.e. a significant difference between groups, favouring the control group, was detected) or 'indeterminate' (i.e. differences between groups were not reported or could not be computed given the study design). To aid interpretation, for behaviours where a majority of positive or negative results were observed, overall results were categorised as either 'positive' or 'negative', respectively. If consistent results were not observed or could not be determined, overall results were categorised as 'mixed'.

## Results

### Study selection

After removing duplicates, 5,594 records were identified, with 158 studies carried forward to the full text screening. Of the 39 studies included in the narrative evidence synthesis, 10 were included in meta-analyses (see Figure 2).




Figure 2. PRISMA flow chart of included studies.

### Study characteristics

Studies were conducted in the United States (13/39; 33%) (Aiello et al., 2010a, 2012; Arbogast et al., 2016a; Azman et al., 2013; Koep et al., 2016; Larson et al., 2009, 2010; Mott et al., 2007; Sandora et al., 2005a; Stebbins et al., 2010; Stedman-Smith et al., 2015a; Updegraff et al., 2011; White et al., 2003), China (5/39; 13%) (Chan et al., 2007; B J Cowling et al., 2009; Benjamin J. Cowling et al., 2008; Liu et al., 2019; Or et al., 2020), Germany (3/39; 13%) (Hübner et al., 2010a; T Suess et al., 2011; Thorsten Suess et al., 2012), Thailand (3/39; 13%) (Apisarnthanarak et al., 2009; Kaewchana et al., 2012; Simmerman et al., 2011b), Australia (2/39; 5%) (MacIntyre et al., 2009; Roberts et al., 2000a), Denmark (2/39; 5%) (Bundgaard et al., 2020; Nandrup-Bus, 2009a), Spain (2/39; 5%) (Azor-Martinez et al., 2016a, 2018a), the United Kingdom (2/39; 5%) (Little et al., 2014; Yardley et al., 2011), with one study each in Australia/Saudi Arabia (Barasheed et al., 2014a), Bangladesh (Ram et al., 2015), Costa Rica (Reyes Fernández et al., 2015a), Finland (Savolainen-Kopra et al., 2012a), France (Canini et al., 2010), the Netherlands (Zomer et al., 2016), and Turkey (Öncü et al., 2019) (see Table 2). Studies had a median of 419 participants (range: 96 to 20,066).

The majority of studies (20/39; 51%) targeted multiple respiratory infections, including (but not limited to) influenza, respiratory syncytial virus and adenovirus (Arbogast et al., 2016b; Azor-Martinez et al., 2018b; Barasheed et al., 2014b; Hübner et al., 2010b; Larson et al., 2009, 2010; Liu et al., 2019; MacIntyre et al., 2009; Mott et al., 2007, p. 20; Nandrup-Bus, 2009b; Öncü et al., 2019; Reyes Fernández et al., 2015b; Roberts et al., 2000b; Sandora et al., 2005b; Savolainen-Kopra et al., 2012b; Stebbins et al., 2010; Stedman-Smith et al., 2015b, p.; White et al., 2003; Yardley et al., 2011; Zomer et al., 2016), with the remaining studies targeting influenza (13/39; 33%) (Aiello et al., 2012; Apisarnthanarak et al., 2009; Azman et al., 2013; Azor-Martinez et al., 2016b; Canini et al., 2010; B J Cowling et al., 2009; Benjamin J. Cowling et al., 2008; Kaewchana et al., 2012; Koep et al., 2016; Little et al., 2014; Or et al., 2020; Ram et al., 2015; Simmerman et al., 2011a), pandemic influenza A (H1N1) (4/39; 10%) (Aiello et al., 2010a; T Suess et al., 2011; Thorsten Suess et al., 2012; Updegraff et al., 2011), SARS (1/39; 3%) (Chan et al., 2007) and SARS-CoV-2 (1/39; 3%) (Bundgaard et al., 2020) (see Table 2).

The majority of studies (28/39; 72%) targeted hand hygiene, with the remaining studies targeting a combination of hand hygiene and/or face mask use (9/39; 23%) (Aiello et al., 2010a, 2012; B J Cowling et al., 2009; Benjamin J. Cowling et al., 2008; Larson et al., 2010; MacIntyre et al., 2009; Simmerman et al., 2011b; T Suess et al., 2011; Thorsten Suess et al., 2012), face mask use only (3/39; 8%) (Barasheed et al., 2014a; Bundgaard et al., 2020; Canini et al., 2010) or a combination of catching of droplets in tissues and hand hygiene (1/39; 3%) (Chan et al., 2007) (see Table 2). Interventions were delivered in participants' own homes (12/39; 31%), nurseries/schools (10/39; 26%), university residence halls/public areas (5/39; 13%), offices (4/39; 10%), online (2/39; 5%), outpatient clinics (1/39; 3%), or an army training centre (1/39; 3%). Four studies did not state the setting for intervention delivery. Studies targeted children and adult household members (16/39; 41%), pre- or school children and/or teachers (8/39; 21%), university staff and/or students (5/39; 13%), office workers (4/39; 10%), community-dwelling adults (3/39; 8%), Hajj pilgrims (1/39; 3%), military trainees (1/39; 3%) or adult members of a social service centre (1/39; 3%).

Study designs used were two-arm, cluster RCTs (11/39; 28%), three-arm, cluster RCTs (10/39; 26%), two-arm RCTs (5/39; 13%), single-arm, pre- and post-intervention studies (5/39; 13%), two-arm, non-randomised cohort studies (2/39; 5%), three-arm RCTs (2/39; 5%), two-arm, pilot RCTs (1/39; 3%) or randomised, cross-over studies (1/39; 3%).

### **Intervention characteristics**

Intervention durations ranged from one day (i.e. one-off interventions) to three years (see Table 3). In studies using a two- or three-arm design (33/39; 85%), comparators included no intervention/usual care (19/33; 58%), educational materials (9/33; 27%), the provision of soap/hand sanitizer (3/33; 9%) or a combination of educational materials and soap/hand sanitizer (2/33; 6%). Interventions were delivered via face-to-face sessions (23/39; 59%), written materials (including books and newsletters) (14/39; 36%), posters/bulletin boards/cue cards (8/39; 21%), cartoons/games (4/39; 10%), videos (4/39; 10%), telephone (3/39; 8%) and/or websites (2/39; 5%). Five studies did not clearly report on the mode of intervention delivery. Where reported, participants received a flat payment for study completion (Aiello et al., 2010a; Benjamin J. Cowling et al., 2008; Cowling, B.J., Chan, K.H., FAng, V.J., Cheng, 2009; Simmerman et al., 2011b; T Suess et al., 2011; Thorsten Suess et al., 2012; White et al., 2003), payment per survey completed (Stebbins et al., 2010), points for study completion (Arbogast et al., 2016a) or points per survey completed (Stedman-Smith et al., 2015a).

We coded 15 different BCTs across intervention descriptions (see Figure 3); however, details on intervention content were typically lacking. Interventions included a median of three BCTs (range: 0 to 6). The most frequently coded BCTs were '12.5 Adding objects to the environment' (17/39; 44%), '4.1 Instruction on how to perform the behaviour' (16/39; 41%) and '5.1 Information about health consequences' (10/39; 26%). Few studies reported targeting a specific mechanism of action. Where reported, interventions were designed to target intentions (Yardley et al., 2011), attitudes (Updegraff et al., 2011; Yardley et al., 2011), subjective norms (Updegraff et al., 2011; Yardley et al., 2011), perceived behavioural control (Liu et al., 2019; Yardley et al., 2011), perceived risk of infection/disease severity (Liu et al., 2019; Updegraff et al., 2011; Yardley et al., 2011), action control (Reyes Fernández et al., 2015a), coping planning (Reyes Fernández et al., 2015a), knowledge (Liu et al., 2019; Or et al., 2020) or skills (Or et al., 2020).




Figure 3. Frequencies of behaviour change techniques (BCTs) coded in published intervention descriptions.

### Acceptability

Fourteen studies reported on the acceptability of interventions. Indicators assessed included mask comfort (Aiello et al., 2012; Barasheed et al., 2014a; Canini et al., 2010; MacIntyre et al., 2009; T. Suess et al., 2011; Thorsten Suess et al., 2012), skin problems/irritation (Azor-Martinez et al., 2016a, 2018a; Little et al., 2014; Nandrup-Bus, 2009a; Sandora et al., 2005a), adverse events (Benjamin J. Cowling et al., 2008), liking/positive impressions (Arbogast et al., 2016a) and ease of understanding (Updegraff et al., 2011), with a small number of participants experiencing discomfort or irritation in the majority of studies that reported these outcomes (see Table 4).

### Practicability

One study (MacIntyre et al., 2009) considered the practicability of scaling up mask fit testing outside the study setting and decided against including routine fit testing as part of the intervention (see Table 4).

### Effectiveness

Studies relied on self-report (25/39; 64%), direct observation (7/39; 18%), a combination of self-report and direct observation (5/39; 13%) or photos/video (2/39; 5%) to examine intervention effectiveness (see Table 4). Outcome variables were heterogeneous across studies (e.g. the frequency or amount of hand sanitizer/soap use per day, the rate of compliance with hand hygiene, or the rate of compliance with face mask use).

### Hand hygiene

.

Overall, the 30 studies pertaining to hand hygiene behaviours (including hand washing and/or hand sanitizer/soap use) had positive results, with 19 studies reporting positive effects (Aiello et al., 2010b, 2012; Apisarnthanarak et al., 2009; Arbogast et al., 2016a; Chan et al., 2007; Benjamin J. Cowling et al., 2008; Kaewchana et al., 2012; Larson et al., 2009; Liu et al., 2019; Mott et al., 2007; Öncü et al., 2019; Or et al., 2020; Simmerman et al., 2011b; Stebbins et al., 2010; T. Suess et al., 2011; Updegraff et al., 2011; White et al., 2003; Yardley et al., 2011; Zomer et al., 2016), three studies reporting negative effects (Aiello et al., 2010b; Larson et al., 2009; Mott et al., 2007), six studies reporting no difference (Aiello et al., 2012; B J Cowling et al., 2009; Koep et al., 2016; Larson et al., 2010; Reyes Fernández et al., 2015a; Stedman-Smith et al., 2015a) and six studies with indeterminate results (Hübner et al., 2010a; Nandrup-Bus, 2009a; Ram et al., 2015; Roberts et al., 2000a; Sandora et al., 2005a; Savolainen-Kopra et al., 2012a) (see Figure 4). It should be noted that some studies reported more than one hand hygiene outcome (e.g. hand washing and hand sanitizer use).




Figure 4. Proportions of reported effects (i.e. positive, negative, no difference or indeterminate) of interventions targeting hand hygiene behaviours and face mask use.

A random-effects meta-analysis ( $k = 6$ ) found a medium, positive effect of interventions on the average frequency of hand hygiene behaviour,  $d = 0.62$ , 95% CI = 0.43–0.80,  $p < .001$  (see Figure 5). However, between-study heterogeneity was high ( $I^2 = 81.2\%$ ).






Figure 5. Forest plot for the standardised mean difference ( $d$ ) in the frequency of hand hygiene behaviour in intervention and control or pre- and post-study comparisons. The comparison in Chan (Chan et al., 2007) pertains to a pre- and post-study comparison; the remaining studies were two- or three-arm RCTs.

#### Face mask use

Overall, the 12 studies pertaining to face mask use reported mixed results, with three studies reporting positive effects (Barasheed et al., 2014a; B J Cowling et al., 2009; Benjamin J. Cowling et al., 2008), two studies reporting no difference (Aiello et al., 2012; MacIntyre et al., 2009), one study reporting negative effects (Aiello et al., 2010a) and six studies with indeterminate results (Bundgaard et al., 2020; Canini et al., 2010; Larson et al., 2010; Simmerman et al., 2011b; T Suess et al., 2011; Thorsten Suess et al., 2012) (see Figure 4).

A random-effects meta-analysis ( $k = 4$ ) found a large, positive effect of interventions on the odds of compliance with face mask use,  $OR = 4.14$ , 95%  $CI = 1.24-13.79$ ,  $p < .001$  (see Figure 6). However, between-study heterogeneity was high ( $I^2 = 89.67\%$ ) and the confidence interval for the pooled effect was wide.




Figure 6. Forest plot for the odds of compliance with face mask use in intervention compared with control arms. The comparison in MacIntyre (MacIntyre et al., 2009) pertains to the surgical mask (intervention) versus the P2 mask arm (control) at the longest point of follow-up.

#### Affordability

Two studies considered the affordability of interventions, with one study discussing the cost of face masks (Bundgaard et al., 2020), which may act as a barrier for wider roll-out, and a second study (Zomer et al., 2016) reporting that although they wanted to provide hand hygiene products to all day care centre groups, they could only afford to do so for a maximum of two groups per centre due to budget restrictions, thus indicating that the selected intervention was not affordable at scale (see Table 4).

#### Spill-over effects

Three studies reported on secondary behaviour change (i.e. positive or negative spill-over to other behaviours), with one study (Zomer et al., 2016) assessing teachers' supervision of children's hand washing (in addition to their own hand washing), a second (Mott et al., 2007) assessing the impact of the intervention on military leaders' hand sanitizer use (in addition to trainees') and a third (Larson et al., 2009) examining multiple hand hygiene behaviours in the same group of participants. The first study reported no change, the second a significant increase in hand sanitizer use and the third a negative impact on hand washing with soap (see Table 4).

#### Equity

Four studies reported on the equity of interventions, with one study (Yardley et al., 2011) reporting that the intervention was equally effective for participants from high and low socioeconomic status groups and three studies (Azman et al., 2013; Azor-Martinez et al., 2016a; Larson et al., 2009) reporting differential intervention effectiveness by educational attainment, parental income or ethnicity, with better outcomes reported in those with high educational attainment, high parental income and from a Black ethnic background (see Table 4).

#### Quality of included studies

One study received an overall rating of 'low risk of bias', with 16 studies rated as 'some concern', 18 as 'high risk of bias', and for four studies, a rating could not be applied (see Table 5).

Table 2. Characteristics of included studies.

Lead author (year)	Country	Target behaviour(s)	Respiratory virus targeted	Study design	Population	Sample size	Mean age (SD)	% Female	% Post-16 educational qualifications	Setting	
1) Aiello (2010)	United States	Face mask use and hand hygiene	Pandemic influenza A (H1N1)	Three-arm, cluster RCT	University students in residence halls	1297 (face mask + hand hygiene = 367; face mask = 378; control = 552)	18.7 (0.8)	66%	100%	University residence halls	I
2) Aiello (2012)	United States	Face mask use and hand hygiene	Influenza A or B	Three-arm, cluster RCT	University students in residence halls	1178 (face mask + hand hygiene = 362; face mask = 420; control = 396)	19.0 (0.9)	55%	100%	University residence halls	I
3) Apisarnthanarak (2009)	Thailand	Hand hygiene	Influenza	Single arm, pre- and post-intervention study	Preschool children	240	5.0 (1.7)	49%	0%	Private kindergarten	I
4) Arbogast (2016)	United States	Hand hygiene	Respiratory syncytial virus, adenovirus, influenza	Two-arm, cluster RCT	Office workers	1386 (intervention = 604; control = 782)	47.0 (0.4)	78%	Not reported	Office buildings	E
5) Azman (2013)	United States	Hand hygiene	Influenza A or B	Two-arm, cluster RCT	Households	3360	Not reported	Not reported	0%	Schools	I
6) Azor-Martinez (2016)	Spain	Hand hygiene	Influenza	Two-arm RCT	School children	1341	8.0 (2.3)	68%	0%	Schools	I
7) Azor-Martinez (2018)	Spain	Hand hygiene	Respiratory viral infections	Three-arm, cluster RCT	Households with a child at a day care centre	911 children	Unclear	Unclear	Unclear	Day care centres	I
8) Barasheed (2014)	Australia/Saudi Arabia	Face mask use	Rhinovirus, influenza, parainfluenza	Two-arm, pilot RCT	Haji pilgrims	164	Not reported	Not reported	Not reported	Not reported	d Is at st
9) Bundgaard (2020)	Denmark	Face mask use	SARS-CoV-2	Two-arm RCT	Community-dwelling adults	6304	47.2 (13.5)	64%	Not reported	Participants' own homes	ac i

											cc o
10) Canini (2010)	France	Face mask use	Influenza	Two-arm, cluster RCT	Households in three regions of France with an index patient	105	26.5 (16.0)	Not reported	Not reported	GP offices	
11) Chan (2007)	China	Catching droplets in tissues and hand hygiene	SARS	Single arm, pre- and post-intervention study	Registered members of a government subsidised social service centre	122	Not reported	63%	20%	Participants' own homes	Te
12) Cowling (2008)	China	Face mask use and hand hygiene	Influenza A or B	Three-arm, cluster RCT	Households with an index patient	198	Not reported	56%	Not reported	Participants' own homes	
13) Cowling (2009)	China	Face mask use and hand hygiene	Influenza A or B	Three-arm, cluster RCT	Households with an index patient	407 index patients	Not reported	51%	Not reported	Participants' own homes	
14) Hübner (2010)	Germany	Hand hygiene	Common cold, influenza	Two-arm RCT	Office workers	134	44.6 (-)	86%	Not reported	Office buildings	
15) Kaewchana (2012)	Thailand	Hand hygiene	Influenza	Two-arm RCT	Households with an index paediatric patient	275 (frequency assessment) + 330 (quality assessment) households	34.7 (13.8)	58%	Not reported	Participants' own homes	1
16) Koep (2016)	United States	Hand hygiene	Influenza	Two-arm, non-randomised cohort study	School children	260	Not reported	46%	0%	Schools	1
17) Larson (2009)	United States	Hand hygiene	Rhinovirus, coronavirus, parainfluenza virus, respiratory syncytial virus, adenovirus, influenza, enterovirus, etc	Single arm, pre- and post-intervention study	Households	422 households	Not reported	Not reported	60%	Participants' own homes	No : (e
18) Larson (2010)	United States	Face mask use and hand hygiene	Influenza, respiratory syncytial virus, parainfluenza,	Three-arm RCT	Households	509	Not reported	52%	54%	Participants' own homes	No : (e

			enterovirus, rhinovirus, adenovirus, metapneumovirus								
19) Little (2015)	United Kingdom	Hand hygiene	Influenza	Two-arm RCT	Community-dwelling adults	20,066	56.6 (13.6)	56%	Not reported	Online	
20) Liu (2019)	China	Hand hygiene	Respiratory viral infections	Single arm, pre- and post-intervention study	Kindergarten teachers	361	29.0 (8.7)	95%	Not reported	Schools	1
21) MacIntyre (2009)	Australia	Face mask use and hand hygiene	Influenza, respiratory syncytial virus, adenovirus, parainfluenza virus, coronavirus, metapneumovirus	Three-arm, cluster RCT	Households with an index paediatric patient	145 families, 290 caregivers	Not reported	Not reported	Not reported	Not reported	c
22) Mott (2007)	United States	Hand hygiene	Respiratory illnesses (e.g. flu, colds)	Three-arm, non-randomised cohort study	Military trainees	2728	20.2 (-)	0%	Unclear	US army training centre	1
23) Nandrup-Bus (2009)	Denmark	Hand hygiene	Respiratory infections	Two-arm, cluster RCT	School children	652	Not reported	51%	0%	Schools	1 : : :
24) Or (2020)	China	Hand hygiene	Influenza	Single arm, pre- and post-intervention study	Households with a child in kindergarten	58 parents; 60 children	Not reported	Parents = 93%; children = 40%	Not reported	Not reported	L p k : : :
25) Ram (2015)	Bangladesh	Hand hygiene	Influenza	Two-arm, cluster RCT	Households	377 index cases; 384 household compounds	121.2 months (181.7)	40%	Not reported	Participants' own homes	
26) Reyes Fernández (2015)	Costa Rica	Hand hygiene	Respiratory infections	Two-arm, cluster RCT	University students	242	21.0 (3.9)	61%	100%	University classrooms	1
27) Roberts (2000)	Australia	Hand hygiene	Respiratory infections	Two-arm, cluster RCT	Pre-school children and day care staff	232 childcare centres; 558 children	Not reported	Not reported	0%	Childcare centres	c
28) Sandora (2005)	United States	Hand hygiene	Respiratory infections	Two-arm, cluster RCT	Households with a child	292 families	36.7 (9.5)	Not reported	91%	Participants' own homes	c

					at a day care centre						
29) Savolainen-Kopra (2012)	Finland	Hand hygiene	Respiratory infections	Three-arm, cluster RCT	Office workers	683	Not reported	Not reported	Not reported	Office buildings	
30) Simmerman (2011)	Thailand	Face mask use and hand hygiene	Influenza	Three-arm RCT	Households	442 index cases; 1147 household contacts	Not reported	59%	Not reported	Participants' own homes	
31) Stebbins (2010)	United States	Hand hygiene	Influenza and related respiratory infections	Two-arm, cluster RCT	School children and teachers	151 teachers	Not reported	Not reported	Not reported	Schools	
32) Stedman-Smith (2015)	United States	Hand hygiene	Respiratory infections	Two-arm, cluster RCT	Office workers	324	Not reported	84%	100%	Office buildings	
33) Suess (2011)	Germany	Face mask use and hand hygiene	Pandemic influenza A (H1N1)	Three-arm, cluster RCT	Households with an index patient during the H1N1 pandemic	147	Index cases = 7.9 (3.3); household contacts = 30.0 (14.2)	52%	Not reported	Participants' own homes	
34) Suess (2012)	Germany	Face mask use and hand hygiene	Pandemic influenza A (H1N1)	Three-arm, cluster RCT	Households with an index patient during the H1N1 pandemic	302	Not reported	Not reported	Not reported	Participants' own homes	
35) Updegraff (2011)	United States	Hand hygiene	Pandemic influenza A (H1N1)	Randomised cross-over study	University students and staff	65 units	Not reported	Not reported	Not reported	Public areas of a university	
36) White (2003)	United States	Hand hygiene	Respiratory infections	Two-arm, cohort study*	University students	430	18.3 (0.7)	62%	0%	University residence halls	
37) Yardley (2011)	United Kingdom	Hand hygiene	Respiratory infections	Two-arm, pilot RCT	Community-dwelling adults	517	49.8 (11.4)	64%	Not reported	Online	
38) Zomer (2016)	The Netherlands	Hand hygiene	Respiratory infections	Two-arm, cluster RCT	Households with a child at a day care centre	71 centres	Not reported	Not reported	Not reported	Not reported	
39) Öncü (2019)	Turkey	Hand hygiene	Respiratory infections	Three-arm, cluster RCT	School children	96	9.2 (1.0)	54%	0%	School laboratory	

Note. \* Unclear if randomised.

Table 3. Characteristics of interventions to change personal protective behaviours.

Lead author (year)	Comparator	Intervention	Intervention duration	BCTs (comparator arm)	BCTs (intervention arm)	Intervention mode of delivery	Theoretical mechanism(s) of action of the intervention	Incentive structure for study participation
1) Aiello (2010)	Educational materials	The face mask group received face masks; written instructions on how to use, store and safely discard masks. The face mask + hand hygiene group also received hand sanitizer	6 weeks	4.1. Instruction on how to perform the behaviour	4.1. Instruction on how to perform the behaviour; 6.1. Demonstration of the behaviour; 8.1. Behaviour practice/rehearsal; 12.5. Adding objects to the environment	Video link, written materials	Not reported	Those with influenza-like illness were offered \$25 for providing a throat specimen
2) Aiello (2012)	Educational materials	The face mask group received face masks; written instructions on how to use, store and safely discard masks. The face mask + hand hygiene group also received hand sanitizer	6 weeks	4.1. Instruction on how to perform the behaviour	4.1. Instruction on how to perform the behaviour; 8.1. Behaviour practice/rehearsal; 12.5. Adding objects to the environment	Written materials	Not reported	Not reported
3) Apisarnthanarak (2009)	NA	Children, teachers and parents received hand hygiene (including sanitizer) education via cartoons and workshops. A single dispenser of alcohol-based hand rub was placed in each nursery room	3 years	NA	2.2 Feedback on behaviour; 2.7 Feedback on outcomes of behaviour; 4.1. Instruction on how to perform the behaviour; 6.1. Demonstration of the behaviour; 8.1. Behaviour practice/rehearsal; 12.5. Adding objects to the environment	Face-to-face workshop, cartoons, written materials	Not reported	Not reported
4) Arbogast (2016)	An educational video about hand hygiene; soap and hand sanitizer provided in toilets	Same as the control group in addition to hand sanitizers being provided in different areas of the office building	13.5 months	4.1. Instruction on how to perform the behaviour; 12.5. Adding objects to the environment	4.1. Instruction on how to perform the behaviour; 6.1. Demonstration of the behaviour; 12.5. Adding objects to the environment	Videos	Not reported	25 'wellness points' were offered to employees who completed both the baseline and post-study survey

5) Azman (2013)	Hand sanitizer	Same as control group in addition to a live demonstration of hand washing behaviour. Information about hand hygiene was sent home	Not reported	12.5. Adding objects to the environment	4.1. Instruction on how to perform the behaviour; 6.1. Demonstration of the behaviour; 12.5. Adding objects to the environment	Face-to-face workshop, written materials	Not reported	Not reported
6) Azor-Martinez (2016)	No intervention	Handwashing workshop; hand hygiene practices were periodically reinforced in the classroom; younger children were supervised during hand hygiene procedures; provision of hand sanitizer	8 months	NA	4.1. Instruction on how to perform the behaviour; 6.1. Demonstration of the behaviour; 8.1. Behaviour practice/rehearsal; 12.5. Adding objects to the environment	Face-to-face workshop	Not reported	Not reported
7) Azor-Martinez (2018)	No intervention	Hand hygiene workshop with instruction on how to correctly perform the behaviour; use of stories, songs, posters; provision of hand sanitizer in one group and liquid soap in the other group; written materials on hand hygiene	8 months	NA	4.1. Instruction on how to perform the behaviour; 6.1. Demonstration of the behaviour; 8.1. Behaviour practice/rehearsal; 12.5. Adding objects to the environment	Face-to-face workshop, written materials	Not reported	Not reported
8) Barasheed (2014)	Hygiene information	In addition to the general hygiene information, face masks and written and verbal instructions on how to use these were provided	5 days	Not reported	4.1. Instruction on how to perform the behaviour; 12.5. Adding objects to the environment	Face-to-face instructions, written materials	Not reported	Not reported
9) Bundgaard (2020)	Weekly e-mails encouraging participants to follow current COVID-19 recommendations	Instructions to wear a mask when outside the home during the next month; provision of 50 three-layer, disposable, surgical face masks with ear loops	1 month	NA	4.1 Instruction on how to perform the behaviour; 12.5 Adding objects to the environment	Written materials	Not reported	Not reported
10) Canini (2010)	No intervention	Provision of face masks and demonstration of how to use them	3 weeks	NA	6.1. Demonstration of the behaviour; 12.5. Adding objects to the environment	Face-to-face instructions	Not reported	Not reported
11) Chan (2007)	NA	Health education	7 days	NA	Not reported	Telephone calls with trained nursing students	Not reported	Not reported
12) Cowling (2008)	No intervention (education about healthy diet and lifestyle)	The face mask group received face masks, information about the efficacy of masks and	9 days	NA	4.1. Instruction on how to perform the behaviour; 5.1. Information about	Not reported	Not reported	At the final home visit, households were



		instruction on how to use and safely dispose of masks; the hand hygiene group received hand sanitizer, liquid soap, information about the efficacy of hand hygiene and demonstration of hand hygiene behaviour			health consequences; 6.1. Demonstration of the behaviour; 12.5. Adding objects to the environment;			reimbursed for their participation with a supermarket voucher worth approximately US \$20
13) Cowling (2009)	No intervention (education about healthy diet and lifestyle)	Hand hygiene; surgical face masks plus hand hygiene	6 days	NA	5.1. Information about health consequences; 12.5. Adding objects to the environment	Not reported	Not reported	At the final home visit, households were reimbursed for their participation with a supermarket voucher worth approximately US \$25
14) Hübner (2010)	No intervention	Provision of hand sanitizer and instruction on how and when to use it at work	12 months	NA	4.1. Instruction on how to perform the behaviour; 12.5. Adding objects to the environment	Not reported	Not reported	Not reported
15) Kaewchana (2012)	30-minute routine health education on influenza infection, nutrition, physical activity, and smoking cessation	30-minute intensive hand washing education; individual training on hand washing; provision of liquid soap; self-monitoring diary; written materials on hand washing techniques	3 months	4.1. Instruction on how to perform the behaviour; 5.1. Information about health consequences	2.3. Self-monitoring of behaviour; 4.1. Instruction on how to perform the behaviour; 5.1. Information about health consequences; 5.3. Information about social and environmental consequences; 8.1. behaviour practice/rehearsal; 12.5. Adding objects to the environment	Face-to-face workshop	Not reported	Not reported
16) Koep (2016)	No intervention	Educational intervention with information about microorganisms; provision of liquid soap and hand sanitizer in toilets and classrooms	Unclear	NA	4.1. Instruction on how to perform the behaviour; 5.1. Information about health consequences; 12.5. Adding objects to the environment	Face-to-face with trained teachers	Not reported	Not reported

17) Larson (2009)	NA	Educational intervention focused on infection control; question and answer fact sheets; two groups of households were randomised to receive hand sanitizer, face masks or both	2 to 20 months	NA	4.1. Instruction of how to perform the behaviour; 5.1. Information about health consequences; 8.1. Behaviour practice/rehearsal; 12.5. Adding objects to the environment	Face-to-face with trained researchers	Not reported	Not reported
18) Larson (2010)	Educational materials on infection control	The hand sanitizer group received education plus hand sanitizer; the hand sanitizer and face mask group received the same interventions plus face masks	19 months	Not reported	4.1. Instruction on how to perform the behaviour; 5.1. Information about health consequences; 8.1. behaviour practice/rehearsal; 12.5. Adding objects to the environment	Face-to-face with trained researchers	Not reported	Not reported
19) Little (2015)	No intervention	Four weekly web-based sessions with new content focused on the role of hand washing, setting up a plan to wash hands, reinforcement of helpful attitudes and norms, addressing negative beliefs, tailored feedback and prompts to login to the website	4 months	NA	1.4. Action planning; 2.2. Feedback on behaviour; 4.1. Instruction on how to perform the behaviour; 7.1. Prompts/cues; 8.1. Behaviour practice/rehearsal	Website	Not reported	Not reported
20) Liu (2019)	NA	Hand hygiene training and information booklet; provision of soap, towels, posters, stickers, books, memory games and diplomas	6 months	NA	4.1 Instruction on how to perform the behaviour; 10.4 Social reward; 12.5 Adding objects to the environment	Face-to-face sessions, posters, stickers, books	Knowledge, perceived susceptibility, perceived severity, perceived behavioural control	Not reported
21) MacIntyre (2009)	No intervention	Information about infection control; provision of either P2 or surgical face masks	Unclear	NA	12.5 Adding objects to the environment	Not reported	Not reported	Not reported
22) Mott (2007)	Hand sanitizer and instructions to wash or sanitize hands after key events (e.g. coughing, sneezing)	In the PI group, hand sanitizer dispensers were installed throughout the training environment; provision of personal hand sanitizer bottles; posters were placed in training facilities to encourage	Not reported	12.5 Adding objects to the environment	4.1 Instruction on how to perform the behaviour; 9.1 Credible source; 12.5 Adding objects to the environment	Face-to-face sessions, posters	Not reported	Not reported

		hand hygiene and foster a sense of pride in staying healthy; instruction on hand hygiene; weekly reminders by drill sergeants to carry, use and refill hand sanitizer bottles. The SI group received personal hand sanitizer bottles and instruction on hand hygiene only						
23) Nandrup-Bus (2009)	NA	Children were required to wash their hands three times per day; training on hand washing and infection control was provided; posters with step-by-step hand washing instructions placed by wash basins; parents were asked to remind children to wash their hands before the first lesson each day	3 months	NA	4.1 Instruction on how to perform the behaviour; 5.1 Information about health consequences	Face-to-face sessions, posters	Not reported	Not reported
24) Or (2020)	No intervention	Children attended 4 weekly sessions with information about infection control and hand hygiene techniques; parents attended a separate session with similar content	4 weeks	NA	4.1 Instruction on how to perform the behaviour; 5.1 Information about health consequences	Face-to-face sessions with an infection control nurse	Knowledge, skills	Not reported
25) Ram (2015)	No intervention	Household compounds were provided with a hand washing station (e.g. water container with a tap, soap); information on infection control and skills training; cue cards placed in a common area in compound courtyards	Tailored; daily intervention visits until 10 days following the resolution of the index case patient's symptoms	NA	4.1 Instruction on how to perform the behaviour; 5.1 Information about health consequences; 12.5 Adding objects to the environment	Face-to-face sessions, cue cards	Not reported	Not reported
26) Reyes Fernández (2015)	No intervention	Instructions on how and when to clean hands and a planning task to help students set action and coping plans	One-off	NA	1.2 Problem solving; 1.4 Action planning; 4.1 Instruction on how to perform the behaviour	A face-to-face session with research assistants, pamphlets	Action control, coping planning	Not reported

27) Roberts (2000)	No intervention	Staff received training in hand washing and were asked to teach the techniques to the children in their care via songs about hand washing to the melodies of nursery rhymes; training was reinforced with fortnightly visits and newsletters	Not reported	NA	4.1 Instruction on how to perform the behaviour; 6.1 Demonstration of the behaviour	Face-to-face training sessions, newsletters	Not reported	Not reported
28) Sandora (2005)	No intervention (educational materials about healthy eating; participants were asked not to use hand sanitizer during the study period)	Provision of alcohol-based hand sanitizer; educational materials (e.g. fact sheets, games, toys) about hand hygiene	5 months	NA	12.5 Adding objects to the environment	Fact sheets, games, toys	Not reported	Not reported
29) Savolainen-Kopra (2012)	No intervention	Both groups received information on infection control. In the soap and water group, toilets were equipped with liquid hand soap. In the hand sanitizer arm, toilets were equipped with both liquid hand soap and alcohol-based hand rub	18 months	NA	4.1 Instruction on how to perform the behaviour; 5.1 Information about health consequences; 12.5 Adding objects to the environment	Not reported	Not reported	Not reported
30) Simmerman (2011)	No intervention (nutritional, physical activity, and smoking cessation education)	The hand washing group received education, instruction on hand washing techniques and a hand washing kit with liquid hand soap. The hand washing + face mask group received the same interventions as the hand washing group in addition to paper surgical face masks, training on how to use them appropriately and information about benefits of use	7 days	NA	4.1 Instruction on how to perform the behaviour; 5.1 Information about health consequences; 12.5 Adding objects to the environment	Face-to-face sessions with trained study nurses	Not reported	Households were compensated with approximately US \$60 in Thai baht
31) Stebbins (2010)	No intervention	Students and staff received training in hand hygiene behaviors; schools placed and maintained supplies of alcohol-based hand sanitizer in all classrooms and common areas;	Not reported	NA	4.1 Instruction on how to perform the behaviour; 5.1 Information about health consequences; 12.5	Face-to-face sessions, videos	Not reported	Teachers were offered a \$5 gift card for completion of each survey

		parents and guardians received educational materials on hand hygiene and home isolation practices			Adding objects to the environment			
32) Stedman-Smith (2015)	A brief training video to promote more effective communication with health care providers, branded key chains, brochures and posters with information about the programme	A brief training video on infection control and demonstration of effective hand washing/gel techniques; hand sanitizer and motivational/educational hand hygiene posters were placed in break rooms, kitchens and conference rooms	Not reported	Not reported	4.1 Instruction on how to perform the behaviour; 12.5 Adding objects to the environment	Video, posters	Not reported	Points toward health promotion items within an employee wellness program were provided for each survey completed
33) Suess (2011)	Educational materials about infection control and recommendation to sleep in a different room than the index patient	Participants in the mask + hand hygiene and mask groups were provided with surgical face masks with ear loops and written information on their correct use; participants in the mask + hand hygiene group were provided with alcohol-based hand rub and instructions on correct use	8 days	Not reported	4.1 Instruction on how to perform the behaviour; 12.5 Adding objects to the environment	Written materials, telephone, face-to-face visits by trained study personnel	Not reported	€150 for study participation
34) Suess (2012)	Educational materials about infection control and recommendation to sleep in a different room than the index patient	Participants in the mask + hand hygiene and mask groups were provided with surgical face masks with ear loops and written information on their correct use; participants in the mask + hand hygiene group were provided with alcohol-based hand rub and instructions on correct use	8 days	Not reported	4.1 Instruction on how to perform the behaviour; 6.1 Demonstration of the behaviour; 12.5 Adding objects to the environment	Written materials, telephone, face-to-face visits by trained study personnel	Not reported	€150 for study participation
35) Updegraff (2011)	Hand sanitizer	Four signs were placed above hand sanitizer units. The perceived susceptibility headline read "Germs are out to get you. Get them first!"; the social norms headline read "Everybody is doing it. Are you?"; the gain-framed headline read "Stay healthy this season."	12 weeks	12.5 Adding objects to the environment	5.1 Information about health consequences; 5.2 Information about social and environmental consequences; 12.5 Adding objects to the environment	Foam boards	Perceived susceptibility, social norms, and attitudes toward the behavior	Not reported

		Sanitize your hands”; and the loss-framed headline read “H1N1. Getting it is as easy as passing me by.” Each sign contained a “fact box” with more detailed information reinforcing the theme						
36) White (2003)	No intervention	Alcohol-gel dispensers were installed in every room, washroom, and dining hall in residence halls; a hand washing message campaign was implemented with bulletin boards and weekly messages to encourage hand washing	10 weeks	NA	12.5 Adding objects to the environment	Bulletin boards	Not reported	Cash incentives totalling a maximum of \$65
37) Yardley (2011)	No intervention	Four weekly web-based sessions with information about the medical team behind the advice (to enhance credibility), infection control, expert recommendations for hand washing frequency and technique, and instructions for picking up a free supply of hand gel from one's local GP practice; a hand washing plan to promote intention formation with situational cueing; tailored feedback to help participants improve their plan; reinforcement of positive attitudes and norms; addressing common negative beliefs; e-mail prompts to login to the website	4 weeks	NA	1.4 Action planning; 2.2 Feedback on behaviour; 5.1 Information about health consequences; 7.1 Prompts/cues; 9.1 Credible source; 12.5 Adding objects to the environment	Website	Intention to wash hands, attitude, subjective norms, perceived behavioural control, perceived risk of infection	Not reported
38) Zomer (2016)	Usual care	Four components: 1) free hand hygiene products with refills for 6 months (e.g. soap, hand sanitizer); 2) a handwashing exercise with 'UV Glow Cream' and an information booklet with the training content; 3) two team training sessions focused on goal	6 months	Not reported	1.1 Goal setting (behaviour); 7.1 Prompts/cues; 8.1 Behavioural practice/rehearsal; 12.5 Adding objects to the environment	Face-to-face sessions, posters, stickers, booklets	Not reported	Not reported

		setting and identifying hand hygiene improvement activities; 4) posters and stickers placed in day care centres acting as reminders to practice hand hygiene						
39) Ōncū (2019)	Hand gel and 'photo shoots' before and after handwashing	The first group received the same intervention as the control group plus information about getting rid of microbes on hands if washing hands 'properly'; the second group received the same interventions as the first plus a 30-minute hand hygiene educational and training session on types of microbes and diseases caused by microbes, in addition to WHO's 9-stage handwashing programme	4 weeks	12.5 Adding objects to the environment	4.1 Instruction on how to perform the behaviour; 5.1 Information about health consequences; 8.1 Behavioural practice/rehearsal; 12.5 Adding objects to the environment	Face-to-face session	Not reported	Not reported

Note. NA = not applicable.

Table 4. APEASE criteria, reach and engagement.

Lead author (year)	Acceptability	Practicability	Affordability	Spill-over effects	Equity	Reach	Engagement	How primary outcome was assessed	Effectiveness
1) Aiello (2010)	Not reported	Not reported	Not reported	Not reported	Not reported	7/15 residence halls were included in the study	Not reported	Self-report	On average, the mask only group wore masks 3.9 hours per day (SD = 3.3) versus 3.0 hours per day (SD = 2.4) in the mask and hand hygiene group. Log mask hours were significantly higher in the mask only group compared with the mask and hand hygiene group at each time point except for week 4 ( $p's < .05$ ). The mask only group washed their hands 8.2 times per day (SD = 9.0) versus 6.1 times per day (SD = 4.8) in the mask and hand hygiene group and 8.8 times per day (SD = 9.3) in the control group. On the log scale, the mask and hand hygiene group washed their hands significantly fewer times per day than the control group from weeks 2 through 4. The mask only group used hand sanitizer 2.3 times per day (SD = 3.5) versus 5.2 times per day (SD = 5.1) in the mask and hand hygiene group and 2.0 times per day (SD = 3.9) in the control group. On



									the log scale, participants in the mask and hand hygiene group reported a significantly greater use of hand sanitizer compared to the mask only and control groups at each week ( $p's < .0001$ ). There were no significant differences between the mask only group and control group in the frequency of hand sanitizer use
2) Aiello (2012)	On average, participants in the two intervention groups rated mask comfort as 4.7/10 (SD = 0.2)	Not reported	Not reported	Not reported	Not reported	5/15 residence halls were included in the study	Not reported	Self-report and observation of mask wearing by trained staff	Participants in the face mask and hand hygiene group and the face mask only group wore masks on average 5.1 hours per day (SD = 2.2) and 5.0 hours per day (SD = 2.2), respectively ( $p > .05$ ). The face mask and hand hygiene group and the face mask only group reported an average use of hand sanitizer 4.5 times per day (SD = 4.1) and 1.3 times per day (SD = 1.8), respectively ( $p < .05$ ). This compares with 1.5 times per day (SD = 2.3) in the control group. The face mask and hand hygiene group and face mask only group washed their hands 5.2 (SD = 3.3) and 5.5 (SD = 3.3) times per day,

									respectively. The control group washed their hands an average of 5.8 times (SD = 5.0) per day (p>.05)
3) Apisarnthanarak (2009)	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Observation of hand hygiene and cough etiquette with a Web camera 1 hour, twice weekly	The observed rate of hand hygiene compliance increased from 21% (50/240) in period 1 to 71% (170/240) in period 2 (p<.001) and to 69% (166/240) in period 3 (p<.001)
4) Arbogast (2016)	Employees in the intervention group were significantly more likely than those in the control group to have a positive impression of the programme because of the presence of alcohol-based hand sanitizers in the workplace (80% vs 69%, p < 0.001). 88% of employees in the intervention group reported liking the products provided	Not reported	Not reported	Not reported	Not reported	1386/1609 employees agreed to participate	Not reported	Self-report	Self-reported hand washing frequency improved significantly over time in the intervention group (p<.05), as did self-reported hand sanitizer use for every activity assessed, including before eating, after sneezing, coughing, handling money, using the restroom, returning to their desk, and interacting with others who may be sick (p's<.05)
5) Azman (2013)	Not reported	Not reported	Not reported	Not reported	Age and ethnicity were significant risk factors for self-reported influenza-like illness (p<.05)	Not reported	Not reported	Self-report	Not reported
6) Azor-Martinez (2016)	One child showed worsening of existing atopic dermatitis due to hand sanitizer	Not reported	Not reported	Not reported	Predictors of a lower rate of absenteeism due to respiratory	1616/1640 children were randomised	Not reported	Self-report	Not reported

	gel use and was excluded from the study				illness included older age, higher parental income, and correct handwashing technique (p<.05)				
7) Azor-Martinez (2018)	One child showed worsening of existing localized atopic dermatitis due to hand sanitizer gel use and was excluded from the study	Not reported	Not reported	Not reported	Not reported	25/52 day care centres contacted were randomised	Not reported	Self-report	Not reported
8) Barasheed (2014)	The most commonly reported reason for not wearing face masks was discomfort (15% of participants)	Not reported	Not reported	Not reported	Not reported	164/4200 pilgrims took part	Not reported	Self-report	Compliance with face mask use was 76% (56/75) in the intervention group and 12% (11/89) in the control group (p<.001)
9) Bundgaard (2020)	Not reported	Not reported	The authors note that costs and availability may reduce the efficacy of face masks to prevent SARS-CoV-2 infection	Not reported	Not reported	17,258/6304 of those responding to recruitment adverts were randomised	Not reported	Self-report	46% of participants in the intervention arm wore the mask as recommended, 47% predominantly as recommended, and 7% not as recommended. Participants used an average of 1.7 masks per weekday and 1.3 per weekend day
10) Canini (2010)	75% of participants in the intervention arm reported discomfort with mask use. The three main causes of discomfort were warmth (45%), respiratory	Not reported	Not reported	Not reported	Not reported	95/105 randomised households completed the study		Self-report	Index patients in intervention households reported wearing a total of 11.0 (SD = 7.2) masks during 4.0 (SD = 1.6) days with an average use of 2.5 (SD = 1.3) masks per day and a duration of use of

	difficulties (33%) and humidity (33%). Children wearing child size face masks reported feeling pain more frequently than those wearing adult face masks (p=.036)								3.7 (SD = 2.7) hours per day
11) Chan (2007)	Not reported	Not reported	Not reported	Not reported	Not reported	182/295 registered members were successfully contacted; 122/182 took part	Not reported	Self-report	Significant improvements were observed with regards to washing hands after sneezing/coughing ( $M_{\text{before}} = 1.26$ , SD = 0.44; $M_{\text{after}} = 1.65$ , SD = 0.89), washing hands with liquid soap ( $M_{\text{before}} = 1.14$ , SD = 0.35; $M_{\text{after}} = 1.53$ , SD = 0.82), and wearing masks in public ( $M_{\text{before}} = 1.31$ , SD = 0.46; $M_{\text{after}} = 2.15$ , SD = 1.12) (all p's < .0001)
12) Cowling (2008)	There were no reported adverse events requiring medical attention	Not reported	Not reported	Not reported	Not reported	198/944 index patients were randomised	Not reported	Self-report	In the face mask group, 45% (10/22) of index cases reported wearing a mask often or always, compared with 30% (22/74) and 28% (9/32) in the control and hand hygiene groups, respectively. In the face mask group, 21% (14/65) of household contacts reported wearing a mask often or always, compared with 1% (2/213) and 4% (4/92) in the control and hand hygiene groups,

									<p>respectively. Index cases in the face mask group used a median of 12 masks (IQR = 6-18) whereas household contacts used a median of 6 masks (IQR = 1-20). 63% (41%) of index cases (household contacts) in the hand hygiene group reported washing their hands often or always after sneezing, coughing or cleaning their nose, compared with 63% (47%) and 31% (27%) in the face mask and control groups, respectively. In the hand hygiene group, households used a median of 56 g (IQR = 27-93) of alcohol from the automatic sanitizer, and a median of 88 g (IQR = 63-149) of liquid hand soap over the course of the study</p>
13) Cowling (2009)	Not reported	Not reported	Not reported	Not reported	Not reported	4,07/2,750 index patients and their households were randomised	Not reported	Self-report	<p>The proportion index cases wearing face masks was 15% in the control group (14/91); 31% in the hand hygiene group (26/85); and 49% (41/83) in the face mask and hand hygiene group. The proportion household contacts wearing face masks was 7% (20/279) in</p>

									the control group; 5% (13/257) in the hand hygiene group; and 26% (67/258) in the face mask and hand hygiene group. The median (IQR) use of liquid soap in the hand hygiene group was 77.6 g (42.4-162.6) and 78.9 (35.2-114.2) in the face mask and hand hygiene group. The median (IQR) hand rub used in the hand hygiene group was 3.2 g (1.1-9.7) in index cases and 1.5 g (0.3-5.3) in contacts. This compared with 1.6 g (0.7-5.1) in index cases and 1.5 g (0.3-3.8) in contacts in the face mask and hand hygiene group
14) Hübner (2010)	Not reported	Not reported	Not reported	Not reported	Not reported	134/850 participants were randomised	Not reported	Self-report	The mean hand disinfection frequency was >5 times daily in 19%, 3-5 times daily in 60%, and 1-2 times daily in 21%
15) Kaewchana (2012)	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Self-report	On day 7, the control and intervention groups reported 3.9 (SD = 2.4) and 5.7 (SD = 3.4) hand washing episodes/day, respectively (p<.001). The percentage of participants who used soap increased from 34% (53/158) to

									88% (139/158) (p<.001)
16) Koep (2016)	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Self-report and objective measure of soap and sanitizer use	No significant differences in hand soap compliance were observed across groups (estimates not reported). Hand sanitizer use increased slightly in the intervention school from 0.04 to 0.06 times per student per day in grades 3 and 4 (p's>.05)
17) Larson (2009)	Not reported	Not reported	Not reported	Hand sanitizer use increased but hand washing with soap decreased	Respondents with college degrees had higher knowledge scores than the other groups, adjusting for baseline scores (p = .04)	Not reported	Not reported	Self-report (questionnaire administered by trained researchers) and objective measures of hand sanitizer and mask use	A significantly greater proportion of participants reported using hand sanitizer at the end of the study (282/422) compared with baseline (6/422), but a lower proportion reported using antibacterial soap at the end of the study (105/422) compared with baseline (191/422)
18) Larson (2010)	Not reported	Not reported	Not reported	Not reported	Not reported	617/672 households that expressed interest in participation met eligibility criteria; 509/617 completed the initial home visit	Not reported	Self-report (questionnaire administered by trained researchers) and objective measures of hand sanitizer and mask use	Participants in the hand sanitizer group used a mean of 12.1 ounces/month and those in the hand sanitizer and face mask group used a mean of 11.6 ounces/month (p=.36). Half of the households with a case of infection reported using masks within 48 hours of symptom onset. Those who used masks

									reported a mean of two masks/day/episode (range: 0–9)
19) Little (2015)	Minor self-reported skin irritation increased among those who did not report problems at baseline. Among individuals who had a skin complaint at baseline, reported skin complaints did not significantly increase over time	Not reported	Not reported	Not reported	Not reported	20,066/80,4897 who received a mailed invitation were randomised	Not reported	Self-report	Not reported
20) Liu (2019)	Not reported	Not reported	Not reported	Not reported	Not reported	12/213 kindergarten clusters received the intervention	Not reported	Self-report	The average self-reported compliance with hand hygiene guidelines was significantly greater after the intervention (9.7, IQR = 0.23) compared with baseline (9.4, IQR = 0.47), $p < .01$ . Teachers reported significantly more hand washing behavior compared with baseline after coughing/sneezing, blowing their nose, changing a diaper, contacting bodily fluids and soiled textiles, going to the toilet, wiping the nose of a child, wiping a child's bottom, and before helping a child with food ( $p$ 's $< .05$ )
21) MacIntyre (2009)	There were no significant	Fit testing for P2 masks	Not reported	Not reported	Not reported	14,5/401 families assessed for	Not reported	Self-report	On day 1 of mask use, 38% (36/94) of



	<p>differences in difficulties with mask use between the P2 and surgical mask groups, but &gt;50% reported concerns, the main one being that wearing a face mask was uncomfortable. Other concerns were that the child did not want the parent wearing a mask.</p> <p>Some participants mentioned that the mask did not fit well and that it was not practical to wear at mealtime or while asleep</p>	<p>was not conducted because this was judged unlikely to be feasible in the general community during a pandemic</p>				eligibility took part			<p>the surgical mask users and 46% (42/92) of the P2 mask users stated that they were wearing the mask “most or all of the time” (p=.37). Adherence dropped to 31% (29/94) and 25% (23/92), respectively, by day 5 of mask use</p>
22) Mott (2007)	Not reported	Not reported	Not reported	<p>Hand sanitizer use increased in leaders in both intervention groups (from 3.0 to 13.4 times/day and from 3.2 to 4.7 times/day, respectively)</p>	Not reported	Not reported	Not reported	Self-report	<p>Post-intervention, there was a decrease in the daily frequency of hand washing in the SI group (from 4.4 to 2.6 times/day) and no change in the PI group (from 4.9 to 5.0 times/day). Hand sanitizer use increased in both intervention groups (from 3.7 to 10.4 times/day and from 4.0 to 6.0 times/day, respectively)</p>
23) Nandrup-Bus (2009)	Three children withdrew from the intervention arm due to skin problems	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Objective measure of soap use	<p>School records showed a usual monthly consumption of 2–2.5 litres of liquid soap at the intervention</p>

									school. During the intervention period, consumption increased to 16 L of liquid soap (but accurate monthly measurement was not possible as soap was continuously replenished). No reliable measurement of soap consumption in the control school was available
24) Or (2020)	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Self-report and observation of hand washing with fluorescent stain gel and photos	After the programme, the percentages of properly washed areas on both hands increased significantly, in particular the wrists (from 0.5% to 82%), $p < .001$
25) Ram (2015)	Not reported	Not reported	Not reported	Not reported	Not reported	377/766 eligible index cases took part	Not reported	The intervention staff weighed the soap each day and replaced it if the bar weighed <20 grams	A median per capita soap consumption of 2.3 g (IQR = 1.7 to 3.2) was observed in the first 12 days of the programme
26) Reyes Fernández (2015)	Not reported	Not reported	Not reported	Not reported	Not reported	242/440 students completed the study	Not reported	Self-report	Self-reported frequency of hand sanitizer use per day, measured on a 5-point Likert scale, increased in both the control (from $M = 1.5$ , $SD = 0.9$ to $M = 1.7$ , $SD = 1.3$ ) and intervention groups (from $M = 1.8$ , $SD = 1.3$ to $M = 2.1$ , $SD = 1.4$ ), $p = .09$

27) Roberts (2000)	Not reported	Not reported	Not reported	Not reported	Not reported	23/26 eligible childcare centres took part	Not reported	Observation by trained researchers	Hand washing compliance in children was divided into 3 groups, corresponding to intervention centers with a score of low (53%-69%; 4 centres), moderate (70%-79%; 4 centres), and high (>80%; 3 centres) compliance
28) Sandora (2005)	Forty-five families reported 112 adverse events related to hand sanitizer use. Seventy-one (63%) of the adverse events were in relation to "dry skin," and 20 (18%) were related to "irritation." Other reported adverse events included "stinging", "smells bad", "dislike it", "allergic reaction", and "too slippery"	Not reported	Not reported	Not reported	Not reported	292/647 families assessed for eligibility took part	Not reported	Self-report	Primary caregivers reported using the hand sanitizer with a median frequency of 5.2 times/day
29) Savolainen-Kopra (2012)	Not reported	Not reported	Not reported	Not reported	Not reported	683/1270 employees considered for eligibility took part	Not reported	Assessment by study nurses	The average amount of soap or disinfectant use per participant was 6.1 g and 6.9 g in the active groups
30) Simmerman (2011)	Not reported	Not reported	Not reported	Not reported	Not reported	465/20,537 paediatric outpatients with influenza-like illness took part	Not reported	Self-report	Participants in the hand washing group reported 4.7 washing episodes/day, compared with 4.9 times/day in the hand washing plus face mask arm and

									3.9 times/day in controls ( $p=.001$ ). Participants in the face mask arm used an average of 12 masks per person, per week (median = 11, IQR = 7–16) and reported wearing face masks a median of 211 minutes/day (IQR = 17–317). Parents wore their masks for a median of 153 (IQR = 40–411) minutes per day, which was greater than other relations (median = 59; IQR = 9–266), index cases themselves (median = 35; IQR = 4–197), or their siblings (median = 17; IQR = 6–107)
31) Stebbins (2010)	Not reported	Not reported	Not reported	Not reported	Not reported	151/167 enrolled teachers took part	Not reported	Self-report	The proportion of students washing their hands more than 3 times per day in the post-flu season was significantly greater in the intervention arm ( $M = 3.7$ ) compared with the control arm ( $M = 3.4$ ), $p=.04$
32) Stedman-Smith (2015)	Not reported	Not reported	Not reported	Not reported	Not reported	324/1708 enrolled office workers took part	Not reported	Self-report	In both the intervention and control group, >80% of employees reported hand sanitizer use at least 25% of the time ( $p>.05$ )
33) Suess (2011)	The majority (60%) of participants did not report any	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Self-report	81% of index cases and 71% of household members from the

	<p>problems when wearing face masks. Of those who reported having removed their masks in transmission-prone situations, 'feeling hot' was the main reason. Other problems mentioned less frequently were pain when wearing the mask and shortness of breath. The majority of adult household contacts in the control and intervention groups perceived wearing face masks as well as intensified hand hygiene as an effective means of preventing transmission of influenza</p>								<p>combined active groups wore a mask 'always' or 'most of the time' when in the same room with either a healthy or infected person. Participants (index cases and contacts combined) in the mask plus hygiene group washed/disinfected their hands significantly more frequently (51/56) compared with those in the mask and control groups combined (59/89) (p=.007)</p>
34) Suess (2012)	<p>The majority of participants (62%) did not report any problems with mask wearing. This proportion was significantly higher in adults (71%) than children (50%) (p = 0.005). The main problem stated by participants (adults as well as children) was "heat/humidity", followed by "pain" and "shortness of</p>	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	<p>The amount of remaining intervention materials was assessed at the end of the study period</p>	<p>Participants in the mask group used a median of 12.9 (IQR = 9.5-16) face masks per individual. Participants in the mask plus hygiene group used a median of 12.6 (IQR = 7.8-14) face masks. The number of hand disinfections per day was 7.4 and 4.1 in index cases in the 2009/10 and 2010/11 cohorts, respectively. The number of hand disinfections per</p>

	breath" when wearing a face mask								day was 8.8 and 7.5 in household contacts in the 2009/10 and 2010/11 cohorts, respectively
35) Updegraff (2011)	There were no significant differences between the four signs in how easy they were to understand. The loss-framed and perceived susceptibility signs elicited significantly more negative affect than the gain-framed and norms signs ( $p$ 's $< .01$ ).	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Grams of sanitizer used per day, measured with a digital scale	All signs resulted in significantly greater hand sanitizer use compared with the no sign condition, but they were not equally effective. Dispensers with the gain-framed signs had the greatest hand sanitizer use, with 66.4% more use than dispensers with no signs ( $p < .001$ ). Loss-framed signs were associated with a 58.4% increase in use compared with no sign ( $p < .001$ ). The social norms signs (44.3% increase) and the perceived susceptibility signs (40.6% increase) were associated with somewhat lower increases in usage compared with the gain-framed and loss-framed signs, but both led to significantly more use than no sign (both $p$ 's $< .01$ ).
36) White (2003)	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Self-report	Over the course of the study, the product group washed their hands 10.4% more frequently than the control group (0.5 times/h vs. 0.4

									times/h; $p < .05$ ). The frequency of hand sanitizer use was also significantly greater in the product group (0.3 times/h vs. 0.0 times/h; $p < .05$ )
37) Yardley (2011)	Not reported	Not reported	Not reported	Not reported	Moderator analyses indicated that the intervention was similarly effective for those from higher and lower socioeconomic status groups.	517/8150 of those invited participated	Of the 324 participants who were randomly assigned to the intervention, 251 (77.5%) progressed to the second session, 219 (67.6%) completed three sessions, and 188 (58.0%) completed all four sessions. The free hand gel was collected by 170/324 (52.5%) eligible participants.	Self-report	Significant differences between groups in hand washing frequency, measured on a 5-point scale, were observed at 4 weeks ( $M = 4.4$ , $SD = 0.9$ vs. $M = 4.0$ , $SD = 0.9$ ; Cohen's $d = 0.42$ ) and at 12 weeks ( $M = 4.1$ , $SD = 1.1$ vs. $M = 4.5$ , $SD = 0.8$ ; Cohen's $d = 0.34$ ), $p$ 's $< .001$
38) Zomer (2016)	Not reported	Not reported	Due to budget restrictions, hand hygiene products were only provided for two groups within each day care centre, even if the centre had more	No significant effect of the intervention was found on supervising children's hand hygiene (36% vs. 32%)	Not reported	71/122 day care centres participated	Of 274 caregivers, 21% (54/261) attended none of the training sessions, 25% (66/261) attended one training session, 29% (75/261)	Direct observation by trained researchers	Hand hygiene compliance in intervention day care centres was 66% vs. 43% in control centres (OR = 6.33, 95% CI = 3.71–10.80)

			than two groups.				attended two training sessions and 25% (66/261) attended all three sessions.		
39) Ōncū (2019)	Not reported	Not reported	Not reported	Not reported	Not reported	96/552 children	Not reported	Photos examined by researchers not directly involved in the study	In the most intensive intervention group, post-intervention hand washing effectiveness scores increased significantly across all regions of both hands (p<.05)

Note. + = positive effect; - = negative effect; / = no change; @ = indeterminate; NA = not applicable.

Table 5. Quality appraisal.



Lead author (year)	1) Bias arising from the randomisation process	2) Bias due to deviations from the intended interventions	3) Bias due to missing outcome data	4) Bias in measurement of the outcome	5) Bias in selection of the reported result	Overall rating
1) Aiello (2010)	Some concerns	Low risk of bias	Some concerns	Low risk of bias	Low risk of bias	Some concerns
2) Aiello (2012)	Some concerns	Low risk of bias	Some concerns	Some concerns	Low risk of bias	Some concerns
3) Apisarnthanarak (2009)	NA	NA	NA	NA	NA	NA
4) Arbogast (2016)	Some concerns	Some concerns	Low risk of bias	Low risk of bias	Some concerns	Some concerns
5) Azman (2013)	Some concerns	Some concerns	Low risk of bias	Some concerns	Some concerns	Some concerns
6) Azor-Martinez (2016)	Some concerns	Some concerns	Some concerns	Some concerns	Low risk of bias	High risk of bias
7) Azor-Martinez (2018)	Some concerns	Some concerns	Low risk of bias	Low risk of bias	Low risk of bias	Some concerns
8) Barasheed (2014)	Some concerns	Some concerns	Some concerns	Some concerns	Some concerns	High risk of bias
9) Bundgaard (2020)	Some concerns	Some concerns	Some concerns	Some concerns	Low risk of bias	High risk of bias
10) Canini (2010)	Some concerns	Some concerns	Low risk of bias	Low risk of bias	Low risk of bias	Some concerns
11) Chan (2007)	NA	Some concerns	Some concerns	Some concerns	Some concerns	High risk of bias
12) Cowling (2008)	Some concerns	Some concerns	Some concerns	Low risk of bias	Low risk of bias	Some concerns
13) Cowling (2009)	Some concerns	Some concerns	Some concerns	Low risk of bias	Low risk of bias	Some concerns
14) Hübner (2010)	Some concerns	Some concerns	Low risk of bias	Some concerns	Some concerns	High risk of bias
15) Kaewchana (2012)	Some concerns	Some concerns	Some concerns	Some concerns	Some concerns	High risk of bias
16) Koep (2016)	Some concerns	Some concerns	Some concerns	Some concerns	Some concerns	High risk of bias
17) Larson (2009)	NA	Some concerns	Some concerns	Some concerns	Some concerns	High risk of bias
18) Larson (2010)	Some concerns	Some concerns	Low risk of bias	Some concerns	Some concerns	High risk of bias
19) Little (2015)	Some concerns	Some concerns	Some concerns	Low risk of bias	Some concerns	High risk of bias
20) Liu (2019)	NA	NA	NA	NA	NA	NA
21) MacIntyre (2009)	Some concerns	Some concerns	Low risk of bias	Some concerns	Low risk of bias	Some concerns

22) Mott (2007)	NA	NA	NA	NA	NA	NA
23) Nandrup-Bus (2009)	Some concerns	Some concerns	Some concerns	Some concerns	Low risk of bias	High risk of bias
24) Or (2020)	NA	NA	NA	NA	NA	NA
25) Ram (2015)	Some concerns	Some concerns	Low risk of bias	Low risk of bias	Low risk of bias	Some concerns
26) Reyes Fernández (2015)	Some concerns	Some concerns	Some concerns	Some concerns	Low risk of bias	High risk of bias
27) Roberts (2000)	Some concerns	Some concerns	Some concerns	Low risk of bias	Some concerns	High risk of bias
28) Sandora (2005)	Low risk of bias	Some concerns	Some concerns	Some concerns	Some concerns	High risk of bias
29) Savolainen-Kopra (2012)	Some concerns	Some concerns	Some concerns	Some concerns	Low risk of bias	High risk of bias
30) Simmerman (2011)	Low risk of bias	Low risk of bias	Some concerns	Some concerns	Low risk of bias	Some concerns
31) Stebbins (2010)	Some concerns	Some concerns	Low risk of bias	Some concerns	Some concerns	High risk of bias
32) Stedman-Smith (2015)	Low risk of bias	Low risk of bias	Some concerns	Some concerns	Some concerns	Some concerns
33) Suess (2011)	Low risk of bias	Low risk of bias	Some concerns	Some concerns	Some concerns	Some concerns
34) Suess (2012)	Low risk of bias	Some concerns	Some concerns	Some concerns	Some concerns	High risk of bias
35) Updegraff (2011)	Low risk of bias	Low risk of bias	Low risk of bias	Low risk of bias	Low risk of bias	Low risk of bias
36) White (2003)	High risk of bias	Low risk of bias	Some concerns	Some concerns	Some concerns	High risk of bias
37) Yardley (2011)	Low risk of bias	Low risk of bias	Some concerns	Some concerns	Low risk of bias	Some concerns
38) Zomer (2016)	Low risk of bias	Some concerns	Low risk of bias	Some concerns	Low risk of bias	Some concerns
39) Ōncü (2019)	Some concerns	Some concerns	Low risk of bias	Low risk of bias	Some concerns	Some concerns

Note. NA = not applicable.

## Discussion

This rapid review of interventions to increase personal protective behaviours to limit the spread of respiratory viruses identified 39 studies conducted across 15 countries. The majority of interventions targeted hand hygiene and/or face mask use, with one intervention targeting the catching of droplets in tissues in addition to hand hygiene. None of the identified interventions focused on avoiding touching the T-Zone, disinfecting surfaces or maintaining physical distancing. Interventions were typically delivered in participants' own homes or in nurseries/schools, targeting children/adult household members or pre- or school children/teachers. Two- or three-arm study designs with passive comparators were typically used. The overall quality of included studies was low, with only one study rated as 'low risk of bias'. The majority of interventions had a face-to-face component and delivered a median of three BCTs; the most frequent were 'Adding objects to the environment', 'Instruction on how to perform the behaviour' and 'Information about health consequences'. Where investigated, interventions were considered acceptable by participants, with a minority reporting issues with mask wear

discomfort or skin irritation from hand hygiene products. Few studies reported the practicability, affordability, spill-over effects or equity of interventions. In a narrative synthesis, interventions targeting hand hygiene behaviour were found to have positive effects and those targeting face mask use had a mixture of positive and negative effects. Random-effects meta-analyses of a small number of studies found positive effects of interventions targeting hand hygiene behaviour and face mask use. However, between-study heterogeneity was high and the confidence interval for the pooled effect of interventions targeting face mask use was wide, partly due to the small number of studies included in the comparison.

#### *Strengths and limitations*

This review was conducted rapidly (July–December 2020) with input on the research questions and review scope from public health and behavioural science experts and lay members as part of a written stakeholder consultation. However, the pragmatic nature of this review, conducted during an ongoing pandemic, also means that it has several important limitations. First, given the expected large number of hand hygiene studies related to gastrointestinal infections, we limited the review to studies explicitly studying behaviour change in relation to respiratory viruses. However, data from interventions targeting personal protective behaviours to prevent gastrointestinal illness are likely to add to our understanding of the acceptability, effectiveness and equity of hand hygiene interventions. Second, evidence indicates that the relative importance of different personal protective behaviours may depend on properties of the specific respiratory virus and context (e.g. fomite transmission may be more pronounced for respiratory syncytial virus compared with coronaviruses) (Boone & Gerba, 2007). However, at the time of planning this rapid evidence review, little was known about SARS-CoV-2. We therefore opted for a broad scope and included interventions targeting personal protective behaviours to limit the spread of any respiratory viral infection. It was also not possible to group the results based on the specific viral infections studied as the majority of studies targeted multiple (as opposed to single) respiratory viral infections and there was little variability in the viral infections targeted. In addition, as the majority of studies targeted a host of different viruses within their interventions, this further limits the conclusions that can be drawn: the perceived susceptibility to different viruses likely differs between, for example, age groups (Rosenstock, 1974) and tailored intervention strategies may therefore be needed for younger (vs. older) adults. However, the current review was unable to address such nuanced questions due to the limited design and reporting of extant studies and the need to synthesise evidence quickly during an ongoing pandemic. Third, our electronic search was restricted to two databases, which may have limited the results, and data extraction was performed by a single reviewer, with a proportion verified by a second reviewer. Fourth, although most of identified studies were two- or three-arm RCTs, they were typically designed to study rates of respiratory infection as their primary outcome, with behaviour change outcomes less clearly reported. This hindered quantitative synthesis, with only a small number of included studies contributing to meta-analyses. Future studies specifically designed to examine the effectiveness of interventions on behavioural outcomes are needed. Fifth, in line with guidelines (Wood et al., 2015), we only coded BCTs when there was clear evidence of their presence; interventions may have included additional BCTs not documented in this review. Sixth, as this was a rapid review with limited resources, we limited our analyses of intervention content to information presented in the published papers and/or supplementary materials, and no attempts were made to contact study authors for access to detailed intervention descriptions. Seventh, as most interventions targeting multiple behaviours (e.g. hand hygiene and face mask use) did not clearly distinguish BCTs that targeted one (but not the other) behaviour, and none of the outcome evaluations considered potential behavioural dependencies (or statistical interactions), it was not possible to consider the extent of BCT overlap and/or behavioural interactions in the present review. Finally, due to the small number of studies available for meta-analysis, we were unable to group studies by, for example, population type, study setting, type of virus, type of outcome assessment, etc., which would have further improved our understanding of intervention effectiveness.

#### *Implications for policy and practice*

Although we caution against drawing firm conclusions due to the low quality of the evidence, positive effects of interventions targeting hand hygiene behaviour and face mask use were observed, with the majority of interventions providing free hand hygiene products and/or face masks to participants in addition to instructions on how to perform the behaviour and information about health consequences. As far as is practicably feasible, authorities should aim to provide free products to staff, clients and visitors during respiratory viral epidemics. The limited range of BCTs detected in published intervention descriptions may suggest a missed opportunity for harnessing techniques indicated by relevant behaviour change theory and evidence. We encourage policymakers and healthcare practitioners to work collaboratively with behavioural scientists to incorporate techniques that theory or evidence predicts are effective for enabling personal protective behaviours (Warren-Gash et al., 2012), such as techniques targeting motivational or self-regulatory processes. For example, while hand hygiene is a well-established, often private and widely accepted protective behaviour that most people have long experience with, face mask wearing is a relatively new (at least in some countries), public behaviour, where there is more debate and uncertainties about the impact of the behaviour among the public and scientists (B J Cowling et al., 2010). It is therefore important to involve behavioural scientists in the development of any new interventions targeting personal protective behaviours in the context of respiratory viral infections to help map out potential influences (e.g. social, self-regulatory) that may help or hinder the target behaviour, acknowledging that different interventions are likely needed for different behaviours.

### Avenues for future research

Findings highlight the need for evaluations of interventions to support people to avoid touching the T-Zone, disinfect surfaces, maintain physical distancing and ensure efficient ventilation. Ventilation is increasingly seen as an important personal protective behaviour but was missed from the present review as it was planned during an earlier epidemic phase when the emphasis was on viral transmission via droplets rather than aerosols (Anderson et al., 2020; Morawska & Milton, 2020). In addition, we need studies designed to detect effects on behavioural outcomes and data on the affordability and equity of interventions to increase personal protective behaviours, particularly in low- and middle-income countries. Although the provision of hand hygiene products and face masks may offset costs related to primary and secondary care or work absenteeism for those with severe respiratory viral illness, the provision of free products at scale may be prohibitively costly. Future research involving health and social care economists should evaluate the cost-effectiveness of different types of interventions to enable personal protective behaviours, including those targeting motivational and self-regulatory processes. We also need further evidence from studies evaluating interventions to improve adherence to face mask use, with unclear results observed at present. Finally, due to the small number of studies with data suitable for meta-analysis, we did not conduct moderator analyses to examine whether, for example, particular BCTs, broader content categories or the unit of randomisation (e.g. individual vs. cluster) were related to intervention effectiveness; this would be important to examine in future meta-analyses with a larger sample size. We did not consider here the use of, for example, the Theoretical Domains Framework (Cane et al., 2012) when coding the mechanisms of action of interventions; this may be useful to consider in future empirical studies and evidence reviews. Finally, although evidence generation during ongoing pandemics is challenging (with a need to balance a pragmatic approach and limited resources with scientific rigour), drawing primarily on studies conducted outside the pandemic setting to inform what behavioural interventions to implement is suboptimal. We therefore recommend that experimental studies of behavioural interventions are prioritised during future respiratory viral pandemics.

### Conclusions

This rapid review identified 39 studies across 15 countries with interventions targeting hand hygiene and/or face mask use. Positive effects of interventions targeting hand hygiene were observed, with unclear results for interventions targeting face mask use. There was a lack of evidence for interventions targeting most behaviours of interest within this review.

### Funding

OP receives salary support from Cancer Research UK (C1417/A22962).

### Declaration of interests

OP, DS, EC, LS, RW and SM have no conflicts of interest to declare.

### Acknowledgements

We gratefully acknowledge the funding listed. The funder had no final role in the study design; in the analysis and interpretation of data; in the writing of the report; or in the decision to submit the paper for publication. All researchers listed as authors are independent from the funder and all final decisions about the research were unrestricted.

### References

- Aiello, A. E., Murray, G. F., Perez, V., Coulborn, R. M., Davis, B. M., Uddin, M., Shay, D. K., Waterman, S. H., & Monto, A. S. (2010a). Mask Use, Hand Hygiene, and Seasonal Influenza-Like Illness among Young Adults: A Randomized Intervention Trial. *The Journal of Infectious Diseases*, 201(4), 491–498. <https://doi.org/10.1086/650396>
- Aiello, A. E., Murray, G. F., Perez, V., Coulborn, R. M., Davis, B. M., Uddin, M., Shay, D. K., Waterman, S. H., & Monto, A. S. (2010b). Mask Use, Hand Hygiene, and Seasonal Influenza-Like Illness among Young Adults: A Randomized Intervention Trial. *Journal of Infectious Diseases*, 201, 491–498. <https://doi.org/10.1086/650396>
- Aiello, A. E., Perez, V., Coulborn, R. M., Davis, B. M., Uddin, M., & Monto, A. S. (2012). Facemasks, hand hygiene, and influenza among young adults: A randomized intervention trial. *PLoS ONE*, 7(1). <https://doi.org/10.1371/journal.pone.0029744>
- Anderson, E. L., Turnham, P., Griffin, J. R., & Clarke, C. C. (2020). Consideration of the Aerosol Transmission for COVID-19 and Public Health. *Risk Analysis*, 40(5), 902–907. <https://doi.org/10.1111/risa.13500>
- Apisarnthanarak, A., Apisarnthanarak, P., Cheevakumjorn, B., & Mundy, L. M. (2009). Intervention with an Infection Control Bundle to Reduce Transmission of Influenza-Like Illnesses in a Thai Preschool. *Infection Control and Hospital Epidemiology*, 30(9), 1–6. <https://doi.org/10.1086/599773>

- Arbogast, J. W., Moore-Schiltz, L., Jarvis, W. R., Harpster-Hagen, A., Hughes, J., & Parker, A. (2016a). Impact of a Comprehensive Workplace Hand Hygiene Program on Employer Health Care Insurance Claims and Costs, Absenteeism, and Employee Perceptions and Practices. *Journal of Occupational and Environmental Medicine*, 58(6). <https://doi.org/10.1097/JOM.0000000000000738>
- Arbogast, J. W., Moore-Schiltz, L., Jarvis, W. R., Harpster-Hagen, A., Hughes, J., & Parker, A. (2016b). Impact of a Comprehensive Workplace Hand Hygiene Program on Employer Health Care Insurance Claims and Costs, Absenteeism, and Employee Perceptions and Practices. *Journal of Occupational and Environmental Medicine*, 58(6). <https://doi.org/10.1097/JOM.0000000000000738>
- Azman, A. S., Stark, J. H., Althouse, B. M., Vukotich, C. J., Stebbins, S., Burke, D. S., & Cummings, D. A. T. (2013). Household transmission of influenza A and B in a school-based study of non-pharmaceutical interventions. *Epidemics*, 5(4), 181–186. <https://doi.org/10.1016/j.epidem.2013.09.001>
- Azor-Martinez, E., Cobos-Carrascosa, E., Seijas-Vasquez, M. L., Fernandez-Sanchez, C., Strizzi, J. M., Torres-Alegrie, P., Santisteban-Martinez, J., & Gimenez-Sanchez, F. (2016a). Hand Hygiene Program Decreases School Absenteeism Due to Upper Respiratory Infections. *Journal of School Health*, 86(12), 873–881.
- Azor-Martinez, E., Cobos-Carrascosa, E., Seijas-Vasquez, M. L., Fernandez-Sanchez, C., Strizzi, J. M., Torres-Alegrie, P., Santisteban-Martinez, J., & Gimenez-Sanchez, F. (2016b). Hand Hygiene Program Decreases School Absenteeism Due to Upper Respiratory Infections. *Journal of School Health*, 86(12), 873–881.
- Azor-Martinez, E., Yui-Hifume, R., Munoz-Vico, F. J., Jimenez-Noguera, E., Strizzi, J. M., Martinez-Martinez, I., Garcia-Fernandez, L., Seijas-Vazquez, M. L., Torres-Alegrie, P., Fernández-Campos, M. A., & Gimenez-Sanchez, F. (2018a). Effectiveness of a Hand Hygiene Program at Child Care Centers: A Cluster Randomized Trial. *Pediatrics*, 142(5), e20181245.
- Azor-Martinez, E., Yui-Hifume, R., Munoz-Vico, F. J., Jimenez-Noguera, E., Strizzi, J. M., Martinez-Martinez, I., Garcia-Fernandez, L., Seijas-Vazquez, M. L., Torres-Alegrie, P., Fernández-Campos, M. A., & Gimenez-Sanchez, F. (2018b). Effectiveness of a Hand Hygiene Program at Child Care Centers: A Cluster Randomized Trial. *Pediatrics*, 142(5), e20181245.
- Barasheed, O., Almasri, N., Badahdah, A.-M., Heron, L., Taylor, J., McPhee, K., Ridda, I., Haworth, E., Dwyer, D., Rashid, H., & Booy, R. (2014a). Pilot Randomised Controlled Trial to Test Effectiveness of Facemasks in Preventing Influenza-like Illness Transmission among Australian Hajj Pilgrims in 2011. *Infectious Disorders - Drug Targets*, 14, 110–116. <https://doi.org/10.2174/1871526514666141021112855>
- Barasheed, O., Almasri, N., Badahdah, A.-M., Heron, L., Taylor, J., McPhee, K., Ridda, I., Haworth, E., Dwyer, D., Rashid, H., & Booy, R. (2014b). Pilot Randomised Controlled Trial to Test Effectiveness of Facemasks in Preventing Influenza-like Illness Transmission among Australian Hajj Pilgrims in 2011. *Infectious Disorders - Drug Targets*, 14, 110–116. <https://doi.org/10.2174/1871526514666141021112855>
- Bish, A., & Michie, S. (2010). Demographic and attitudinal determinants of protective behaviours during a pandemic: A review. *British Journal of Health Psychology*, 15(4), 797–824. <https://doi.org/10.1348/135910710X485826>
- Blake, P., & Wadhwa, D. (2020). 2020 Year in Review: The impact of COVID-19 in 12 charts. World Bank Blog.
- Boone, S. A., & Gerba, C. P. (2007). Significance of Fomites in the Spread of Respiratory and Enteric Viral Diseases. *APPL. ENVIRON. MICROBIOL.*, 73, 10.
- Brooks, S. K., Webster, R. K., Smith, L. E., Woodland, L., Wessely, S., Greenberg, N., & Rubin, G. J. (2020). The psychological impact of quarantine and how to reduce it: Rapid review of the evidence. *The Lancet*, 395(10227), 912–920. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8)
- Bundgaard, H., Bundgaard, J. S., Raaschou-Pedersen, D. E. T., von Buchwald, C., Todsén, T., Norsk, J. B., Pries-Heje, M. M., Vissing, C. R., Nielsen, P. B., Winslow, U. C., Fogh, K., Hasselbalch, R., Kristensen, J. H., Ringgaard, A., Andersen, M. P., Bakkegaard Goecke, N., Trebbien, R., Skovgaard, K., Benfield, T., ... Iversen, K. (2020). Effectiveness of Adding a Mask Recommendation to Other Public Health Measures to Prevent SARS-CoV-2 Infection in Danish Mask Wearers. *Annals of Internal Medicine*.
- Cane, J., O'Connor, D., & Michie, S. (2012). Validation of the theoretical domains framework for use in behaviour change and implementation research. *Implementation Science*, 7(1), 37.
- Canini, L., Andréoletti, L., Ferrari, P., Angelo, D. R., Blanchon, T., Lemaitre, M., Filleul, L., Ferry, J. P., Desmaizieres, M., Smadja, S., Valleron, A. J., & Carrat, F. (2010). Surgical mask to prevent influenza transmission in households: A cluster randomized trial. *PLoS ONE*, 5(11), 1–6. <https://doi.org/10.1371/journal.pone.0013998>
- Carey, R. N., Connell, L. E., Johnston, M., Rothman, A. J., Bruin, M. De, Kelly, M. P., & Michie, S. (2019). Behavior Change Techniques and Their Mechanisms of Action: A Synthesis of Links Described in Published Intervention Literature. *Annals of Behavioral Medicine*, 53, 693–707. <https://doi.org/10.1093/abm/kay078>

- Chan, S. S. C., So, W. K. W., Wong, D. C. N., Lee, A. C. K., & Tiwari, A. (2007). Improving older adults' knowledge and practice of preventive measures through a telephone health education during the SARS epidemic in Hong Kong: A pilot study. *International Journal of Nursing Studies*, 44, 1120–1127. <https://doi.org/10.1016/j.ijnurstu.2006.04.019>
- Cheung, M. W. L. (2019). A Guide to Conducting a Meta-Analysis with Non-Independent Effect Sizes. *Neuropsychology Review*, 29, 387–396.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Hillsdale, NJ.
- Cowling, B. J., Chan, K.-H., Fang, V. J., Cheng, C. K. Y., Fung, R. O. P., Wai, W., Sin, J., Seto, W. H., Yung, R., Chu, D. W. S., Chiu, B. C. F., Lee, P. W. Y., Chiu, M. C., Lee, H. C., Uyeki, T. M., Houck, P. M., Peiris, J. S. M., & Leung, G. M. (2009). Facemasks and Hand Hygiene to Prevent Influenza Transmission in Households: A Cluster Randomized Trial. *Annals of Internal Medicine*, 151, 437–446.
- Cowling, B. J., Zhou, Y., Ip, D. K. M., Leung, G. M., & Aiello, A. E. (2010). Face masks to prevent transmission of influenza virus: A systematic review. *Epidemiology and Infection*, 138(4), 449–456. <https://doi.org/10.1017/S0950268809991658>
- Cowling, Benjamin J., Fung, R. O. P., Cheng, C. K. Y., Fang, V. J., Chan, K. H., Seto, W. H., Yung, R., Chiu, B., Lee, P., Uyeki, T. M., Houck, P. M., Peiris, J. S. M., & Leung, G. M. (2008). Preliminary findings of a randomized trial of non-pharmaceutical interventions to prevent influenza transmission in households. *PLoS ONE*, 3(5). <https://doi.org/10.1371/journal.pone.0002101>
- Cowling, B. J., Chan, K. H., Fang, V. J., Cheng, C. K. Y. (2009). Facemasks and Hand Hygiene to Prevent Influenza Transmission in Households. *Annals of Internal Medicine*, 151, 437–446.
- Edwards, R., Charani, E., Sevdalis, N., Alexandrou, B., Sibley, E., Mullett, D., Loveday, H. P., Drumright, L. N., & Holmes, A. (2012). Optimisation of infection prevention and control in acute health care by use of behaviour change: A systematic review. *The Lancet Infectious Diseases*, 12(4), 318–329. [https://doi.org/10.1016/S1473-3099\(11\)70283-3](https://doi.org/10.1016/S1473-3099(11)70283-3)
- Ferguson, N. M., Cummings, D. A. T., Fraser, C., Cajka, J. C., Cooley, P. C., & Burke, D. S. (2006). Strategies for mitigating an influenza pandemic. *Nature*, 442(7101), 448–452. <https://doi.org/10.1038/nature04795>
- Haby, M. M., Chapman, E., Clark, R., Barreto, J., Reveiz, L., & Lavis, J. N. (2016). What are the best methodologies for rapid reviews of the research evidence for evidence-informed decision making in health policy and practice: A rapid review. *Health Research Policy and Systems*, 14(1), 1–12. <https://doi.org/10.1186/s12961-016-0155-7>
- Hübner, N.-O., Hübner, C., Wodny, M., Kampf, G., & Kramer, A. (2010a). Effectiveness of alcohol-based hand disinfectants in a public administration: Impact on health and work performance related to acute respiratory symptoms and diarrhoea. *BMC Infectious Diseases*, 10(250).
- Hübner, N.-O., Hübner, C., Wodny, M., Kampf, G., & Kramer, A. (2010b). Effectiveness of alcohol-based hand disinfectants in a public administration: Impact on health and work performance related to acute respiratory symptoms and diarrhoea. *BMC Infectious Diseases*, 10(250).
- Huis, A., van Achterberg, T., de Bruin, M., Grol, R., Schoonhoven, L., & Hulscher, M. (2012). A systematic review of hand hygiene improvement strategies: A behavioural approach. *Implementation Science*, 7(1), 1–14. <https://doi.org/10.1186/1748-5908-7-92>
- International Monetary Fund. (2020). *World Economic Outlook Update*.
- Kaewchana, S., Simmerman, M., Somrongsong, R., Suntarattiwong, P., Lertmaharit, S., & Chotipitayusunondh, T. (2012). Effect of Intensive Hand Washing Education on Hand Washing Behaviors in Thai Households with an Influenza-Positive Child in Urban Thailand. *Asia-Pacific Journal of Public Health*, 24(4), 577–585. <https://doi.org/10.1177/1010539510393728>
- Killingley, B., & Nguyen-Van-Tam, J. (2013a). Routes of influenza transmission. *Influenza and Other Respiratory Viruses*, Suppl. 2, 42–51. <https://doi.org/10.1111/irv.12080>
- Koep, T. H., Jenkins, S., Hammerlund, M. E. M., Clemens, C., Fracica, E., Ekker, S. C., Enders, F. T., Huskins, W. C., & Pierret, C. (2016). Promotion of Influenza Prevention Beliefs and Behaviors through Primary School Science Education. *Journal of Community Medicine and Health Education*, 6(3). <https://doi.org/10.4172/2161-0711.1000444>
- Kwok, Y. L. A., Gralton, J., & McLaws, M. L. (2015). Face touching: A frequent habit that has implications for hand hygiene. *American Journal of Infection Control*, 43(2), 112–114. <https://doi.org/10.1016/j.ajic.2014.10.015>
- Larson, E. L., Ferng, Y.-H., McLoughlin, J. W., Wang, S., & Morse, S. S. (2009). Effect of Intensive Education on Knowledge, Attitudes, and Practices Regarding Upper Respiratory Infections Among Urban Latinos. *Nursing Research*, 58(3), 150–157.
- Larson, E. L., Ferng, Y.-H., Wong-McLoughlin, J., Wang, S., Haber, M., & Morse, S. S. (2010). Impact of Non-Pharmaceutical Interventions on URIs and Influenza in Crowded, Urban Households. *Public Health Reports*, 125, 178–191.

- Little, P., Stuart, B., Hobbs, F. D. R., Moore, M., Barnett, J., Popoola, D., Middleton, K., Kelly, J., Mullee, M., Raftery, J., Yao, G., Carman, W., Fleming, D., Stokes-lampard, H., Williamson, I., Joseph, J., Miller, S., & Yardley, L. (2014). An internet-delivered handwashing intervention to modify influenza-like illness and respiratory infection transmission (PRIMIT): A primary care randomised trial. *The Lancet*, 386(10004), 1631–1639. [https://doi.org/10.1016/S0140-6736\(15\)60127-1](https://doi.org/10.1016/S0140-6736(15)60127-1)
- Liu, X., Zhao, Z., Hou, W., Polinder, S., van Beeck, E. F., Zhang, Z., Zhou, Y., Liu, G., Xie, X., Cheng, J., Richardus, J. H., & Erasmus, V. (2019). A multimodal intervention to improve hand hygiene compliance via social cognitive influences among kindergarten teachers in China. *PLoS ONE*, 14(5), e0215824.
- Luangasanatip, N., Hongsuwan, M., Limmathurotsakul, D., Lubell, Y., Lee, A. S., Harbarth, S., Day, N. P. J., Graves, N., & Cooper, B. S. (2015). Comparative efficacy of interventions to promote hand hygiene in hospital: Systematic review and network meta-analysis. *BMJ (Online)*, 351. <https://doi.org/10.1136/bmj.h3728>
- Lunn, P., Belton, C., Lavin, C., McGowan, F., Timmons, S., & Robertson, D. (2020). Using Behavioural Science to Help Fight the Coronavirus. *Economic and Social Research Institute*, 1–24.
- MacIntyre, C. R., Cauchemez, S., Dwyer, D. E., Seale, H., Cheung, P., Browne, G., Fasher, M., Wood, J., Gao, Z., Booy, R., & Ferguson, N. (2009). Face mask use and control of respiratory virus transmission in households. *Emerging Infectious Diseases*, 15(2), 233–241. <https://doi.org/10.3201/eid1502.081167>
- Mbakaya, B. C., Lee, P. H., & Lee, R. L. T. (2017). Hand hygiene intervention strategies to reduce diarrhoea and respiratory infections among schoolchildren in developing countries: A systematic review. *International Journal of Environmental Research and Public Health*, 14(4), 1–14. <https://doi.org/10.3390/ijerph14040371>
- Michie, S., Atkins, L., & West, R. (2014). *The Behaviour Change Wheel: A Guide to Designing Interventions*. Silverback Publishing.
- Michie, S., Richardson, M., Johnston, M., Abraham, C., Francis, J., Hardeman, W., Eccles, M. P., Cane, J., & Wood, C. E. (2013). The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: Building an international consensus for the reporting of behavior change interventions. *Annals of Behavioral Medicine*, 46(1), 81–95. <https://doi.org/10.1007/s12160-013-9486-6>
- Michie, S., Rubin, J., & Amlôt, R. (2020, February 28). Behavioural science must be at the heart of the public health response to covid-19. *BMJ Opinion*.
- Michie, S., West, R., & Amlôt, R. (2020, March 3). Behavioural strategies for reducing covid-19 transmission in the general population. *BMJ Opinion*.
- Michie, S., West, R., Amlôt, R., & Rubin, J. (2020, March 11). Slowing down the covid-19 outbreak: Changing behaviour by understanding it. *BMJ Opinion*.
- Moore, G. F., & Evans, R. E. (2017). What theory, for whom and in which context? Reflections on the application of theory in the development and evaluation of complex population health interventions. *SSM - Population Health*, 3, 132–135. <https://doi.org/10.1016/j.ssmph.2016.12.005>
- Morawska, L., & Milton, D. K. (2020). It Is Time to Address Airborne Transmission of Coronavirus Disease 2019 (COVID-19). *Clinical Infectious Diseases*, 71, 2311–2313. <https://doi.org/10.1093/cid/ciaa939>
- Mott, P. J., Sisk, B. W., Arbogast, J. W., Ferrazzano-Yaussy, C., Bondi, C. A. M., & Sheehan, J. J. (2007). Alcohol-Based Instant Hand Sanitizer Use in Military Settings: A Prospective Cohort Study of Army Basic Trainees. *Military Medicine*, 172, 1170–1176.
- Nandrup-Bus, I. (2009a). Mandatory handwashing in elementary schools reduces absenteeism due to infectious illness among pupils: A pilot intervention study. *American Journal of Infection Control*, 37(10), 820–826. <https://doi.org/10.1016/j.ajic.2009.06.012>
- Nandrup-Bus, I. (2009b). Mandatory handwashing in elementary schools reduces absenteeism due to infectious illness among pupils: A pilot intervention study. *American Journal of Infection Control*, 37(10), 820–826. <https://doi.org/10.1016/j.ajic.2009.06.012>
- Norris, E., Finnerty, A. N., Hastings, J., Stokes, G., & Michie, S. (2019). A scoping review of ontologies related to human behaviour change. *Nature Human Behaviour*, 3(2), 164–172. <https://doi.org/10.1038/s41562-018-0511-4>
- Olena Doronina, R. N., Jones, D., Martello, M., Biron, A., & Lavoie-Tremblay, M. (2017). A Systematic Review on the Effectiveness of Interventions to Improve Hand Hygiene Compliance of Nurses in the Hospital Setting. *Journal of Nursing Scholarship*, 49(2), 143–152. <https://doi.org/10.1111/jnu.12274>
- Öncü, E., Köksöy Vayisoğlu, S., Lafci, D., Yurtsever, D., Ravlı Bulut, E., & Peker, E. (2019). Comparison of Interactive Education Versus Fluorescent Concretization on Hand Hygiene Compliance Among Primary School Students: A Randomized Controlled Trial. *The Journal of School Nursing*, 35(5), 337–347. <https://doi.org/10.1177/1059840518785447>
- Or, P. P.-L., Ching, P. T.-Y., & Chung, J. W.-Y. (2020). Can Flu-Like Absenteeism in Kindergartens Be Reduced Through Hand Hygiene Training for Both Parents and Their Kindergarteners? *Journal of Primary Care & Community Health*, 11, 1–6. <https://doi.org/10.1177/2150132719901209>

- Ram, P. K., Divita, M. A., Khatun-e-Jannat, K., Islam, M., Krytus, K., Cercone, E., Munir Sohel, B., Ahmed, M., Rahman, Q., Rahman, M., Yu, J., Brooks, W. A., Azziz-Baumgartner, E., Fry, A. M., & Luby, S. P. (2015). Impact of Intensive Handwashing Promotion on Secondary Household Influenza-Like Illness in Rural Bangladesh: Findings from a Randomized Controlled Trial. *PLoS ONE*, 10(6), e0125200. <https://doi.org/10.1371/journal.pone.0125200>
- Reyes Fernández, B., Lippke, S., Knoll, N., Moya, E. B., & Schwarzer, R. (2015a). Promoting action control and coping planning to improve hand hygiene. *BMC Public Health*, 15(964). <https://doi.org/10.1186/s12889-015-2295-z>
- Reyes Fernández, B., Lippke, S., Knoll, N., Moya, E. B., & Schwarzer, R. (2015b). Promoting action control and coping planning to improve hand hygiene. *BMC Public Health*, 15(964). <https://doi.org/10.1186/s12889-015-2295-z>
- Roberts, L., Smith, W., Jorm, L., Patel, M., Douglas, R. M., & McGilchrist, C. (2000a). Effect of Infection Control Measures on the Frequency of Upper Respiratory Infection in Child Care: A Randomized, Controlled Trial. *Pediatrics*, 105(4), 738–742.
- Roberts, L., Smith, W., Jorm, L., Patel, M., Douglas, R. M., & McGilchrist, C. (2000b). Effect of Infection Control Measures on the Frequency of Upper Respiratory Infection in Child Care: A Randomized, Controlled Trial. *Pediatrics*, 105(4), 738–742.
- Rosenstock, I. M. (1974). Historical origins of the health belief model. *Health Education Monographs*, 2(4), 328–335.
- SafeLives. (2020). *Domestic abuse and COVID-19*.
- Sandora, T. J., Taveras, E. M., Shih, M.-C., Resnick, E. A., Lee, G. M., Ross-Degnan, D., & Goldmann, D. A. (2005a). A Randomized, Controlled Trial of a Multifaceted Intervention Including Alcohol-Based Hand Sanitizer and Hand-Hygiene Education to Reduce Illness Transmission in the Home. *Pediatrics*, 116(3), 587–594. <https://doi.org/10.1542/peds.2005-0199>
- Sandora, T. J., Taveras, E. M., Shih, M.-C., Resnick, E. A., Lee, G. M., Ross-Degnan, D., & Goldmann, D. A. (2005b). A Randomized, Controlled Trial of a Multifaceted Intervention Including Alcohol-Based Hand Sanitizer and Hand-Hygiene Education to Reduce Illness Transmission in the Home. *Pediatrics*, 116(3), 587–594. <https://doi.org/10.1542/peds.2005-0199>
- Savolainen-Kopra, C., Haapakoski, J., Peltola, P. A., Ziegler, T., Korpela, T., Anttila, P., Amirousefi, A., Huovinen, P., Huvinen, M., Noronen, H., Riikkala, P., Roivainen, M., Ruutu, P., Teirila, J., Vartiainen, E., & Hovi, T. (2012a). Hand washing with soap and water together with behavioural recommendations prevents infections in common work environment: An open cluster-randomized trial. *Trials*, 13(10).
- Savolainen-Kopra, C., Haapakoski, J., Peltola, P. A., Ziegler, T., Korpela, T., Anttila, P., Amirousefi, A., Huovinen, P., Huvinen, M., Noronen, H., Riikkala, P., Roivainen, M., Ruutu, P., Teirila, J., Vartiainen, E., & Hovi, T. (2012b). Hand washing with soap and water together with behavioural recommendations prevents infections in common work environment: An open cluster-randomized trial. *Trials*, 13(10).
- Simmerman, J. M., Suntarattiwong, P., Levy, J., Jarman, R. G., Kaewchana, S., Gibbons, R. V., Cowling, B. J., Sanasuttipun, W., Maloney, S. A., Uyeki, T. M., Kamimoto, L., & Chotipitayasanondh, T. (2011a). Findings from a household randomized controlled trial of hand washing and face masks to reduce influenza transmission in Bangkok, Thailand. *Influenza and Other Respiratory Viruses*, 5, 256–267. <https://doi.org/10.1111/j.1750-2659.2011.00205.x>
- Simmerman, J. M., Suntarattiwong, P., Levy, J., Jarman, R. G., Kaewchana, S., Gibbons, R. V., Cowling, B. J., Sanasuttipun, W., Maloney, S. A., Uyeki, T. M., Kamimoto, L., & Chotipitayasanondh, T. (2011b). Findings from a household randomized controlled trial of hand washing and face masks to reduce influenza transmission in Bangkok, Thailand. *Influenza and Other Respiratory Viruses*, 5, 256–267. <https://doi.org/10.1111/j.1750-2659.2011.00205.x>
- Stebbins, S., Stark, J. H., & Vukotich Jr, C. J. (2010). Compliance With a Multilayered Nonpharmaceutical Intervention in an Urban Elementary School Setting. *Journal of Public Health Management and Practice*, 16(4), 316–324.
- Stedman-Smith, M., DuBois, C. L. Z., Grey, S. F., Kingsbury, D. M., Shakya, S., Scofield, J., & Slenkovich, K. (2015a). Outcomes of a Pilot Hand Hygiene Randomized Cluster Trial to Reduce Communicable Infections Among US Office-Based Employees. *Journal of Occupational and Environmental Medicine*, 57(4), 374–380. <https://doi.org/10.1097/JOM.0000000000000421>
- Stedman-Smith, M., DuBois, C. L. Z., Grey, S. F., Kingsbury, D. M., Shakya, S., Scofield, J., & Slenkovich, K. (2015b). Outcomes of a Pilot Hand Hygiene Randomized Cluster Trial to Reduce Communicable Infections Among US Office-Based Employees. *Journal of Occupational and Environmental Medicine*, 57(4), 374–380. <https://doi.org/10.1097/JOM.0000000000000421>
- Suess, T., Remschmidt, C., Schink, S., Luchtenberg, M., Haas, W., Krause, G., & Buchholz, U. (2011). Facemasks and intensified hand hygiene in a German household trial during the 2009/2010 influenza A (H1N1) pandemic: Adherence and tolerability in children and adults. *Epidemiology & Infection*, 139, 1895–1901. <https://doi.org/10.1017/S0950268810003006>
- Suess, Thorsten, Remschmidt, C., Schink, S. B., Schweiger, B., Nitsche, A., Schroeder, K., Doellinger, J., Milde, J., Haas, W., Koehler, I., Krause, G., & Buchholz, U. (2012). The role of facemasks and hand hygiene in the prevention of influenza transmission in households: Results from a cluster randomised



- trial; Berlin, Germany, 2009–2011. *BMC Infectious Diseases*, 12(26). <https://doi.org/10.1186/1471-2334-12-26>
- The Cochrane Collaboration. (2011). *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0. [Updated March 2011]*. (J. Higgins & S. Green, Eds.).
- Tricco, A. C., Antony, J., Zarin, W., Striffler, L., Ghassemi, M., Ivory, J., Perrier, L., Hutton, B., Moher, D., & Straus, S. E. (2015). A scoping review of rapid review methods. *BMC Medicine*, 13, 224. <https://doi.org/10.1186/s12916-015-0465-6>
- Updegraff, J. A., Emanuel, A. S., Gallagher, K. M., & Steinman, C. T. (2011). Framing Flu Prevention—An Experimental Field Test of Signs Promoting Hand Hygiene During the 2009–2010 H1N1 Pandemic. *Health Psychology*, 30(3), 295–299. <https://doi.org/10.1037/a0023125>
- Viechtbauer, W. (2010). Conducting Meta-Analyses in R with the metafor Package. *Journal of Statistical Software*, 36(3), 1–48.
- Warren-Gash, C., Fragaszy, E., & Hayward, A. C. (2012). Hand hygiene to reduce community transmission of influenza and acute respiratory tract infection: A systematic review. *Influenza and Other Respiratory Viruses*, 7(5), 738–749. <https://doi.org/10.1111/irv.12015>
- West, R., & Michie, S. (2020). Routes of transmission of SARS-CoV-2 and behaviours to block it: A summary. *Qeios*, 1–4.
- West, R., Michie, S., Rubin, J., & Amlôt, R. (n.d.). How understanding principles of behaviour change can help limit the spread of COVID-19. *Submitted*.
- West, R., Michie, S., Rubin, J., & Amlôt, R. (2020, April 3). Don't touch the T-Zone—How to block a key pathway to infection with SARS-CoV-2. *BMJ Opinion*.
- White, C., Kolbe, R., Carlson, R., Lipson, N., & Dolan, M. (2003). The effect of hand hygiene on illness rate among students in university residence halls. *American Journal of Infection Control*, 31(6), 364–370. <https://doi.org/10.1067/mic.2003.76>
- Wood, C. E., Richardson, M., Johnston, M., Abraham, C., Francis, J., Hardeman, W., & Michie, S. (2015). Applying the behaviour change technique (BCT) taxonomy v1: A study of coder training. *Translational Behavioral Medicine*, 5, 134–148. <https://doi.org/10.1007/s13142-014-0290-z>
- World Health Organisation. (2017). *Rapid reviews to strengthen health policy and systems: A practical guide* (A. C. Tricco, E. V Langlois, & S. E. Straus, Eds.). <https://doi.org/10.1111/j.1365-2966.2008.12859.x>
- World Health Organisation. (2019). *Non-pharmaceutical public health measures for mitigating the risk and impact of epidemic and pandemic influenza*.
- Yardley, L., Miller, S., Schlotz, W., & Little, P. (2011). Evaluation of a Web-Based Intervention to Promote Hand Hygiene: Exploratory Randomized Controlled Trial. *Journal of Medical Internet Research*, 13(4), e107. <https://doi.org/10.2196/jmir.1963>
- Zomer, T. P., Erasmus, V., Looman, C. W., van Beeck, E. F., Tjon-a-Tsien, A., Richardus, J. H., & Voeten, H. A. C. M. (2016). Improving hand hygiene compliance in child daycare centres: A randomized controlled trial. *Epidemiology & Infection*, 144, 2552–2560. <https://doi.org/10.1017/S0950268816000911>

**Supplementary data:** available at <https://doi.org/10.32388/OJXD82.2>

## Declarations

**Funding:** The author(s) received no specific funding for this work.

**Potential competing interests:** The author(s) declared that no potential competing interests exist.