

Mastoparans

Carlos Henrique Marchiori¹

¹ Instituto Federal Goiano

Potential competing interests: No potential competing interests to declare.

Co-authors: Marco Vinícios de Oliveira Santana² and Klebert de Paula Malheiros³.

^{2,3}Instituto Marco Santana, Goiânia, Goiás, Brazil.

The mastoparans, the most widely described class of peptides isolated from the venom of different wasp species, were named after their first recognized target, mast cells. They are lysine-rich peptides and perform several biological functions, including the release of mediators through activation of G-protein receptors, stimulation of phospholipase A2 and C, mobilization of Ca²⁺ to the sarcoplasmic reticulum, involvement in cell death by necrosis or apoptosis, and antimicrobial activity. Studies carried out with different types of mastoparans showed that some peptides have low hemolytic activity, but high antibiotic activity (Figure 1) [1-4].




Figure 1. Biochemistry of mastoparan. a α -Helical secondary structure of mastoparan-X isolated from *Vespa xanthoptera* (NCBI-Protein). Histamine released by b G protein binding and c direct endosome binding.

Source: https://www.researchgate.net/figure/Biochemistry-of-mastoparan-a-a-Helical-secondary-structure-of-mastoparan-X-isolated-from_fig2_343165039.

Several studies have shown that mastoparans interact with natural and artificial biological membranes, assuming an amphipathic α -helix conformation, which may cause an increase in permeability, which is the primary cause of the effect of releasing mediators of cell activation. The need to understand the mechanism of action of peptides, such as mastoparanes, has opened up a vast field for research on the structures

[[= and functions of these compounds. Many authors claim that the structure strongly interferes with the intensity and/or type of biological activity of biologically active peptides [4-7].

Therefore, their structures must be determined, not only to explain their biological activities but also to favor the

Mastoparan $\text{H}_2\text{N-INLKALAALAKKIL-CONH}_2$

FUB86 $\text{H}_3\text{C}-(\text{CH}_2)_{11}-\text{N}(\text{H})-\text{CH}(\text{NH}_2)-(\text{CH}_2)_4-\text{NH}_2$

FUB132 $\text{H}_3\text{C}-(\text{CH}_2)_{11}-\text{N}(\text{H})-\text{CH}(\text{NH}_2)-(\text{CH}_2)_4-\text{N}(\text{H})-\text{CH}(\text{NH}_2)-(\text{CH}_2)_4-\text{N}(\text{H})-\text{CH}(\text{NH}_2)-(\text{CH}_2)_4-\text{NH}_2$

Figure 2. Main biological activities exhibited by mastoparans. Sources: DOI: 10.3389/fmolb.2022.824989 PMID: 35813822; PMCID: PMC9263278.

Structurally, wasp peptides are presented in an α -helix conformation, having 10 to 14 amino acid residues in their primary sequence. They are rich in hydrophobic amino acids such as leucine, isoleucine, and alanine, as well as some lysine in their primary sequences. Although the amino acid sequences of these peptides are relatively conserved, substitutions of some residues seem to alter the biological activities of these peptides, proving to be an interesting basis for future studies using pharmacological applications (Figure 3) [10-12].




Figure 3. Mrgprb2-mediated mast cell activation by mastoparans leading to neutrophil recruitment and *Staphylococcus aureus* (Bacillales: Staphylococcaceae) clearance in a mouse dermonecrotic-infected lesion. ROS—reactive oxygen species; NETs—neutrophil extracellular traps. Sources: DOI: 10.3389/fmolb.2022.824989 PMID: 35813822; PMCID: PMC9263278.

Different amino acid sequences of the mastoparan peptide isolated from the social wasp species *Polybia paulista* Ihering, 1896 were identified. The polybia-MPI and polybia-CP peptides, synthesized from the venom of the wasp *P. paulista*, show potent antibacterial activity against strains of *Escherichia coli*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, and *Bacillus subtilis*. Two other peptides with antibacterial activity, called Polybina I and II, were also isolated from the same species of wasp. The venom of *Polybia occidentalis* (Olivier, 1791), a social wasp, has anticoagulant and fibrinogen-degrading pharmacological properties. Anticoagulation occurs at different stages of the coagulation process in the intrinsic, extrinsic, and specific pathways. The venom can inhibit platelet aggregation and destroy plasma fibrinogens [12-14].

References

- [1] Ángel M. The venom of a Brazilian wasp (*Polybia paulista*) can kill cancer cells [Internet]. Madrid: EL PAÍS; @2015 [cited 2024 Mar 22]. Available from https://brasil.elpais.com/brasil/2015/09/01/ciencia/1441100654_055665.html.
- [2] Souza CL, et al. Revisiting *Polybia paulista* wasp venom using shotgun proteomics – Insights into the N-linked glycosylated venom proteins. 2019; 200(30): 60-73.
- [2] Vila-Farrés X, et al. Sequence-activity relationship, and mechanism of action of mastoparan analogues against extended-drug resistant *Acinetobacter baumannii*. European Journal of Medicinal Chemistry. 2015; 101: 34-40.

- [3] Santana CJC, Pires Júnior OR, Fontes W, Palma MS, Castro MS. Mastoparans: A group of multifunctional α -helical peptides with promising therapeutic properties. *Frontiers in Molecular Biosciences*. 2022; 24(9): 824989. Source: doi: 10.3389/fmolb.2022.824989. PMID: 35813822; PMCID: PMC9263278.
- [4] Wine JFC. Trypanocidal action of a mastoparan from *Polybia paulista* and its possible mechanism of action [Ph.D. dissertation]. Fortaleza: Federal University of Ceará; 2015.
- [5] Abd El-Wahed A., et al.. Wasp venom biochemical components and their potential in biological applications and nanotechnological interventions. *Toxins (Basel)*. 2021;13(3): 206.
- [6] Arifuzzaman M, et al. MRGPR-mediated activation of local mast cells clears cutaneous bacterial infection and protects against reinfection. *Science Advances*. 2019; 5: eaav0216.
- [7] Chahdi A, Choi WS, Kim YM, Beaven MA. Mastoparan selectively activates phospholipase d2 in cell membranes. *Journal of Biological Chemistry*. 2003; 278: 12039–12045.
- [8] Chen X, et al. Evaluation of the bioactivity of a mastoparan peptide from wasp venom and of its analogues designed through targeted engineering. *International Journal of Biological Sciences*. 2018; 14: 599–607.
- [9] Colella F, Scillitani G, Pierri CL. Sweet as honey, bitter as bile: Mitochondriotoxic peptides and other therapeutic proteins isolated from animal tissues, for dealing with mitochondrial apoptosis. *Toxicology*. 2021; 447: 152612.
- [10] Silva AMB. Pro-necrotic activity of cationic mastoparan peptides in human glioblastoma multiforme cells via membranolytic action. *Molecular Neurobiology*. 2018; 55: 5490–5504.
- [11] Silva AMB. Pro-necrotic activity of cationic mastoparan peptides in human glioblastoma multiforme cells via membranolytic action. *Molecular Neurobiology*. 2018; 55: 5490–5504.
- [12] da Silva AVR, et al. The effects of the c-terminal amidation of mastoparans on their biological actions and interactions with membrane-mimetic systems. *Biochimica et Biophysica Acta (BBA) - Biomembranes*. 2014; 1838, 2357–2368.
- [13] das Neves RC, et al. Antimicrobial and antibiofilm effects of peptides from venom of social wasp and scorpion on multidrug-resistant *Acinetobacter baumannii*. *Toxins*. 2019; 11: 216.
- [14] de Azevedo RA, et al. Mastoparan induces apoptosis in b16f10-nex2 melanoma cells via the intrinsic mitochondrial pathway and displays antitumor activity in vivo. *Peptides*. 2015; 68: 113–119.