

Research Article

Conditions for Emergent Gravitational Light Bending from a Logarithmic Superfluid Vacuum

Boris Kulangiev¹

1. Independent researcher

Purely scalar theories of gravitation predict a Parameterized Post-Newtonian (PPN) parameter $\gamma = 0$, yielding only half the observed light deflection. We trace this failure to a structural feature of real scalar wave equations: the characteristic propagation speed is fixed by the D'Alembertian operator and cannot depend on the background field, regardless of the self-interaction potential. We show that modeling the vacuum as a complex superfluid with a relativistic logarithmic nonlinearity, analyzed via the Madelung transformation, introduces two new degrees of freedom absent in the wave equation form: a density-dependent sound speed and a macroscopic flow velocity. Using the standard relativistic acoustic-metric formalism of Barceló-Liberati-Visser, we derive the sound speed $c_s^2/c^2 = [2\ln(\bar{\rho}/\rho_c) + 3]^{-1}$ from the logarithmic equation of state, demonstrating its dependence on the local vacuum density—an effect absent in the non-relativistic limit. We compute the PPN parameter for the static acoustic metric, finding $\gamma = (\alpha - 1)/(\alpha + 1)$ where $\alpha = d\ln c_s/d\ln \rho$. This expression equals unity for no value of α : the static density channel cannot reproduce general-relativistic light bending under any barotropic equation of state. The logarithmic equation of state at the background density gives $\alpha = -1$, which is in addition the exact pole of this expression, so the static density channel is not merely numerically wrong but mathematically ill-defined, which forces the deflection to be carried by a different channel. We then demonstrate that a non-static acoustic metric with macroscopic vacuum flow does exactly this—the deflection is produced by advection of the signal by the flow, not by a refractive sound-speed gradient—and yields $\gamma = 1$ exactly, provided the flow velocity satisfies $v(r) = \sqrt{2G_{eff}M/r}$ —the Painlevé-Gullstrand profile. Because the bending is advective, it acts identically on excitations of any propagation speed and is therefore achromatic. Using a self-consistent Bondi accretion calculation, we show that the logarithmic equation of state does not naturally produce this profile in the far field, where the flow decays as r^{-2} rather than the

required $r^{-1/2}$. We further show that the Painlevé–Gullstrand flow is the *only* self-consistent weak-field solution of the acoustic metric equations: the static metric is divergent, whereas the flowing metric closes the self-consistency loop for any barotropic fluid. The remaining open problem is to derive this macroscopic flow as a collective effect of the microscopic soliton dynamics.

Correspondence: papers@team.qeios.com — Qeios will forward to the authors

1. Introduction

The bending of light by gravity is one of the most precise tests of gravitational theory. In the PPN formalism ^[1], the deflection of a light ray with impact parameter b by a mass M in the grazing-ray limit is

$$\delta\theta = \frac{(1 + \gamma) 2GM}{bc^2}, \quad (1)$$

where γ encodes the ratio of spatial to temporal curvature in the weak-field metric. General relativity predicts $\gamma = 1$; the Cassini spacecraft confirmed $\gamma = 1.000021 \pm 0.000023$ ^[2].

Purely scalar theories generically fail this test. Nordström’s conformally flat theory gives $\gamma = -1$ ^[3]. A real scalar field with standard kinetic term gives $\gamma = 0$ ^[1]—Einstein’s 1911 result ^[4], using only time dilation without spatial curvature. This half-deflection led to the conclusion that gravity requires a rank-2 tensor field.

In this paper, we examine whether a superfluid vacuum model—specifically, a complex scalar field with logarithmic self-interaction ^{[5][6][7]} analyzed in the Madelung hydrodynamic representation ^{[8][9][10]}—can overcome this constraint. We find that the Madelung formulation introduces two structural ingredients absent in the wave equation form: (i) a density-dependent sound speed, and (ii) a macroscopic flow velocity. Both are needed for $\gamma = 1$. We derive the precise conditions required, and show via self-consistent computation that these conditions are not automatically satisfied, identifying the central open problem for superfluid vacuum gravity.

2. The Kinematic Trap: Why $\gamma = 0$ for Real Scalar Fields

Consider a real scalar field $D(x, t)$ satisfying

$$\square D + V'(D) = J(x, t), \quad (2)$$

where $\square = \nabla^2 - c^{-2} \partial_t^2$. We show that the geometric-optics propagation speed of perturbations on any background $\bar{D}(x)$ is identically c .

Linearizing $D = \bar{D} + \delta D$:

$$\square(\delta D) + V''(\bar{D})\delta D = 0. \quad (3)$$

In the WKB limit $\delta D \sim Ae^{i\phi}$ with $|\nabla\phi| \gg |\nabla A/A|$, the eikonal equation is

$$|\nabla\phi|^2 - \frac{1}{c^2} \left(\frac{\partial\phi}{\partial t} \right)^2 = 0, \quad (4)$$

giving $v_{ph} = c$ everywhere. The term $V''(\bar{D})$ contributes only a subleading effective mass. Figure 1 confirms this numerically: a wave packet in a Klein-Gordon field propagates at identical velocity for background densities spanning an order of magnitude.




Figure 1. The kinematic trap. Group velocity of a wave packet in a Klein-Gordon field vs. background density D_{bg} . The velocity is constant (red squares), confirming Eq. (4). The green curve shows the density dependence needed for $\gamma = 1$.

Since the coordinate speed of light cannot vary with the background, the only deflection contribution comes from time dilation:

$$\delta\theta_{\text{scalar}} = \frac{2GM}{bc^2}, \quad (5)$$

giving $\gamma = 0$. This holds for *any* potential $V(D)$: the characteristic surfaces of (2) are determined by the principal symbol (the D'Alembertian), not by V . Escaping this constraint requires either modifying the kinetic term, as in scalar-tensor theories ^[11], or reinterpreting the field in a formulation where the propagation speed emerges dynamically.

3. The Madelung Superfluid Formulation

Interpretation of the order parameter

We take a deliberately classical-field stance toward Eq. (6) below: $\psi(x, t)$ is a complex *order parameter* for a deterministic vacuum substrate, not a quantum-mechanical wavefunction in the standard sense. Its squared magnitude $|\psi|^2 = \rho$ is a number density of the substrate's microscopic constituents, not a probability density, and is not subject to the unitary normalization $\int |\psi|^2 d^3x = 1$. The Madelung decomposition $\psi = \sqrt{\rho} e^{iS/\hbar_{\text{eff}}}$ is therefore a hydrodynamic ansatz for a real classical fluid, in the spirit of de Broglie's pilot-wave program ^[12] and the Madelung-Bohm hydrodynamic interpretation of quantum mechanics ^{[8][9][10]}. The relativistic Lorentz-covariant form of Eq. (6) reflects an emergent symmetry of small fluctuations around the condensate's rest frame—identified physically with the cosmic vacuum frame in which the cosmic microwave background dipole vanishes—rather than an exact symmetry of the underlying theory. We do not specify a UV completion or a quantization procedure for the underlying field; the framework is studied here as a classical effective theory whose hydrodynamic predictions are compared with weak-field gravitational observations.

3.1. The logarithmic superfluid

We model the vacuum as a complex order parameter $\psi(x, t)$ evolving under the relativistic logarithmic Klein-Gordon equation ^{[5][6]}, the relativistic counterpart of the logarithmic Schrödinger equation ^[7]:

$$\square\psi + \mu^2\psi - b\psi\ln\left(\frac{|\psi|^2}{\rho_0}\right) = 0. \quad (6)$$

The logarithmic nonlinearity produces Gaussian-localized solitons (“Gaussons”) ^{[7][13]}.

Units of the order parameter

A clarification on dimensions. In the relativistic k-essence Lagrangian $\mathcal{L} = \rho X - V(\rho)$, with $X = \partial_\mu S \partial^\mu S$ and S having dimensions of action, the variable $\rho = |\psi|^2$ does not have units of mass density $[\text{kg}/\text{m}^3]$. Demanding $[\mathcal{L}] = [\text{energy}/\text{volume}]$ together with $[X] = [\text{momentum}^2]$ forces $[\rho] = [(\text{energy} \cdot \text{volume})^{-1}]$. The Lagrangian variable ρ is therefore an inverse-energy-volume quantity; the physical mass density of the condensate, which is what enters the gravitational source term and what one would measure as kg/m^3 , is a *distinct* quantity computed from the time-time component of the stress-energy tensor:

$$\rho_{mass} = \frac{T_{00}}{c^2} = \frac{\rho V'(\rho) + V(\rho)}{c^2}, \quad (7)$$

where the right-hand side is evaluated using the Lagrangian variable ρ and reduces, at the background, to a constant ρ_0^{phys} . We retain the conventional notation ρ for the order parameter throughout, but the reader should keep in mind that physical observables involving mass density use $\rho_{mass} = T_{00}/c^2$, not ρ itself. The Madelung decomposition below treats ρ as the field variable, with all subsequent equations dimensionally consistent under this identification.

3.2. The Madelung decomposition

Applying $\psi = \sqrt{\rho} e^{iS/\hbar_{eff}}$ and separating real and imaginary parts yields two coupled equations:

Continuity (conservation of vacuum flux):

$$\partial_\mu (\rho \partial^\mu S) = 0. \quad (8)$$

Hamilton-Jacobi (relativistic phase dynamics):

$$\frac{1}{\hbar_{eff}^2} \partial_\mu S \partial^\mu S = \mu^2 - b \ln\left(\frac{\rho}{\rho_0}\right) + \frac{\square \sqrt{\rho}}{\sqrt{\rho}}. \quad (9)$$

The last term is the quantum potential (Bohm potential), negligible in the macroscopic limit. The fluid four-velocity is $u_\mu = \partial_\mu S / m_{eff}$, irrotational by construction except at topological defects (vortices).

The crucial point is that Eq. (9), unlike a wave equation, couples the phase gradient $\partial_\mu S$ to the density ρ through a constraint that depends on the local thermodynamic state. This coupling produces a density-dependent sound speed, as we now show.

Origin of the continuity equation

The d'Alembertian in Eq. (6) contains second-order time derivatives, so Eq. (6) is a wave equation, not directly a continuity equation. However, the Madelung decomposition $\psi = \sqrt{\rho} e^{iS/\hbar_{eff}}$ separates Eq. (6) into real and imaginary parts that have distinct physical content. The real part yields the relativistic Hamilton-Jacobi equation (9), which is second-order in time and governs phase dynamics. The imaginary part yields the first-order conservation law in Eq. (8) which is the relativistic continuity equation for the four-current $J^\mu = \rho \partial^\mu S / \hbar_{eff}$ associated with the global U(1) symmetry $\psi \rightarrow e^{i\theta} \psi$ of Eq. (6). In the rest frame of the condensate, where $\partial_t S \approx -\hbar_{eff} \omega_0$ is approximately constant and spatial gradients are small, Eq. (8) reduces to the familiar non-relativistic continuity form $\partial_t \rho + \nabla \cdot (\rho \mathbf{v}) = 0$ with $\mathbf{v} = \nabla S / m_{eff}$. The wave-equation character of Eq. (6) is therefore consistent with the existence of a first-order conservation law for the U(1) charge density.

4. Relativistic Sound Speed from the Logarithmic Equation of State

For a relativistic scalar field with Lagrangian density $\mathcal{L} = \rho X - V(\rho)$ where $X \equiv \partial_\mu S \partial^\mu S$ and $\rho = |\psi|^2$, the Hamilton-Jacobi equation in the macroscopic limit reads $X = V'(\rho)$, which allows the density to be integrated out in favor of X . The resulting effective Lagrangian $\mathcal{L}_{eff}(X) = \rho(X)X - V(\rho(X))$ is of k-essence type, with pressure $p = \mathcal{L}_{eff}$ and energy density $\epsilon = 2X\mathcal{L}'_{eff} - \mathcal{L}_{eff}$ [14]. Using the Barceló-Liberati-Visser definition $c_s^2 = dp/d\epsilon$ for relativistic barotropic fluids [15], one finds

$$\frac{c_s^2}{c^2} = \frac{\rho_0 V''(\rho_0)}{\rho_0 V''(\rho_0) + 2V'(\rho_0)}. \quad (10)$$

As a check, the radiation-gas potential $V(\rho) = \frac{1}{2}g\rho^2$ gives $V' = g\rho$, $V'' = g$, $\rho V'' = g\rho = V'$, and Eq. (10) recovers the standard result $c_s^2/c^2 = 1/3$.

For the logarithmic potential $V(\rho) = -b\rho \ln(\rho/\rho_c)$, where the minus sign reflects the attractive (self-binding) interaction in Eq. (6):

$$V' = -b \left[\ln\left(\frac{\rho}{\rho_c}\right) + 1 \right], V'' = -\frac{b}{\rho}, \rho_0 V'' = -b. \quad (11)$$

Substituting (the minus signs cancel in the ratio):

$$\boxed{\frac{c_s^2}{c^2} = \frac{1}{2\ln(\bar{\rho}/\rho_c) + 3}} \quad (12)$$

The coupling b cancels, and c_s depends on the local vacuum density $\bar{\rho}$ through the logarithm. Requiring $c_s = c$ at the background density ρ_∞ fixes $\rho_c = e\rho_\infty$. This is not an arbitrary normalization: the condition $\rho_c = e\rho_\infty$ is equivalent to $V'(\rho_\infty) = 0$, i.e. the background sits at the extremum of the logarithmic potential. The same extremum condition fixes the equation-of-state exponent $\alpha = -1$ at the background and, in the cosmology companion, the equation-of-state value $w = -1$ as the exact coupling-independent ratio P/ε at $V'(\rho_\infty) = 0$. The location $\alpha = -1$ is therefore structurally determined by the vacuum residing at the minimum of its self-interaction potential, not placed there by a choice of scale; a different ρ_c would describe a vacuum displaced from its own potential extremum, which is not the physical ground state. The precise physical value of ρ_∞ —which determines the absolute scale of the framework—is fixed by the gravitational coupling derivation in companion work, where dimensional self-consistency forces ρ_∞ to the Planck mass density; we use the present convention throughout this paper and defer the cosmological identification to dedicated work.

This density dependence is a relativistic effect. In the non-relativistic logarithmic Schrödinger equation, the denominator of (10) is dominated by $|V'(\rho_0)| \approx m_{eff}c^2 \gg |\rho_0 V''|$ (where ρ_0 here denotes the background value of the Lagrangian variable, not the physical mass density), yielding $c_s^2 \approx |\rho_0 V''|/(2m_{eff}c^2) \propto b/(m_{eff}c^2)$, independent of density ^[13]. In the relativistic regime, V' and $\rho_0 V''$ are comparable, and the density dependence survives (Fig. 2). Equation (12) also reveals a hard minimum density: at $\bar{\rho} = \rho_c e^{-3/2}$ the sound speed diverges, and below this threshold $c_s^2 < 0$, signaling breakdown of the hydrodynamic description.




Figure 2. (a) Sound speed vs. density. The relativistic logarithmic fluid (green) has density-dependent c_s , while the real scalar KG and non-relativistic BEC give constant propagation speed. A singularity at $\bar{\rho}/\rho_c = e^{-3/2} \approx 0.22$ marks the minimum physical density. (b) The equation of state exponent $\alpha = d \ln c_s / d \ln \rho$, crossing $\alpha = -1$ at $\bar{\rho} = \rho_\infty$.

The equation of state exponent, which controls the PPN structure, is:

$$\alpha \equiv \frac{d \ln c_s}{d \ln \rho} = -\frac{1}{2 \ln(\bar{\rho}/\rho_c) + 3}, \quad (13)$$

which is dimensionless and unaffected by the choice of units. Equations (10) and (12) are written as the dimensionless ratio c_s^2/c^2 , so that $c_s = c$ at the background density; elsewhere in this paper c_s denotes the physical sound speed in units $c = 1$ (as in the acoustic metric and the accretion equations below), consistent with the same convention. At the cosmological background $\bar{\rho} = \rho_\infty$ (where $\ln(\rho_\infty/\rho_c) = -1$): $\alpha = -1$.

Comparison with non-relativistic SVT

We note that this density-dependent sound speed differs from the result obtained in the non-relativistic superfluid vacuum theory (NR-SVT) program [16], where the first-sound speed of the logarithmic superfluid is found to be density-independent: $c_s^2 = -Db/\hbar$ with b approximately constant. The two results are not in conflict; they describe different regimes. NR-SVT analyzes the non-relativistic logarithmic Schrödinger equation, in which the rest-mass term dominates the dispersion relation and the sound speed reduces to a fixed scalar function of the coupling. The relativistic logarithmic equation (6) used here couples the phase dynamics to the density through the d'Alembertian operator, producing the density dependence of Eq. (12). The density dependence is therefore a relativistic effect,

absent in the non-relativistic limit. The density dependence plays two distinct roles in the framework, which should not be conflated. First, it is what renders the *static* density channel pathological at the background exponent $\alpha = -1$ (Sec. 5), forcing the deflection onto the flow channel; the relativistic starting point is therefore essential, since the non-relativistic limit has density-independent c_s and no such structure. Second, the density dependence governs propagation through large-scale variations of the vacuum density itself—for example, the changes in local c_s , the associated frequency shift, and the refraction experienced by light crossing the boundary of a cosmological underdensity such as the KBC void—a genuinely refractive, cosmological effect treated in the companion cosmology paper [17], whose observability is subject to the Volovik condition (whether atomic and detector scales rescale together with c_s). The deflection by a compact mass measured by Cassini [2], by contrast, occurs at essentially uniform density and is carried by the flow (Sec. 6), not by a refractive c_s gradient.

5. The Static Acoustic Metric and Its Limitations

5.1. PPN parameter from the static metric

For a static irrotational fluid with density $\bar{\rho}(x)$ and sound speed $c_s(x)$, the acoustic metric is [15][18]:

$$g_{tt} = -\bar{\rho}c_s, g_{ij} = \frac{\bar{\rho}}{c_s} \delta_{ij}. \quad (14)$$

With $c_s \propto \bar{\rho}^\alpha$, a small density perturbation $\bar{\rho} = \rho_\infty(1 + \epsilon)$ gives:

$$\frac{\delta g_{tt}}{g_{tt}} = (1 + \alpha)\epsilon, \quad (15)$$

$$\frac{\delta g_{rr}}{g_{rr}} = (1 - \alpha)\epsilon. \quad (16)$$

To extract γ we first rescale the acoustic metric so that the background is Minkowski-normalized, $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$, as required by the standard PPN form [1]. With $c_s(\rho_\infty) = c$ this fixes the conformal factor to unity at the background and the perturbations of Eqs. (15)–(16) become directly comparable to the PPN expansion $g_{00} = -1 + 2U + \mathcal{O}(U^2)$, $g_{ij} = (1 + 2\gamma U)\delta_{ij}$ [1]. Reading $2U$ from the temporal perturbation and $2\gamma U$ from the spatial perturbation gives

$$2U = -\frac{\delta g_{tt}}{g_{tt}} = -(1 + \alpha)\epsilon, 2\gamma U = +\frac{\delta g_{rr}}{g_{rr}} = (1 - \alpha)\epsilon. \quad (17)$$

The relative minus sign is essential and is easily lost. Because g_{tt} is negative while g_{rr} is positive, a positive *fractional* change of g_{tt} corresponds to a *decrease* of $|g_{tt}|$ and hence to a negative contribution to

U , whereas the spatial component carries no such flip. Identifying γ with the bare ratio of fractional changes $(\delta g_{rr}/g_{rr})/(\delta g_{tt}/g_{tt})$ would discard this sign. Taking the ratio of the two expressions in Eq. (17) yields

$$\boxed{\gamma_{static} = \frac{\alpha - 1}{\alpha + 1}} \quad (18)$$

This is the PPN parameter for any static barotropic acoustic metric with equation of state exponent α (Fig. 3), and it carries two immediate consequences.

First, $\gamma_{static} = 1$ has *no* solution: the condition $\alpha - 1 = \alpha + 1$ reduces to $-1 = 1$. No barotropic equation of state whatsoever reproduces the general-relativistic value from the static density channel; $\gamma = 1$ is approached only as $\alpha \rightarrow \pm\infty$ and is never attained. The obstruction is therefore not specific to the logarithmic model.

Second, at $\alpha = 0$ the sound speed is density-independent, the metric (14) is genuinely conformally flat, and Eq. (18) returns $\gamma_{static} = -1$ —precisely the Nordström value quoted in Sec. 1 [3]. This serves as an independent check on the sign, as does the deflection factor

$$1 + \gamma_{static} = \frac{2\alpha}{\alpha + 1}, \quad (19)$$

which vanishes at $\alpha = 0$, in agreement with the elementary observation that a spatially constant c_s produces no refractive bending.




Figure 3. (a) Static PPN parameter $\gamma(\alpha) = (\alpha - 1)/(\alpha + 1)$. The logarithmic fluid at $\bar{\rho} = \rho_\infty$ has $\alpha = -1$, a pole of γ_{static} : the static acoustic metric is ill-defined. The general-relativistic value $\gamma = 1$ (dashed) is never attained for any finite α ; it is approached only as $\alpha \rightarrow \pm\infty$. At $\alpha = 0$ the sound speed is constant, the metric is conformally flat, and $\gamma = -1$, the Nordström value. (b) The deflection factor $(1 + \gamma) = 2\alpha/(\alpha + 1)$, which vanishes at $\alpha = 0$ —constant c_s , hence no refractive bending—and diverges at the pole.

5.2. The static acoustic metric is pathological

Equation (18) reveals a fundamental obstruction. For the logarithmic equation of state at the background density, $\alpha = -1$, and the denominator of γ_{static} vanishes:

$$\gamma_{static} \rightarrow \infty \quad \text{at} \quad \alpha = -1. \quad (20)$$

The static acoustic metric does not merely give the wrong numerical value of γ ; the spatial metric perturbation diverges relative to the temporal one, and no finite weak-field expansion around the static background is possible. Physically, the pole signals that the static configuration is not a stable background for perturbations: an arbitrarily small density gradient produces an arbitrarily large metric response.

It is useful to separate two statements about the static metric that are easily conflated. Writing Eq. (14) as $g_{\mu\nu} = (\bar{\rho}/c_s) \text{diag}(-c_s^2, 1, 1, 1)$ shows that it is conformally flat *only* when c_s is spatially constant. A density gradient makes $|g_{tt}|/g_{rr} = c_s^2(x)$ position-dependent, so the static metric of an inhomogeneous condensate is *not* conformally flat, and it does deflect light. What holds in general is the weaker statement that the conformal factor $\bar{\rho}/c_s$ cancels from the null condition entirely, so that null propagation is governed by $c_s(x)$ alone through the refractive index $n = c/c_s$, with $n - 1 = -\alpha\epsilon$ at first order.

The static channel is obstructed twice over. The logarithmic model sits at the pole $\alpha = -1$, where no finite weak-field expansion exists. Away from the pole, Eq. (18) never reaches $\gamma = 1$ for any α , so no alternative barotropic equation of state repairs the static channel either: at the special point $\alpha = 0$ one recovers the conformally flat Nordström case, with $\gamma = -1$ and vanishing deflection, and elsewhere the ratio of spatial to temporal curvature is wrong by construction. Any consistent treatment must therefore regularize by introducing dynamical content beyond the static density channel; this is the role of the Painlevé–Gullstrand flow developed in Sec. 6.

6. The Flowing Acoustic Metric and the Path to $\gamma = 1$

6.1. The Painlevé–Gullstrand structure

The static metric exhausts only one of the two degrees of freedom provided by the Madelung decomposition. The second—the macroscopic flow velocity $\mathbf{v} = \nabla S/m_{eff}$ —contributes off-diagonal terms. For a spherically symmetric radial inflow $v(r)$ in a fluid with $\bar{\rho}/c_s \approx \text{const}$ and $c_s \approx c$, the acoustic metric takes the Painlevé–Gullstrand form ^{[18][19][20]}:

$$ds^2 = -c^2 dt^2 + (dr + v(r)dt)^2 + r^2 d\Omega^2. \quad (21)$$

If the flow velocity satisfies the free-fall profile:

$$v(r) = \sqrt{\frac{2G_{eff}M}{r}}, \quad (22)$$

then the standard coordinate transformation $dt_s = dt - vdr/(c^2 - v^2)$ converts (21) to the Schwarzschild metric:

$$ds^2 = -\left(1 - \frac{2G_{eff}M}{rc^2}\right) c^2 dt_s^2 + \frac{dr^2}{1 - 2G_{eff}M/(rc^2)} + r^2 d\Omega^2, \quad (23)$$

which gives $\gamma = 1$ exactly. The off-diagonal flow term breaks the conformal flatness of the static metric, introducing genuine spatial curvature through the $g_{rr} = (1 - v^2/c^2)^{-1}$ factor. That this specific velocity profile is not imposed by fiat but follows from a Bernoulli self-consistency condition is established in Sec. 7.2. The physical mechanism is frame-dragging by the inflowing vacuum: light propagating against the current is slowed (increased effective refractive index), while the spatial geometry is distorted by the kinetic energy of the flow. Both effects contribute to the deflection, and their combined weight gives the full GR value.

6.2. Self-consistency test: Bondi accretion

The Painlevé–Gullstrand argument assumes $v(r) = \sqrt{2G_{eff}M/r} \propto r^{-1/2}$. We now ask: does this profile emerge self-consistently from the fluid equations with the logarithmic equation of state?

For steady-state spherical accretion, the continuity equation (8) gives $4\pi r^2 \bar{\rho} v = \dot{M} = \text{const.}$ If $\bar{\rho}$ is approximately constant (required for the Painlevé–Gullstrand form), then:

$$v(r) \propto \frac{1}{r^2} \quad (\text{continuity with } \bar{\rho} \approx \text{const}). \quad (24)$$

This is incompatible with the required $v \propto r^{-1/2}$. The discrepancy grows as $(r_s/r)^{3/2}$, where $r_s = GM/(2c^2)$ is the sonic radius.

We verify this by solving the full Bondi accretion equations [21]:

$$\frac{dv}{dr} = \frac{v(2c_s^2/r - GM/r^2)}{v^2 - c_s^2} \quad (25)$$

with $c_s^2(\bar{\rho})$ given by Eq. (12) and $\bar{\rho}$ from continuity. The solution, passing through the sonic point ($v = c_s$ at $r = r_s$), is shown in Fig. 4.




Figure 4. (a) Bondi accretion velocity (solid gold) vs. the Painlevé–Gullstrand requirement

$v = \sqrt{2GM/r}$ (dashed red). In the subsonic far field, $v_{\text{Bondi}} \propto r^{-2}$, falling far below the $r^{-1/2}$ profile.

(b) Ratio $v_{\text{Bondi}}/v_{\text{PG}}$. At the solar limb ($r \sim 5 \times 10^5 GM/c^2$), the ratio is $\approx 2.8 \times 10^{-9}$.

This establishes that single-soliton Bondi accretion cannot produce the PG profile required by the acoustic-metric consistency argument. The resolution is that the PG profile must arise from collective, many-soliton dynamics, not from single-source accretion—a point we return to in Sec. 7.2.

7. Discussion

7.1. The circularity objection

A natural objection: by placing c in the D'Alembertian of Eq. (6), we assumed Lorentz invariance. Is $\gamma = 1$ then circular?

No. The D'Alembertian alone guarantees only that the asymptotic propagation speed is c . As shown in Sec. 2, the real scalar equation $\square D + V'(D) = 0$ also contains c in its D'Alembertian, yet gives $v = c = \text{const}$ regardless of background, yielding $\gamma = 0$. Lorentz invariance is necessary but not sufficient for $\gamma = 1$.

What promotes γ from 0 to 1 is the combination of three distinct structural features, none of which is present in the real-scalar theory:

1. *The Madelung decomposition* promotes the field content from one real degree of freedom (D) to two real degrees of freedom (density ρ and phase S), breaking the rigidity of the principal symbol.
2. *The logarithmic nonlinearity*, in the *relativistic* regime, produces a density-dependent sound speed with $\alpha = -1$ at the background (Eq. 13).
3. *The Painlevé-Gullstrand flow* provides a second dynamical channel (the macroscopic flow velocity) that regularizes the static-metric divergence at $\alpha = -1$ and delivers $\gamma = 1$ for any barotropic equation of state.

Each ingredient is individually insufficient: the real-scalar KG with the same D'Alembertian gives $\gamma = 0$; the Madelung decomposition alone in the non-relativistic limit gives $\alpha = 0$ and thus no refractive bending; the static acoustic metric alone at $\alpha = -1$ is pathological. Lorentz invariance is the foundation; the complex field structure, the relativistic log nonlinearity, and the macroscopic flow together build the house.

7.2. The logical chain

Our results establish a clear logical chain:

The kinematic trap is absolute

Any real scalar field equation with a D'Alembertian kinetic term gives $\gamma = 0$, regardless of the self-interaction potential (Sec. 2). This is a property of the equation's characteristic surfaces, not of any

particular solution.

The Madelung formulation breaks the trap

The complex field's Madelung decomposition introduces two degrees of freedom—density ρ and flow velocity $\mathbf{v}$ —that are hidden in the wave equation form. The relativistic logarithmic equation of state yields a density-dependent sound speed (Eq. 12), which is the prerequisite for any refractive bending mechanism.

The static acoustic metric is pathological

The PPN parameter of a static barotropic acoustic metric is $\gamma = (\alpha - 1)/(\alpha + 1)$ (Eq. 18), which equals unity for no value of α : no barotropic equation of state reproduces general-relativistic deflection from the static density channel. For the logarithmic equation of state at the background density, $\alpha = -1$, placing γ_{static} exactly at the pole. The static acoustic metric is therefore mathematically ill-defined, not merely numerically wrong.

Background flow is necessary and regularizes the pathology

The Painlevé-Gullstrand form of the flowing acoustic metric reproduces the Schwarzschild geometry and $\gamma = 1$ exactly, provided $v(r) = \sqrt{2G_{eff}M/r}$ (Sec. 6). The off-diagonal flow term $-2vdrdt$ cannot be removed by a conformal rescaling and carries the full $\gamma = 1$ result for any barotropic equation of state, independent of α . In this way the flow regularizes the divergence of the static case: the dynamical content that the static metric could not carry in the (pathological) density channel is transferred to the (well-defined) flow channel.

The PG flow is the only self-consistent solution

In the superfluid vacuum framework, the gravitational potential is not externally specified—it is the acoustic metric. This imposes a self-consistency requirement: the density perturbation $\epsilon = \delta\rho/\rho_0$ and flow velocity $v(r)$ must produce an acoustic metric whose effective Newtonian potential $\Phi_{eff} = -\frac{1}{2}\delta g_{tt}/g_{tt}$ matches the potential that generated ϵ and v in the first place.

For the static solution (Solution A: $v = 0, \epsilon \neq 0$), the divergence of γ_{static} at $\alpha = -1$ means that no finite density perturbation ϵ produces a finite weak-field metric: the self-consistency equation admits no regular solution in this channel.

For the flowing solution (Solution B: $v = \sqrt{2G_{eff}M/r}$, $\rho \approx \rho_0$), the Bernoulli equation gives $\frac{1}{2}v^2 + \Phi_{eff} = 0$ at leading order, and the acoustic metric in Schwarzschild coordinates gives $g_{tt} \approx -(1 - 2U)$, $g_{rr} \approx (1 + 2U)$, yielding $\Phi_{eff} = -U$ and $\gamma = 1$. The Bernoulli relation $v^2 = 2U$ then closes the loop: Φ_{eff} produces the flow that produces Φ_{eff} . This solution is self-consistent for any barotropic equation of state, because the Painlevé-Gullstrand acoustic metric is identically the Schwarzschild metric regardless of the specific form of $c_s(\rho)$.

The flow carries the potential, not the density well

Linearizing the Hamilton-Jacobi equation (9) about the background with $|\mathbf{v}| \ll c$ and $b > 0$, and using the mostly-plus signature fixed by the d'Alembertian, gives $\ln(\rho/\rho_\infty) = -(m_{eff}^2/b\hbar_{eff}^2)|\mathbf{v}|^2$ to leading order—a genuine density depletion $\delta\rho \equiv \rho_\infty - \rho > 0$ in the well, with $\delta\rho/\rho_\infty \propto v^2$. This depletion is real but does not source gravity. Expanding the temporal acoustic-metric component with $c_s \propto \rho^\alpha$ gives $\delta g_{tt}/g_{tt} = (1 + \alpha)\epsilon - v^2/c_s^2$, and at the background exponent $\alpha = -1$ the density coefficient $(1 + \alpha)$ vanishes, so the depletion decouples from g_{tt} entirely. The potential is therefore carried by the flow channel alone, $\Phi_{eff} = -\frac{1}{2}v^2$, and requiring this to reproduce the Newtonian potential $-G_{eff}M/r$ forces $v(r) = \sqrt{2G_{eff}M/r}$. Because $\Phi_{eff} \propto v^2$, a $1/r$ potential admits only $v^2 \propto 1/r$: the Painlevé-Gullstrand profile is the unique flow that carries a Newtonian potential. This argument never invokes a Poisson density well, and so remains consistent at the $\alpha = -1$ pole—unlike a density-well derivation, which the pole would invalidate.

The continuity constraint

The radially inward PG flow $\mathbf{v} = -\sqrt{2GM/r}\hat{\mathbf{r}}$ has $\nabla \cdot \mathbf{v} = -\frac{3}{2}\sqrt{2GM}r^{-3/2} \neq 0$, corresponding to condensate accumulation $\partial\rho/\partial t = -\rho_0\nabla \cdot \mathbf{v} > 0$. The accumulation rate is small: at the solar limb, $|\nabla \cdot \mathbf{v}|^{-1} \sim 10^3$ s, but this timescale applies to fractional changes $\delta\rho/\rho_0 = 2U \sim 10^{-6}$, so the absolute density change per observation epoch is negligible. The PG flow is a valid quasi-steady approximation on all timescales short compared to the gravitational collapse time. A fully steady-state treatment requires a mechanism that redirects the inflowing condensate without local accumulation; candidate mechanisms include two-fluid counterflow, vortex-core absorption, and geometric redirection into non-radial outflow. The choice of mechanism is deferred to future work.

Single-particle vs. collective flow

The Bondi accretion solution for the logarithmic equation of state does not produce the required velocity profile in the far field. Continuity enforces $v \propto r^{-2}$ for nearly-incompressible steady flow, while $\gamma = 1$ requires $v \propto r^{-1/2}$. This discrepancy applies to single-soliton configurations. However, the self-consistency argument above does not depend on the single-particle velocity field. The PG flow must arise as a *collective* effect of $N \sim M/m_{\text{particle}}$ solitons: individual particles contribute microscopic density perturbations that sum to the macroscopic potential $\Phi = -GM/r$, and the resulting vacuum inflow adjusts to maintain self-consistency of the acoustic metric. In this picture, the Bondi calculation describes the flow around a single vortex core (irrelevant at astrophysical distances), while the PG flow describes the macroscopic vacuum response to the collective mass distribution. The relationship between the microscopic and macroscopic descriptions—analogue to the Feynman-Onsager relation in rotating superfluids—remains an open problem.

Two open problems remain. First, deriving the macroscopic PG flow from the microscopic soliton dynamics: showing that a collection of N vortex-Gaussons in the logarithmic condensate produces a collective vacuum inflow $v = \sqrt{2GM/r}$ as a mean-field effect, analogous to how a lattice of quantized vortices in superfluid helium produces solid-body rotation [20]. Second, establishing the steady-state mechanism: identifying whether the two-fluid counterflow (superfluid infall balanced by normal-component outflow) or vortex-core absorption provides the sink term required for continuity. The first problem is amenable to a Thomas-Fermi mean-field calculation; the second requires understanding the quantum depletion of the condensate in the presence of topological defects.

7.3. Relation to prior work

The superfluid vacuum program of Zloschastiev [16][22][23] derives emergent spacetime metrics from the logarithmic condensate but does not compute PPN parameters explicitly. The analogue gravity program of Barceló, Liberati, and Visser [24][15] proves that fluid perturbations propagate on acoustic metrics but does not construct specific models matching solar system tests, and its experimental realizations—sonic horizons and Hawking-like emission in Bose-Einstein condensates [25][26], and rotating black-hole analogues in condensed and photon-fluid systems [27][28]—probe the kinematics of these effective metrics rather than their match to gravitational observations. The view of Lorentz invariance and relativistic dynamics as emergent low-energy features of a condensed substrate [29] is the conceptual

setting for the present preferred-frame treatment. The present work connects these programs by deriving the explicit PPN constraints and identifying the gap between the static and flowing acoustic metrics.

The flowing acoustic metric's equivalence to the Painlevé–Gullstrand form of Schwarzschild was noted by Unruh [18] and developed extensively by Visser [19] and Volovik [20]. Our contribution is to test this identification against self-consistent fluid dynamics, revealing that the velocity profile is not automatically produced.

The construction here is relativistic: the effective metric arises from a relativistic logarithmic field rather than a non-relativistic condensate, which is what makes the sound speed density-dependent (Sec. 4) and the cosmological-propagation regime nontrivial. Curved and expanding effective geometries have been engineered in Bose-Einstein condensates [30], providing a laboratory analogue of the large-scale density-variation regime discussed above.

7.4. *Observational tests passed by the framework*

What "light" is, the dispersion relation, and achromaticity bounds

Within the framework, light is not separate from the condensate: photons are long-wavelength excitations of the same relativistic logarithmic Klein-Gordon field whose Madelung decomposition produces the acoustic metric. The framework is a single-field theory; there is no independent electromagnetic medium "on top" of the vacuum. The same complex order parameter ψ that gives the acoustic metric through its hydrodynamic envelope also carries photonic excitations as its transverse phonon-like modes. The U(1) gauge structure of electrodynamics is an internal symmetry of ψ , with the derivation of Maxwell's equations from the gauged log-KG field forming a separate manuscript of the program. The relevant dispersion relation follows directly from the linearized log-KG equation.

Linearizing the log-KG equation about the background ($\rho = \rho_0$, $S_0 = -m_{eff}c^2t$) yields two Bogoliubov branches. Because the d'Alembertian is second order in both space and time, each is strictly linear in k^2 :

$$\omega^2 = c^2 k^2 \quad (\text{gapless}), \quad \omega^2 = c^2 k^2 + 2bc^2 \quad (\text{gapped}). \quad (26)$$

The gapless branch is the photonic excitation: its dispersion is *exactly* $\omega = ck$, with no higher-order correction. (This is unlike the non-relativistic Bogoliubov spectrum, whose k^4 term arises from the Schrödinger equation's first-order time derivative; the relativistic kinetic operator admits no such term.) The second branch is a massive mode gapped at $\omega_{gap} = c\sqrt{2b}$ —the healing/Planck scale—and is

irrelevant to low-energy electromagnetic propagation. Consequently the speed of light, gravitational waves, and acoustic phonons all equal c at the background *to all orders*, in exact agreement with the GW170817 constraint $|c_{gw} - c|/c < 10^{-15}$ [31].

The continuum theory therefore predicts exactly achromatic light deflection and Shapiro delay: the deflection of Eq. (1) is frequency-independent at every order, so Cassini Shapiro-delay timing [2], gravitational lensing, and gamma-ray-burst time-of-flight bounds are satisfied identically rather than by numerical suppression. Any vacuum dispersion would originate not in the continuum log-KG but in the discrete, granular structure of the condensate at the healing scale $\xi \sim \ell_P$ (fixed by the framework’s gravitational-coupling derivation in companion work)—a genuinely Planck-suppressed ultraviolet effect outside the scope of the present continuum treatment. Whether such granularity induces a quadratic correction $\Delta v/c \sim (E/E_P)^2$ consistent with current Lorentz-invariance bounds is a question for a microscopic completion of the framework, left to dedicated work.

It is essential to the light-bending result that this gapless photon, like every excitation of the condensate, propagates on the same flowing acoustic metric: its kinetic term is built with the acoustic metric, so the photon is advected by the background flow $v(r)$ exactly as the c_s -phonon is. The deflection of Sec. 6 is this advection, and because the off-diagonal flow term $-2vdrdt$ acts on v^2 alone—common to all excitations and independent of propagation speed—the $\omega = ck$ photon and the c_s -phonon acquire the same potential $\Phi_{eff} = -\frac{1}{2}v^2$ and the same $\gamma = 1$. Exact achromaticity and $\gamma = 1$ are thus the same statement: a speed-independent (advective) deflection acts identically at all frequencies.

The PG flow and preferred-frame tests

The Painlevé-Gullstrand flow $v(r) = \sqrt{2GM/r}$ takes the value $\sim 10^{-3}c$ at the solar limb, raising the question of why high-precision local experiments do not detect a Sagnac-like “ether wind.” The answer is the same structural feature that delivers $\gamma = 1$. The PG metric is identically Schwarzschild — the transformation from PG coordinates to standard Schwarzschild coordinates is a passive change of variables that cannot alter any invariant observable. The PPN preferred-frame parameters $\alpha_1, \alpha_2, \alpha_3$ vanish in Schwarzschild geometry by construction [1], and they continue to vanish in any coordinate system describing the same geometry, including the PG flow representation. The framework therefore predicts $\alpha_1 = \alpha_2 = \alpha_3 = 0$ at the same order as $\gamma = 1$: as a direct consequence of the PG-Schwarzschild equivalence, not as a separate calculation.

This resolves the apparent puzzle. The PG flow is a real macroscopic feature of the framework — the condensate genuinely supports the velocity field $v(r)$ around the Sun — but its observable consequences are precisely those of Schwarzschild gravity. Laboratory rulers and clocks situated in the solar system are in free fall in the same gravitational field; by construction, $v(r)$ is the free-fall velocity at radius r . An Earth-bound experiment cannot detect the underlying flow for the same reason a freely-falling observer in general relativity cannot detect the gravitational field they are falling through, expressed here in the language of an explicit medium picture rather than a geometric one. The framework’s preferred rest frame is identified observationally with the CMB rest frame (in which the CMB dipole vanishes); this defines the global rest frame against which the macroscopic vacuum dynamics are described, while local physics in the solar system reproduces the standard PPN bounds. Refinements at higher post-Newtonian order, where the medium picture and the geometric picture could in principle diverge, are not derived in this work.

The weak equivalence principle

The MICROSCOPE measurement of the Eötvös parameter, $\eta(Ti, Pt) = [-1.5 \pm 2.3(stat) \pm 1.5(syst)] \times 10^{-15}$ [32], is the tightest current bound on equivalence-principle violation. In any theory in which gravitation arises from coupling to a substrate rather than from a postulated geometric metric, this bound constrains composition-dependent terms in the matter-substrate coupling. The framework’s leading-order universality, established above through the PG-Schwarzschild equivalence, automatically passes this test. The remaining theoretical question is whether composition-dependent corrections to the soliton-condensate coupling — arising from differences in electromagnetic binding energy, nuclear binding energy, or spin-orbit structure between test materials — generate an η_{12} within the MICROSCOPE confidence interval, or whether they predict a specific non-zero value that future experiments could resolve. Computing η_{Pt-Ti} explicitly from the framework’s matter-substrate coupling is deferred to dedicated work; it requires the U(1) gauging of the condensate and the soliton internal-energy decomposition, both developed in companion follow-up manuscripts of this program.

Shapiro time delay

The recovery of $\gamma = 1$ via the Painlevé–Gullstrand structure automatically reproduces the Shapiro time delay. Once the acoustic metric in the PG frame is shown to be identically Schwarzschild (Sec. 6), all

weak-field tests parameterized by the same PPN coefficients $\{\gamma, \beta, \alpha_i, \dots\}$ [11] are reproduced together as a consequence: the coordinate transformation from PG to standard Schwarzschild coordinates is a passive change of variables that cannot alter any invariant observable. The Cassini measurement of γ from Shapiro delay timing [12] is therefore reproduced without additional input, with no framework-specific dispersion in the long-wavelength limit (see also discussion of the electromagnetic sector below).

Asymptotic constraints

The GW170817 constraint ($|c_{gw} - c|/c < 10^{-15}$ [31]) requires that the asymptotic sound speed equal c exactly, fixing a combination of the equation of state parameters. The second PPN parameter β , controlling perihelion precession, takes the GR value $\beta = 1$ by the same coordinate-equivalence argument as γ . The full post-Newtonian expansion to 2PN and higher orders, which probes the framework beyond the linearized acoustic-metric regime, is left for future work.

7.5. Limitations and open problems

Conformal flatness and strong fields

The acoustic metric (14) is conformally flat ($g_{ij} \propto \delta_{ij}$). General relativity admits nontrivial Weyl curvature. The present framework reproduces the weak-field PPN limit but cannot describe strong-field phenomena: binary pulsar orbital decay [33], black hole ringdown, or nonlinear gravitational wave dynamics. Whether extensions incorporating vorticity or background flow can access the strong-field regime is an open question.

The quantum potential

The Bohm potential $Q = \hbar^2 \nabla^2 \sqrt{\rho} / (2m \sqrt{\rho})$ was neglected in the macroscopic limit. At the soliton coherence scale, Q becomes important and governs quantum-mechanical particle behavior. A complete theory must show a smooth transition between quantum and gravitational regimes—analogueous to the phonon-to-hydrodynamic crossover in superfluid helium [20].

8. Conclusion

We have shown that the failure of scalar gravity to produce the observed light bending is a property of the wave equation representation, not of scalar fields per se. The Madelung decomposition of a relativistic

logarithmic superfluid introduces two degrees of freedom—a density-dependent sound speed and a macroscopic flow velocity—that together provide the structural ingredients needed for $\gamma = 1$.

The static acoustic metric of the logarithmic superfluid is mathematically pathological: the equation of state exponent $\alpha = -1$ at the background density places the static PPN parameter $\gamma_{static} = (\alpha - 1)/(\alpha + 1)$ exactly at the pole. No finite weak-field expansion around the static background exists, and no other barotropic equation of state would rescue the static channel, since $\gamma_{static} = 1$ has no solution. By contrast, the Painlevé–Gullstrand flow $v = \sqrt{2G_{eff}M/r}$ regularizes this pathology: the resulting acoustic metric is identically Schwarzschild in Painlevé–Gullstrand coordinates, and $\gamma = 1$ follows for *any* barotropic equation of state. This selects the flowing solution as the unique physically admissible weak-field configuration.

The Bondi accretion calculation shows that a single spherically symmetric source does not produce the PG profile: far from the sonic radius the flow decays as r^{-2} , falling nearly nine orders of magnitude below the $r^{-1/2}$ requirement at the solar limb. The remaining open problems are therefore to derive the macroscopic PG flow as a collective mean-field effect of the microscopic soliton dynamics—analogueous to the Feynman–Onsager relation in rotating superfluids—and to identify the steady-state mechanism (two-fluid counterflow or vortex-core absorption) that resolves the continuity constraint. Both are amenable to analytical treatment and form the subject of forthcoming work.

Statements and Declarations

Data and Code Availability

The Klein–Gordon wave packet simulation code (Figure 1) and the Bondi accretion integration code (Figure 4) are available at https://github.com/kulangiev/kulangiev.github.io/tree/main/papers/paper_1 under Creative Commons Attribution 4.0 (CC BY 4.0) licence. All numerical parameters, initial conditions, and post-processing routines used to generate the figures are documented in the repository.

Acknowledgments

The author thanks C. Barceló, S. Liberati, and M. Visser for the acoustic metric framework. The methodological stance of this paper—treating the order parameter as a real classical field whose hydrodynamic behavior underlies emergent gravitational phenomena—is indebted to the Madelung–de Broglie–Bohm tradition, particularly as developed in P. R. Holland’s exposition of the quantum theory

of motion. Claude (Anthropic) was used as an editorial and computational tool during manuscript preparation. All scientific content, equations, and interpretations are the author's own.

References

1. ^{a, b, c, d, e, f}Will CM (2014). "The Confrontation between General Relativity and Experiment." *Living Rev Relativ.* **17**:4. doi:[10.12942/lrr-2014-4](https://doi.org/10.12942/lrr-2014-4).
2. ^{a, b, c, d}Bertotti B, Iess L, Tortora P (2003). "A Test of General Relativity Using Radio Links with the Cassini Spacecraft." *Nature.* **425**:374.
3. ^{a, b}Nordström G (1913). "Zur Theorie der Gravitation vom Standpunkt des Relativitätsprinzips" [On the Theory of Gravitation from the Standpoint of the Relativity Principle]. *Ann Phys (Leipzig).* **347**:533.
4. ^aEinstein A (1911). "Über den Einfluß der Schwerkraft auf die Ausbreitung des Lichtes" [On the Influence of Gravity on the Propagation of Light]. *Ann Phys (Leipzig).* **340**:898.
5. ^{a, b}Rosen G (1969). "Dilatation Covariance and Exact Solutions in Local Relativistic Field Theories." *Phys Rev.* **183**:1186.
6. ^{a, b}Bartkowski K, Górka P (2008). "One-Dimensional Klein-Gordon Equation with Logarithmic Nonlinearities." *J Phys A: Math Theor.* **41**:355201.
7. ^{a, b, c}Białynicki-Birula I, Mycielski J (1979). "Gaussons: Solitons of the Logarithmic Schrödinger Equation." *Phys Scr.* **20**:539.
8. ^{a, b}Madelung E (1927). "Quantentheorie in hydrodynamischer Form" [Quantum Theory in Hydrodynamic Form]. *Z Phys.* **40**:322.
9. ^{a, b}Bohm D (1952). "A Suggested Interpretation of the Quantum Theory in Terms of "Hidden" Variables. I." *Phys Rev.* **85**:166.
10. ^{a, b}Holland PR (1993). *The Quantum Theory of Motion*. Cambridge University Press.
11. ^aBrans C, Dicke RH (1961). "Mach's Principle and a Relativistic Theory of Gravitation." *Phys Rev.* **124**:925.
12. ^ade Broglie L (1927). "La mécanique ondulatoire et la structure atomique de la matière et du rayonnement" [Wave Mechanics and the Atomic Structure of Matter and Radiation]. *J Phys Radium.* **8**:225.
13. ^{a, b}Avdeenkov AV, Zloshchastiev KG (2011). "Quantum Bose Liquids with Logarithmic Nonlinearity: Self-Sustainability and Emergence of Spatial Extent." *J Phys B.* **44**:195303.
14. ^aBabichev E, Mukhanov V, Vikman A (2008). "k-Essence, Superluminal Propagation, Causality and Emergent Geometry." *JHEP.* **02**:101.

15. ^{a, b, c}Barceló C, Liberati S, Visser M (2005). "Analogue Gravity." *Living Rev Relativ.* **8**:12.
16. ^{a, b}Zloshchastiev KG (2023). "Derivation of Emergent Spacetime Metric, Gravitational Potential and Speed of Light in Superfluid Vacuum Theory." *Universe.* **9**:234.
17. ^ΔKulangiev BB (2026). "Cosmological Implications of the Logarithmic Superfluid Vacuum." Preprint. <https://kulangiev.github.io/#paper3>.
18. ^{a, b, c}Unruh WG (1981). "Experimental Black-Hole Evaporation?" *Phys Rev Lett.* **46**:1351.
19. ^{a, b}Visser M (1998). "Acoustic Black Holes: Horizons, Ergospheres and Hawking Radiation." *Class Quant Grav.* **15**:1767.
20. ^{a, b, c, d}Volovik GE (2003). *The Universe in a Helium Droplet.* Oxford.
21. ^ΔBondi H (1952). "On Spherically Symmetrical Accretion." *Mon Not Roy Astron Soc.* **112**:195.
22. ^ΔZloshchastiev KG (2020). "An Alternative to Dark Matter and Dark Energy: Scale-Dependent Gravity in Superfluid Vacuum Theory." *Universe.* **6**:180.
23. ^ΔZloshchastiev KG (2011). "Spontaneous Symmetry Breaking and Mass Generation as Built-In Phenomena in Logarithmic Nonlinear Quantum Theory." *Acta Phys Polon B.* **42**:261.
24. ^ΔBarceló C, Liberati S, Visser M (2001). "Analogue Models for FRW Cosmologies." *Class Quant Grav.* **18**:1137.
25. ^Δ[authors] (2010). "Realization of a Sonic Black Hole Analog in a Bose-Einstein Condensate." *Phys Rev Lett.* **105**:240401.
26. ^Δ[authors] (2009). "Numerical Observation of Hawking Radiation from Acoustic Black Holes in Atomic Bose-Einstein Condensates." *Phys Rev A.* **80**:043601.
27. ^Δ[authors] (2018). "Rotating Black Hole Geometries in a Two-Dimensional Photon Superfluid." *Optica.* **5**:1099.
28. ^Δ[authors] (2019). "Quantum Analogue of a Kerr Black Hole and the Penrose Effect in a Bose-Einstein Condensate." *Phys Rev B.* **99**:214511.
29. ^Δ[authors] (2003). "Emergent Relativity." *Int J Mod Phys A.* **18**:831.
30. ^Δ[authors] (2022). "Curved and Expanding Spacetime Geometries in Bose-Einstein Condensates." *Phys Rev A.* **106**:033313.
31. ^{a, b}Abbott BP et al. (2017). "Gravitational Waves and Gamma-Rays from a Binary Neutron Star Merger: GW170817 and GRB 170817A." *Astrophys J Lett.* **848**:L13.
32. ^ΔTouboul P et al. (MICROSCOPE Collaboration) (2022). "MICROSCOPE Mission: Final Results of the Test of the Equivalence Principle." *Phys Rev Lett.* **129**:121102.

33. ^ΔTaylor JH, Fowler LA, McCulloch PM (1979). "Measurements of General Relativistic Effects in the Binary Pulsar PSR 1913+16." *Nature*. 277:437.

Declarations

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.