

v1: 10 March 2024

Research Article

Evaluation of Diabetes Risk Score Tool for Detecting Undiagnosed Type 2 Diabetes Mellitus in Referral Clinics at Primary Health Care Centers in Sudan: A Cross-Sectional Study

Preprinted: 3 January 2024

Peer-approved: 10 March 2024

© The Author(s) 2024. This is an Open Access article under the CC BY 4.0 license.

Qeios, Vol. 6 (2024)
ISSN: 2632-3834

Sahar M. Fadl¹, Ammar H. Abbas², Manal A. Elemam^{3,4}, Omer Kheir⁵

1. Diabetes Prevention and Control Program General Directorate of Health Programs and Non-Communicable Disease, Ministry of Health, Riyadh, Saudi Arabia; 2. Preventive Medicine, Prince Sultan Military Medical City (PSMMC), Riyadh, Saudi Arabia; 3. Ministry of Health Saudi Arabia, Riyadh, Saudi Arabia; 4. General Directorate of Health Programs and Non-Communicable Disease, Ministry of Health, Riyadh, Saudi Arabia; 5. Johns Hopkins Aramco Healthcare, Saudi Arabia

Introduction: The prevalence of diabetes in Sudan is increasing, but suitable risk assessment and screening tools to identify at-risk individuals are lacking.

Objective: To evaluate the Diabetes Risk Score (DRS) tool for detecting undiagnosed type 2 diabetes mellitus.

Methods: In this cross-sectional study, 214 individuals were recruited from primary health care referral centers in Khartoum State. Attendees were interviewed to fill out the DRS questionnaire. Random blood glucose and glycosylated hemoglobin (HbA1c) were tested. Descriptive statistics and sensitivity analyses were used to test the applicability of the DRS in Sudan.

Results: The prevalence of undiagnosed Type 2 diabetes was found to be 14%. Regarding blood tests, random blood glucose was normal in 93% of the participants (cutoff: ≤ 140 mg/dl). The HbA1c result was normal in 86% of the participants (cutoff: $< 6.4\%$). The DRS was high in 40.2% (cutoff: ≥ 33), while 59.8% were considered to have moderate to low risk. The DRS had a sensitivity of 83.33% and a specificity of 66.85%. The positive and negative predictive values were 29.07% and 96.09%, respectively. The area under the curve (AUC) for detecting undiagnosed Type 2 diabetes was 0.751 (95% confidence interval: 0.662–0.840).

Conclusion: The DRS tool was found to be applicable with reference to the HbA1c test for predicting undiagnosed Type 2 diabetes.

Correspondence: papers@team.qeios.com — Qeios will forward to the authors

Introduction

Diabetes mellitus (DM) is a serious chronic disease that has emerged as a worldwide public health problem. It is considered one of the four priority non-communicable diseases (NCDs) requiring global action. The incidence and prevalence of diabetes have been steadily increasing [1]. Notably, the Middle East and North Africa (MENA) region has the second-highest rate of diabetes among all International Diabetes Federation (IDF) regions. The prevalence of diabetes in this region is 9.2%, but nearly half of the cases (49%) are undiagnosed. The IDF has announced that the number of people with diabetes worldwide will reach 693 million by 2045 [2].

According to the Central Intelligence Agency (CIA), the total estimated population of Sudan is 43,087,468. Khartoum State has three major cities with a combined population of 7,380,479 ^[3]. According to IDF 2019 statistics, Sudan is one of the 19 territories of the IDF-MENA region, and the prevalence of diabetes among Sudanese adults is 10.9% ^[4].

A recent study found that the incidence rates of type 1 diabetes mellitus in Khartoum State among children under the age of five and those aged 16 to 19 were 8.4 and 7.7/100,000, respectively.

The overall prevalence of diabetes was 6.0% in 2016, according to the STEPwise Surveillance (STEPS) survey of non-communicable diseases. In Khartoum State, the prevalence was 11.6% ^[5].

A cross-sectional survey in Gadarif State measured the prevalence of newly diagnosed diabetes as 10.0% ^[6]. Diabetes prevalence was significantly higher in urban areas than in rural areas. The prevalence of undiagnosed diabetes in North Africa is high compared to the overall prevalence of diabetes, ranging from 18 to 75% ^[7]. Several studies have recommended active screening for individuals older than 45 years, as well as those with hypertension or unexplained weight loss ^[8]. In settings with poor resources, selective multistage screening is encouraged by the World Health Organization (WHO). The implementation of the Package of Essential Non-communicable Disease Interventions (PEN) at the primary care level includes recommendations for the screening of individuals older than 40 years, as well as younger individuals who have risk factors ^[9].

WHO PEN is the minimal standard for NCDs that improves national capacity to incorporate and scale up care of heart disease, stroke, cardiovascular risk, diabetes, cancer, asthma, and chronic obstructive pulmonary disease in primary health care in low-resource settings. PEN highlights simplified clinical protocols adapted for primary healthcare settings, enabling health staff to provide prompt, effective, and complete care and promote behavior change for those at risk or currently afflicted by NCD. ^[10]

Screening for type 2 DM (T2DM) involves a combination of risk assessment and biochemical tests to confirm the diagnosis of new cases. Screening tests are combined to improve performance and provide more accurate results. Screening tests for T2DM can be applied separately (a questionnaire followed by blood glucose measurement if a high-risk score is reached) or simultaneously ^[11]. Screening tests are usually followed by diagnostic tests (fasting blood glucose and/or oral glucose tolerance tests) using standard criteria to make a definitive diagnosis.

Several potential approaches are available to screen for diabetes ^[11]. The entire population may be screened, or selective or targeted screening can be performed for subgroups who have already been identified as having a relatively high risk concerning age, body weight, ethnic origin, etc. Opportunistic screening may also be carried out at a time when people meet health care professionals for reasons other than diabetes.

Sudan still has weaknesses regarding early detection and prevention strategies. The strategy that is currently being implemented relies on diagnostic criteria for diabetes and laboratory confirmation through healthcare providers according to the Sudan DM Guidelines of 2011, which were developed by the Federal Ministry of Health Sudan (FMoH) and the NCD Directorate. There are numerous advantages to implementing a simple and non-invasive screening tool for the early detection of borderline and undiscovered diabetic cases. Such simple interventions could reduce healthcare expenditures by either reversing the occurrence of the disease or delaying the appearance of disease complications. Thus, the objectives of this study were to evaluate the Diabetes Risk Score (DRS) tool for detecting T2DM among undiagnosed individuals in a Sudanese setting based on its sensitivity and receiver operating characteristic curve.

Materials and Methods

Study Design

This analytical, cross-sectional, health-facility-based study was performed at referral primary health care centers (RPHCCs) selected from localities in Khartoum State.

Study population and sampling procedures

A total of 632 primary health care centers are available to provide preventive and curative health services for the population of Khartoum ^[12]. The first-stage study population was chosen from 74 RPHCCs that have a high attendance rate and provide an advanced package of services. Participants were eligible for inclusion if they were adults aged 18 years or older and were not known to have diabetes or previously diagnosed with diabetes (participants must not have received a previous diagnosis of diabetes mellitus from a healthcare provider). The exclusion criteria were pregnancy (a female individual who has confirmed pregnancy through a positive pregnancy test result), the use of metformin or other glucose-modifying medications (an individual who is currently prescribed and actively taking Metformin or another glucose-modifying drug for the management of diabetes mellitus or other metabolic conditions), and critical illness (individuals who exhibit clinical signs and symptoms indicative of severe illness, typically requiring urgent medical intervention or hospitalization).

Sample size

Three-stage random cluster sampling was adopted. The first stage was the division of Khartoum State into seven localities, and the second stage was the selection of targeted RPHCCs in these localities. The third stage was the selection of targeted attendees from each health center. The sample size was calculated as follows^[13]:

$$n = z^2 pq / d^2 * deff$$

Where:

- n = the desired sample size
- z = the confidence coefficient, 1.96
- p = 10.9% (the proportion of unknown diabetics derived from the prevalence of diabetes in Sudan according to IDF 2018)
- p = 100-10.9 = 89.1
- q = 1-p = 1-0.89 = 0.11
- d = desired margin of error, 0.05
- deff = design effect, 1.5

The sample size was calculated as 226 participants, which was divided proportionally among the 7 localities. The total number of selected health centers in all of Khartoum State was 10. The study used probability proportional to size to calculate the number of attendees in each RPHCC. Finally, the selection of the sample unit (attendees) was done using systematic random sampling during the sample collection at each selected RPHCC.

Data Collection Procedures

Data were gathered through face-to-face interviews, measurements of weight, height, and waist circumference, and blood spot samples. An adapted DRS questionnaire was used, which consisted of the 12 original questions about the main risk factors for T2DM extracted from the CANRISK tool ^[14] in addition to three added questions to reflect cultural and nutritional habits that are believed to influence the risk of Type 2 diabetes among the Sudanese population. The questionnaire was pre-tested and administered to 20 non-diabetic patients at Alferdos Primary Healthcare Centre. All investigations were tested. Out of twenty, two had high diabetes scores. The findings have been excluded from the study.

Diabetes risk scores were considered as the outcome variable. Based on the original score applied in First Nations and Métis communities in Canada, the participants were divided according to diabetes risk scores into low, moderate, and high-risk groups for those with scores of less than 21, 21 to 32, and 33 or more, respectively ^[14].

To measure the sensitivity and specificity in regard to HbA1c readings, DRS scores were also categorized using binary outcomes: scores less than 33 were considered as negative DRS, while scores of 33 or more were considered as positive DRS. The height in centimeters and weight in kilograms were used to calculate the body mass index (BMI) to assess general obesity, while the waist circumference was used for central obesity. Height, weight, BMI, and waist circumference were all considered measurable independent variables ^[15].

A random blood glucose test was performed through a capillary blood sample using a glucometer (FreeStyle Lite, Abbott Diabetes Care Inc., Alameda, CA) and pin-prick lancets. The cutoff point was set as 140 mg/dl to distinguish between high and low readings. Due to its availability at the facility-based level and reasonable cost, the test was used as a proxy indicator for blood glucose levels in the study. The most recent HbA1c test was used to determine the average blood glucose levels over the previous three months with a point-of-care device (Clover A1C- HbA1c Analyzer®).

According to the American Diabetes Association, the following HbA1c cut-off points were established ^[16]. The risk of developing DM or prediabetes when the HbA1c result was less than 5.7%, moderate risk was indicated by a result between 5.7% and 6.4%, and high risk was indicated by a result of more than 6.4%. The HbA1c test result was also used as a binary outcome as follows: HbA1c of 6.4% or more was considered positive for T2DM, and HbA1c less than 6.4% was considered negative for T2DM.

Statistical analysis

The Statistical Package for the Social Sciences (SPSS) version 25 for Windows was used for analyses. Data were coded, entered, cleaned, and documented according to the category of risk scores on the questionnaire. Descriptive analysis was performed for the dependent and independent variables of the study population using percentages, tables, and figures. The sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and ROC curves were generated using cross-tabulation of the outcome (DRS) with the reference test (HbA1c).

Ethical consideration

The Sudan Medical Specialization Board and the Major Research Committee of the Khartoum State Ministry of Health provided official ethics approval. Written consent was obtained from the participants after explaining the purposes and objectives of the research. The blood test used was noninvasive and required a minimum-risk procedure with a pinprick technique. The participants were made aware of their right to leave the study at any moment without suffering any consequences. Data confidentiality was guaranteed, and only the principal investigator had access to personal information, which was kept private. For those who were determined to have Type 2 diabetes, a referral note was given to seek medical care and advice. General lifestyle advice was delivered to those who had low risk scores.

Results

A total of 214 outpatient attendees were enrolled, resulting in a response rate of 94.7% (5.3% did not consent to participate in HbA1c testing). The participants were 22.9% male and 77.1% female. Regarding sociodemographic characteristics, 77.6% of the study population were married, and most (40.7%) were originally from the central region. The age group of 38–47 years had the highest percentage of participants (37.9%), while the age group of 58–67 years had the lowest percentage (9.3%). The respondents' occupation, household size, length of time living in Khartoum, and income are shown in Table 1.

Characteristics	n (%) (N=214)
Sex	
Female	165 (77.1%)
Male	49 (22.9%)
Age groups in year	
18-27	32 (15.0%)
28-37	48 (22.4%)
38-47	81 (37.9%)
48-57	27 (12.6%)
58-67	20 (9.3%)
<68	6 (2.8%)
Place of origin	
Central Region	87 (40.7%)
North	46 (21.5%)
South	5 (2.3%)
East	6 (2.8%)
West	70 (32.7%)
Duration of residency in Khartoum state	
Less than 15 Years	30 (14%)
15- 30 Years	111 (51.9%)
31- 45 Years	40 (18.7%)
More than 45 Years	33 (15.4%)
Marital Status	
Married	166 (77.6%)
Unmarried	48 (22.4%)
No. of family members	
Less than 3 members	35 (16.4%)
Between 3 & 5	66 (30.8%)
More than 5 members	113 (52.8%)
Occupation	
Employed	110 (51.4%)
Not employed	104 (48.6%)
Level of Education	
University and above	84 (39.3%)
Secondary	106 (49.5%)
Primary or Illiterate	24 (11.2%)
Family Income to Expenditure	
Income more than expenditure	7 (3.3%)
Income equal to expenditure	64 (29.9%)
Income less than expenditure	143 (66.8%)

Table 1. Socio-demographic characteristics among attendees of Referral Primary Health Care Centers at Khartoum State

The participants' medical histories indicated various conditions. There were 71 participants (31.2%) who had a history of high blood pressure or were taking medication for hypertension, while 41.1% (88) of the participants had a family history of diabetes, and 13.6% (29) had a history of high glucose levels at some point in time Table 2.

Characteristics of participants		Frequency	Percentage
History of high BP or medication	No	143	66.8 %
	Yes	71	31.2%
Family History of DM	No	126	58.9%
	Yes	88	41.1%
History of high glucose level ever before	No	185	86.4%
	Yes	29	13.6%

Table 2. Medical history among attendees of Referral Primary Health Care Centers at Khartoum State, Sudan, n=214

Regarding BMI, 34.6% ^[74] were within the normal BMI range, while 29.4% were overweight, and 36% were obese. Furthermore, 27.3% of women had a normal waist circumference of less than 80 cm, 26.1% had a circumference between 80 and 88 cm, and 46.7% had a circumference of 88 cm or more. Among men ^[27], 55% had a regular waist circumference (less than 94 cm), while 24.6% had a waist circumference between 94 and 102 cm, and approximately 20.4% ^[12] had a waist circumference of 102 cm or higher. Only 9% of the study population were smokers, and 76% were physically inactive Table 3.

Characteristics of participants	n (%)
BMI N=214	
Normal Weight (18-Less than 25)	74 (34.6%)
Overweight (25-Less than 30)	63 (29.4%)
Obesity (30 and above)	77 (36.0%)
Waist circumference <u>for Women</u> N=165	
Less than 80 cm	45 (27.3%)
80-88 cm	43 (26.1%)
88 cm and more	77 (46.7%)
Waist circumferenc <u>for Men</u> N=49	
Less than 94cm	27 (55.1%)
94-102 cm	12 (24.5%)
102 cm and more	10 (20.4%)
Vegetables consumption N=214	
More than two times per week	198 (92.5 %)
Less than twice per week	16 (7.5 %)
Fruits consumption N=214	
More than two times per week	73 (34.1 %)
Less than twice per week	141 (65.9 %)
Kisra-Asida-Gurrasa -Bread N=214	
More than two times per week	161 (73.8 %)
Less than twice per week	53 (26.2%)
Smoking N=214	
YES	19 (9%)
NO	195 (91%)
Physical Activity N=214	
Yes	52 (24%)
No	162 (76%)

Table 3. Anthropometric Measurements, eating habits, smoking, and physical activity among attendees of Referral Primary Health Care Centers at Khartoum State at Khartoum State.

Concerning RBG, the majority of the participants (93%) had normal values (cutoff: < 140 mg/dl). Eighty-six percent were negative for diabetes according to the HbA1c result (cutoff: < 6.4%). Regarding DRS, 59.8% were considered negative (cutoff: < 33) Table 4.

Categorical and binary outcome variables with their corresponding cutoff values among attendees of Referral Primary Health Care Centers at Khartoum State, Sudan .						
Finding	Categorical outcome	Frequency	Percent	Binary outcome	Frequency	Percent
Random blood glucose	≤140g/dl	199	93.0 %	Negative (≤140g/dl)	199	93.0 %
	>140g/dl	15	7.0 %	Positive (>140g/dl)	15	7.0 %
HbA1c	<5.7	166	77.6 %	Negative (<6.4%)	184	86 %
	5.8 – 6.4	18	8.4 %			
	≥6.4	30	14.0 %	Positive (≥6.4%)	30	14.0 %
DRS	less than 21	70	32.7 %	Negative (<33)	128	59.8 %
	21 – 33	58	27.1 %			
	33 and more	86	40.2 %	Positive (≥33)	86	40.2 %

Table 4. Binary outcome variables with their corresponding cutoff values among attendees of Referral Primary Health Care Centers at Khartoum State.

[†] e Abbreviations RBG, Random blood glucose, HbA1c, hemoglobin A1c, DRS, Diabetes Risk Score

[‡] N = 214

HbA1c tests were used as a reference test in this study. Thirty participants were considered diabetic by both tests (14.0%) Table 5.

Diabetes Risk Score	HbA1c		Total
	Positive (≥6.4%)	Negative (<6.4%)	
High Score(33 and more)	25	61	86
Low score (less than 33)	5	123	128
Total	30	184	214

Table 5. Cross-tabulation for DRS with HbA1c (Sensitivity Analysis) among attendees of Referral Primary Health Care Centers at Khartoum State.

[†] Percentage according to the number of patients

[‡] N = 214

[§] Sensitivity = $25/30 \times 100 = 83.33\%$

[¶] Specificity = $123/184 \times 100 = 66.85\%$

The sensitivity and specificity of the DRS in relation to the reference test were 83.33% and 66.8%, respectively. The total PPV was 29.07%, and the NPV was 96.09%. The area under

the curve (AUC) for the DRS was 0.751 (95% confidence interval (CI): 0.662–0.840) (Figure 1).




Figure 1. It shows the Area under the Curve for the Diabetes Risk Score (sensitivity against 1-specificity) among attendees of the Referral Primary Health Care Centers at Khartoum State, Sudan. The AUC for the Diabetes Risk Score = 0.751 (95% CI: 0.662–0.840)

Discussion

This study evaluated the DRS as a screening tool by predicting the prevalence of undiagnosed diabetes, which was measured at 14%. This finding is consistent with the prevalence of diabetes in the MENA region (9.2%) [2]. The result is supported by IDF 2019 statistics for Sudan [4], as well as the Khartoum State STEPS 2016 survey (11.6% with a CI of 9.1–14.1%) [5]. The prevalence of newly diagnosed participants was nearly identical to that of a recent study conducted in Gadarif State in 2019, which revealed a prevalence of 10.0%.

Men had a significantly higher risk than women. In comparison, according to the findings of a Saudi study, women had higher scores than men in both moderate and high-risk categories [17], although none of the models from Gulf regions addressed gender [18]. The DRS has been widely implemented as a low-cost and valid screening tool in many countries to detect those who are at risk of developing T2DM. This risk prediction model enables early detection, prevention, and intervention [14].

The majority of participants (72.4%) were 18 to 44 years old, which is similar to the CANRISK study performed with a South Asian population. Regardless of the model, the odds of dysglycemia increased with age, and there were significantly higher odds in the older age groups. Both results demonstrate a significant relationship between age and the risk of diabetes [14]. Regarding BMI, the majority of the participants were overweight or obese, with rates of 29.4% and 36%, respectively. According to the DRS, 40.2% had a high-risk DRS (≥ 33), while 59.8% had a low-risk DRS (< 33). This risk score is widely

recommended for use in low-resource settings as one of the major approaches for screening programs ^[14].

The sensitivity and specificity of our DRS were 83.33% and 66.85%, respectively. In the Eastern Mediterranean region, many similarities related to the DRS have been reported. Studies performed in Saudi Arabia, Kuwait, United Arab Emirates, and Oman reported sensitivities and specificities of 76.6% and 52.1%, respectively ^{[18][19][20][21]}. In 2015, there were approximately 1.5 million new diabetes cases among adults according to the 2017 National Diabetes Statistics Report of the Centers for Disease Control. Adults aged 45 to 64 years were the most prevalent age group for diabetes diagnoses ^[22].

The AUC result shows that 95% of DRSs were acceptable and had good predictability for preventing undiagnosed diabetes. Thus, the DRS could be used at the community level. This finding is similar to that of a Canadian study, in which the AUC was 0.80 for South Asians with a slight reduction to 0.75 among First Nations/Métis populations ^[14]. The accuracy of the DRS in this study is consistent with the previous observations of the CANRISK and FINDRISC surveys, in which the DRS performed reasonably well in identifying patients with elevated blood glucose levels, with AUC curves ranging from 0.69 to 0.85% ^[23]. The PPV and NPV results are similar to the results of a Métis population, in which the PPV was 30%, and the NPV was 90% with a cutoff point of 33. In the Canadian South Asian population study, the PPV was 28%, and the NPV was 93% ^[14].

A considerable number of people in Khartoum City were at risk of developing T2DM. The questionnaire used is reliable, valuable, and easy to use as a screening tool. The prevalence of diabetes among undiagnosed participants was considerable. Less than half of the participants had high DRS results.

The sensitivity, specificity, and AUC of the DRS tool showed that it is an accurate method that is suitable for application in the screening of diabetes in the health system in Sudan. The main recommendations of this study are to adopt the DRS tool as an easy, affordable, and accessible tool for diabetes screening in populations at the primary healthcare level and to adopt further confirmation by blood tests for DRS in moderate and high-risk populations to reduce the economic burden on the health system. This study comprises a representative sample of primary health care centers from Khartoum State's major geographical regions. In part due to the limitations of the cross-sectional investigation, we were unable to assess DRS' effectiveness in predicting T2DM events. Therefore, conduct a follow-up investigation to assess the predictive validity of the DRS. Another drawback is that the population demographics were tilted toward females, which may have significantly undermined the risk score's performance in the male gender. Also, more research is needed to examine a larger Sudanese population to test variables related to Sudanese culture and the risks of developing diabetes, which may limit the applications of the DRS on a larger population scale. Furthermore, the variable employment can be adjusted to better suit certain categories while also providing generic information. The research work's future scope and motivation can be used to predict the necessity of it.

Appendix

Diabetes Risk Score Questionnaire for undiagnosed Diabetic client attending Referral Primary Health Care Centers at Khartoum State			
Do you have Diabetes: 1- Yes 2- No			
Serial Number:			
Health Center:			
Locality:			
Target population: Adult 18 and above, Not Known Diabetic, who is attending Family Medicine Or General Clinics at Referral Primary Health Care Centre (RPHCC) at Khartoum State.			
No	Questions	Answers	Scores for individual questions
Answer the questions and Choose the corresponding score for each question in Pink, then Add the total scores at the end			
1	Place of origin	0-Central Region 1- North 2-South 3-East 4- West	Origin of participants mother and father: If the origin South Asia.....11 If the origin Africa.....5 If the origin of Arab, Turk.....3 If the origin of white0
2	Duration of residency in Khartoum in years		
3	Occupation	0-Employed 1-Not Employed	If yes, Specify:
4	Marital status	0-Married 1- Not married	
5	Education	0- University and above 1- Secondary or Primary 2- Illiterate	00 1 2 2 6
6	Age Years	Less than 44.....0 45-54.....6 55-64.....14 65 and more.....15
7	Sex	0- Female 1- Male	00 16
8	Head of the Family financially	
9	No of Family member	
10	Family income .to. Expenditure	Less /Equal/ More	
11	Currently smoking (Cigarette/Tumbak /Shishah)	0-No..... If No, ever smoke: 0- Yes 1- No	00 11
12	Do you usually do Physical Activity 30 minutes/day	0- Yes If Yes, what kind:..... 1- No	00 11
13	Usually how many days/week you eat Vegetables	0-More than two times per week. 1- Less than twice per week	00 11
14	Usually how many days/week you eat fruits	0-More than two per week..... 1- Less than twice per week	00 11
15	Usually how many days/week you eat: Kiswa-Asida / Gurrasa / Bread Baladi with Raddah / Bread Afrangi / Rice	0-More than two times per week 1-Less than twice per week	
16	Family History of DM (Mother, Father, Sisters, Brothers & Kids)	0- No 1- Yes	00 16
17	History of high glucose level	0- No 1- Yes	00 114
18	History of high BP or medication	0- No 1- Yes	00 14
19	Ladies: History of large baby 4.1 kg	0- No 1- Yes	00 11
20	BMI	Height Cm Weight..... Kg	Less than 25.....0 25-29.....6 30 and more14
21	Waist Circumference Cm	WOMEN Less than 80.....0 80-88.....4 88 and more.....6 MEN Less than 94.....0 94-102.....4 102 and more.....6	
Total Score for question (1,5,6,7,11,12,13,14,16,17,18,19,20,21)			21 and less Low Risk 21-33 Moderate 33and more High Risk
Test	HbA1c%	5.7and less Normal 5.7-6.4 Pre-diabetes 6.4 & more Diabetes

No.	Locality	N (total population)	Selected centers /locality	N (desired population)	Selected centers
1	Khartoum	32150	2	57	Al ferdos Sameer
2	Jebel Aulia	18656	2	61	Al Ingath 1 Kalakla Guba
3	Omdurman	4592	1	15	Awad Hussain
4	Karrari	8192	1	20	Al Harah no 38
5	Khartoum north	17778	2	44	Al Ingath Khartoum north
6	Ombada	8408	1	21	Al Harah no 52
7	East Nile	9964	1	25	Martyr Nada Health Center

Table A1. The distribution of the sample size was proportional to the 7 localities.

Abbreviations

- AUC: Area under the Curve.
- BMI: Body Mass Index.
- CANRISK: Canadian Diabetes Risk Assessment Questionnaire.
- CBS: Community Based Screening.
- CDC: Centers for Disease Control and Prevention.
- CHAD: Community Health Awareness of Diabetes.
- DRS: Diabetes Risk Score.
- FBG: Fasting Blood Glucose.
- FINDRISC: Finnish Diabetes Risk Score.
- GLV: Green Leafy Vegetables.
- HbA1c: Glycosylated Hemoglobin A1C.
- HTN: Hypertension.
- IDF: International Diabetes Federation.
- MENA: Middle East and North Africa.
- NCDs: Non-Communicable Diseases NPV Negative Predictive Value.
- PPV: Positive Predictive Value OGTT Oral Glucose Tolerance Test.
- PPS: Probability Proportional Sampling.
- RAQ: Risk Assessment Questionnaires.
- RBG: Random Blood Glucose.
- ROC: Receiver Operating Characteristic.
- RPHC: Referral Primary Health Care Centers.
- SPSS: Statistical Package for Social Sciences.
- STEPS: STEPwise approach to Surveillance.
- T2DM: Type 2 Diabetes Mellitus.
- WC: Waist Circumference.
- WHO: World Health Organization.
- WHO/PEN: World Health Organization / Package of Essential Non-communicable Disease Interventions.

Notes

Third, due to the nature of the cross-sectional study, we were unable to evaluate the performance of DRS in predicting DM and preDM events. We will try to conduct a follow-

up study and assess the predictive validity of NCDRS further.

Validation of the risk score in the same population and the use of a larger sample size would have further enhanced the generalizability of our results.

Statements and Declarations

Ethics

The Evaluation of Diabetes Risk Score Tool for Detecting Undiagnosed Type 2 Diabetes Mellitus in Attendees of Referral Clinics at Primary Health Care Centers, Sudan study was approved by the Sudan Medical Specialization Board, and the Khartoum State Ministry of Health's Major Research Committee provided official ethics approval. The researcher obtained written consent from the participants after explaining the research purposes and objectives.

Data Availability

The raw data supporting the conclusions of this article may be made available by the authors upon reasonable request, pending ethical considerations regarding participant privacy. Inquiries can be directed to the corresponding author.

Author Contributions

- SF: Principal investigator was actively involved in the planning, conducting, data collection, and design of the study, also in writing the paper.
- AA: Was involved in the data analysis.
- ME: Was involved in data collection, interpretation of the data, as well as writing the paper. Author Agreement
- OK: Actively involved in the planning, conception, and design of the study, also revising the paper. All authors have approved the final manuscript.

References

1. [△]Roglic G (2016). "WHO Global report on diabetes: A summary." *Int J Non-Commun Dis.* 1:3–8.
2. [△][△]Cho NH, Shaw JE, Karuranga S, Huang Y, da Rocha Fernandes JD, Ohlrogge AW, Malanda B (2018). "IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045." *Diabetes Res Clin Pract.* 138:271–281.
3. [△]United Nations (2002). *World population ageing, 1950–2050*. New York: United Nations.
4. [△][△]IDF Middle East & North Africa (2019). "Sudan 2019." IDF Middle East & North Africa. <https://idf.org/our-network/regions-members/middle-east-and-north-africa/members/47-sudan.html>.
5. [△][△]Federal Ministry of Health, Sudan (2016). "Sudan STEP-wise survey for non-communicable diseases risk factors, 2016 report." Federal Ministry of Health, Sudan. <https://extranet.who.int/ncdsmicrodata/index.php/catalog/438>.
6. [△]Omar SM, Musa IR, ElSouli A, Adam I (2019). "Prevalence, risk factors, and glycaemic control of type 2 diabetes mellitus in eastern Sudan: a community-based study." *Ther Adv Endocrinol Metab.* 10:2042018819860071.
7. [△]Bos M, Agyemang C (2013). "Prevalence and complications of diabetes mellitus in Northern Africa, a systematic review." *BMC Public Health.* 13:387.
8. [△]Noor SK, Bushara SO, Sulaiman AA, Elmadhoun WM, Ahmed MH (2015). "Undiagnosed diabetes mellitus in rural communities in Sudan: prevalence and risk factors." *East Mediterr Health J.* 21(3):164–70.
9. [△]Mendis S, Al Bashir I, Dissanayake L, Varghese C, Fadhil I, Marhe E, Sambo B, Mehta F, Elsayad H, Sow I, Algae M, Tennakoon H, Truong LD, Lan le TT, Huiuinato D, Hewageegana N, Fahal NA, Mebrhatu G, Tshering G, Chestnov O (2012). "Gaps in capacity in primary care in lo

- w-resource settings for implementation of essential noncommunicable disease interventions." *Int J Hypertens*. 2012;584041.
10. [△]World Health Organization (2017). Tackling NCDs: best buys' and other recommended interventions for the prevention and control of noncommunicable diseases. Geneva (CH): World Health Organization.
 11. ^{△, b}World Health Organization (2003). Screening for type 2 diabetes: report of a World Health Organization and International Diabetes Federation meeting. World Health Organization.
 12. ^{△, b}Central Bureau of Statistics (Sudan) (2018). Central Bureau of Statistics (Sudan). <http://ghdx.healthdata.org/organizations/central-bureau-statistics-sudan>.
 13. [△]Charan J, Biswas T (2013). "How to calculate sample size for different study designs in medical research?" *Indian J Psychol Med*. 35(2):121–6. doi:10.4103/0253-7176.116232. PMID 24049221.
 14. ^{△, b, c, d, e, f}Agarwal G, Jiang Y, Rogers Van Katwyk S, Lemieux C, Orpana H, Mao Y, Hanley B, Davis K, Leuschen L, Morrison H (2018). "Effectiveness of the CANRISK tool in the identification of dysglycemia in First Nations and Métis in Canada." *Health Promot Chronic Dis Prev Can*. 38(2):55–63.
 15. [△]National Institutes of Health (1998). "Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults--The Evidence Report." *Obes Res*. 6 Suppl 2:51S–209S. PMID 9813653.
 16. [△]American Diabetes Association (2011). "Diagnosis and classification of diabetes mellitus." *Diabetes Care*. 34 Suppl 1(Suppl 1):S62–9. doi:10.2337/dc11-S062. PMID 21193628.
 17. [△]Alghwiri AA, Alghadir A, Awad H, Anwer S (2018). "Comparison of diabetes risk estimate in the cities of Riyadh and Amman." *Medicine (Baltimore)*. 97(40):e12689. doi:10.1097/MD.00000000000012689.
 18. ^{△, b}Ng SW, Zaghloul S, Ali HI, Harrison G, Popkin BM (2011). "The prevalence and trends of overweight, obesity and nutrition-related non-communicable diseases in the Arabian Gulf States." *Obes Rev*. 12(1):1–13. doi:10.1111/j.1467-789X.2010.00750.x. PMID 20546144.
 19. [△]Sulaiman N, Mahmoud I, Hussein A, Elbadawi S, Abusnana S, Zimmet P, Shaw J (2018). "Diabetes risk score in the United Arab Emirates: a screening tool for the early detection of type 2 diabetes mellitus." *BMJ Open Diabetes Res Care*. 6(1):e000489.
 20. [△]Al Khalaf MM, Eid MM, Najjar HA, Alhajry KM, Doi SA, Thalib L (2010). "Screening for diabetes in Kuwait and evaluation of risk scores." *East Mediterr Health J*. 16(7):725–31.
 21. [△]Al-Lawati JA, Tuomilehto J (2007). "Diabetes risk score in Oman: a tool to identify prevalent type 2 diabetes among Arabs of the Middle East." *Diabetes Res Clin Pract*. 77(3):438–44. doi:10.1016/j.diabres.2007.01.013. PMID 17306410.
 22. [△]McGuire S (2014). "Centers for Disease Control and Prevention. State indicator report on Physical Activity, 2014. Atlanta, GA: U.S. Department of Health and Human Services; 2014." *Adv Nutr*. 5(6):762–3.
 23. [△]Robinson CA, Agarwal G, Nerenberg K (2011). "Validating the CANRISK prognostic model for assessing diabetes risk in Canada's multi-ethnic population." *Chronic Dis Inj Can*. 32(1):19–31. PMID 22153173.

Declarations

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.