

Is creeping abandon of human cancer defences evolutionarily favoured?

Alexander D. Rahm¹, Pierre Pratley²

¹ Université de la Polynésie Française

² Royal Tropical Institute

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.

Abstract

Among the animal species on which observations are available, humans have a uniquely high lifetime risk to suffer from cancer - over 38%, compared to less than 10% for all observed other species (except species suffering from environmental pollution). Peto's paradox shows that this cannot simply be explained by mathematical models which view cancer genesis as a stochastic process, with resulting risks polynomial in lifespan and body mass - whales have a longer lifespan and about 30 times the human body mass, however their cancer risk remains constant throughout their life rather than increasing sharply after female reproductive age as observed in humans. Rather, it is well documented in the literature that species-specific tumour suppression mechanisms allow for large lifespan and body mass. Chimpanzees, being closely related to humans, have a very low cancer risk, and hence the weakness of human cancer defence is likely to have resulted from the specific development of homo sapiens. As this weakness appears past the reproductive years, a prominent hypothesis blames it to antagonistic pleiotropy. However, homo sapiens having lived in small tribes during most of its development, natural selection is likely to also have acted at the level of tribes, and higher degrees of inbreeding would quite certainly have been detrimental to a tribe. And males of high social status can attract new reproductive partners again and again until an age that has seen several generations grow, which in case of a not-so-large tribe would have considerably narrowed down its genetic pool. Furthermore, lowering tumour suppression activities might save calories and hence benefit tribes with limited food production; and individuals suffering from cancer after female reproductive age could still have made contributions to parental/grandparental care, while no more being attractive as a reproductive partner. So, we arrive at the mentioned hypothesis: Is creeping abandon of human cancer defences evolutionarily favoured?

Alexander D. Rahm¹ and Pierre Pratley²

¹ *Laboratoire GAATI, University of French Polynesia, BP 6570 - 98702 Faa'a, French Polynesia*

² *KIT Royal Tropical Institute, Global Health Department, Mauritskade 64, 1092 AD Amsterdam, The Netherlands*

1. Context

Among the animal species on which observations are available, humans have a uniquely high lifetime risk to suffer from cancer - over 38%, compared to less than 10% for all observed other species ^{[1][2][3][4]} (except species suffering from environmental pollution ^[5]). Obviously many cases of human cancer are due to carcinogens which we eat, breathe and receive in other ways through our modern lifestyle, which were not available to prehistoric humans, nor are they to observed animals. However, and though with some uncertainty, the literature lets us suspect ^[6] that these modern carcinogens do not explain all of the high modern human cancer mortality, so a significant share of it may be due to differences in species-specific cancer defenses. Peto's paradox ^[7] shows that this cannot simply be explained by mathematical models which view cancer genesis as a stochastic process, with resulting risks polynomial in lifespan and body mass - whales have a longer lifespan and about 30 times the human body mass, however their cancer risk remains constant throughout their life rather than increasing sharply after female reproductive age as observed in humans ^[8]. Rather, it is well documented in the literature that species-specific tumour suppression mechanisms allow for large lifespan and body mass ^{[9][10]}. Chimpanzees, as the extant species most closely related to humans (and with particularly matching cancer genes ^[11]), have a very low cancer risk ^[6], and hence the weakness of human cancer defence (which is supported by reduced apoptotic function compared to chimpanzee and macaque cells ^[12]) is likely to have resulted from the specific development of *homo sapiens* (see also a study on oncogene development since the chimpanzee/human last common ancestor ^[13]). As this weakness appears past the reproductive years, a prominent hypothesis blames it to *antagonistic pleiotropy* ^{[14][10]} (with as a consequence the development of menopause to protect descendants ^[15]). We want to argue in favour of a different hypothesis here, the one stated in the title of this paper.

2. Setting of the hypothesis

We base our hypothesis on the assumption that *homo sapiens* have lived in small hunter-gatherer tribes during most of the species development. In hunter-gatherer tribes of the 20th century, on which anthropological literature is available, men did hunt individually, and there were no social differences between them ^[16]. However those modern tribes were marginalized into territories unsuitable for agriculture and pastoralism, and little animals were available for them to hunt, while cave paintings tell us that in contrast, early humans did attack mega-fauna like mammoths, bears, etc., which they had to do in gangs. So we suppose that each tribe was growing around a hunting gang, and as a consequence, males born into the tribe would become members of the hunting gang, and women would have had the opportunity to move to a different tribe once reaching adulthood. This immediately breaks the egalitarian social structure and gives a claim for the boss of the hunting gang to become also chieftain of the tribe, and we have some evidence for such tribal structures from the historic fact that patriarchal societies were much more frequent than matriarchal ones. Then natural selection is likely to have acted at the level of tribes, because the paternal genes of the tribe all come from the same kin, and the survival of the kin's genes is linked to the survival of the tribe. Then higher degrees of inbreeding would quite certainly have been detrimental to a tribe. And males of high social status can attract new reproductive partners again and again until an age that has seen several generations grow ^[17], which in case of a not-so-large tribe would have considerably narrowed down

its genetic pool. Of course, as skeleton findings suggest that highly aged individuals were rather rare among prehistoric humans (among Early and Middle Pleistocene Homo, 42 old adults and 166 young adults have been found ^[18]), one could just dismiss the influence of such individuals on the genetics of a tribe. But extended lifespans are possible for hunter-gatherers according to observations in modern tribes ^[19], and lowering tumour suppression activities as part of a state of ageing would actually have reduced the chances of individuals to reach a high age in the first place, so to contribute explanation to the age distribution of prehistoric skeleta (mortality due to infectious diseases should have decreased in adult age when all regionally circulating germs were known to the immune system, and mortality due to predation should have decreased as well for adults experienced with predating animals). Furthermore, as tumour suppression involves killing a lot of suspect cells, lowering tumour suppression activities might save a decisive amount of calories (so to account for sometimes observed lower appetite of aged persons) and hence benefit tribes with unreliable food collection; and individuals suffering from cancer after female reproductive age could still have made contributions to parental/grandparental care, while no more being attractive as a reproductive partner. Now if epigenetic deactivation of tumour suppressor genes as part of the telomere-triggered senescence process is coded on a dominant gene, then individuals within a tribe would not have been able to betray a tribe-wide lowering of tumour suppression activities, because the men of a tribe do in our setting all carry that gene, and it would remain active when combined with the genes of women entering the tribe from outside.

3. The hypothesis

Therefore, we think that research on the question

Is creeping abandon of human cancer defences evolutionarily favoured?

could facilitate cancer prevention. In a study which modeled cancer defence activities as species- and life-history-dependent but constant over age ^[20], it was already suggested that a model with age-dependent cancer defence activities could provide a better explanation for Peto's paradox; and in view of the above argumentation, we would expect human cancer defences to reduce their activity with progressing age.

A positive answer to our question would then of course open the question why creeping abandon of cancer defences cannot be observed in chimpanzees, which also live in patriarchal groups --- does their pattern have to do with not enough chimps living into old age, or do they have a more favorable diet free of modern day carcinogens, or does the fact that reproductive value increases or remains high into older age drive greater anti-cancer adaptations?

4. Suggested experiments

1. The hypothesis could be tested directly by analyzing cell kernels of present-day aged persons, and checking if tumour suppressor genes have been deactivated epigenetically.
2. Among the DNA found in prehistoric human remains, oncogenes and tumour suppressor genes could be tracked. This would potentially give some indications on whether the former increased and the latter decreased over a long

timespan. So far, mainly agricultural societies of just a few hundred generations ago have been studied from an evolutionary perspective for genetic markers of cancer defense [21], but DNA can now be collected and analyzed from much older human remains [22].

3. The life cycle events in a prehistoric tribe could be modeled stochastically, along with resources (particularly food) accessible to the tribe and influencing its prosperity, keeping track of the health of individuals throughout the modeled years, and a sample of their genes to track the effects of incest over generations (via detrimental recessive genes); then epigenetic mechanisms which decrease the cancer defences after female reproductive age could be introduced in a variant of this Monte Carlo simulation. This could potentially make it more plausible that creeping abandon of cancer defences increases the evolutionary competitiveness of a tribe. The authors can implement such an *in silico* model themselves, but will wait for eventual comments (maybe from people interested in collaborating on *in silico* experiments on this) before doing so. It is obvious that such a simulation will involve so many parameters that they cannot just be varied in order to find plausible values for them by conclusions from the output they produce, but instead the tribes need to be modeled using anthropological literature about 20th century hunter-gatherer tribes, in order to make informed guesses.

Other References

- Wolf A.M. (2010), The tumor suppression theory of aging, *Mechanisms of Ageing and Development* 200: 111583.

References

1. ^ Chu P-Y., Zhuo Y-X., Wang F-I., Jeng C-R., Pang V.F., Chang P-H., Chin S-C., Liu C-H. (2012), *Spontaneous neoplasms in zoo mammals, birds, and reptiles in Taiwan – A 10-year survey. Animal Biology* 62: 95-110.
2. ^ Takayama S., Thorgeirsson U.P., Adamson R.H. (2008), *Chemical carcinogenesis studies in nonhuman primates. Proc. Jpn. Acad., Ser. B* 84: 176-188.
3. ^ Wood C.E., Osborne A.L., Starost M.F., Tarara R.P., Hill L.R., Wilkinson L.M., Geisinger K.R., Feiste E.A., Cline J.M. (2006), *Hyperplastic and neoplastic lesions of the mammary gland in macaques. Vet. Pathol.* 43: 471-483.
4. ^ Xia H-J., Chen C-S. (2011), *Progress of non-human primate animal models of cancers. Zoological Research* 32(1): 70-80.
5. ^ Martineau D., Lemberger K., Dallaire A., Labelle P., Lipscomb T.P., Michel P., Mikaelian I. (2002), *Cancer in wildlife, a case study: beluga from the St. Lawrence estuary, Québec, Canada. Environmental Health Perspectives* 110(3): 285-292.
6. ^{a, b} Varki N.M., Varki A. (2015), *On the apparent rarity of epithelial cancers in captive chimpanzees. Philos Trans R Soc Lond B Biol Sci.* 370(1673): 20140225.
7. ^ Peto, R. (2015), *Quantitative implications of the approximate irrelevance of mammalian body size and lifespan to lifelong cancer risk. Phil. Trans. R. Soc. B* 370: 20150198.

8. [^]Nunney L., Maley CC., Breen M., Hochberg ME., Schiffman JD. (2015), *Peto's paradox and the promise of comparative oncology*. *Phil. Trans. R. Soc. B* 370: 20140177.
9. [^]Albuquerque T.A.F., Drummond do Val L., Doherty A., de Magalhães J.P. (2018), *From humans to hydra: patterns of cancer across the tree of life*. *Biol. Rev.* 93: 1715-1734.
10. ^{a, b}Laconi E., Marongiu F., DeGregori J. (2020), *Cancer as a disease of old age: changing mutational and microenvironmental landscapes*. *Br J Cancer* 122: 943-952.
11. [^]Puente X-S., Velasco G., Gutiérrez-Fernández A., Bertranpetit J., King M-C., López-Otín C. (2006), *Comparative analysis of cancer genes in the human and chimpanzee genomes*. *BMC Genomics* 7:15.
12. [^]Arora G., Mezencev R., McDonald J. (2012), *Human Cells Display Reduced Apoptotic Function Relative to Chimpanzee Cells*. *PLOS ONE* 7(9): e46182.
13. [^]Huang J., Zhong Y., Makohon-Moore A., White T., Jasin M., Norell M., Wheeler W., Iacobuzio-Donahue C. (2022), *Evidence for reduced BRCA2 functional activity in Homo sapiens after divergence from the chimpanzee-human last common ancestor*, *Cell Reports* 39(5): 110771. <https://doi.org/10.1016/j.celrep.2022.110771>
14. [^]Hamilton W.D. (1966), *The moulding of senescence by natural selection*. *J. Theor. Biol.* 12: 12-45.
15. [^]Thomas F., Giraudeau M., Renaud F., Ujvari B., Roche B., Pujol P., et al. (2019), *Can postfertile life stages evolve as an anticancer mechanism?* *PLoS Biol.* 17(12): e3000565.
16. [^]Ember C.R. (2020), *Hunter-Gatherers*. C. R. Ember, ed. *Explaining Human Culture*. *Human Relations Area Files*, <https://hraf.yale.edu/ehc/summaries/hunter-gatherers>
17. [^]The Telegraph (2009), *Silvio Berlusconi's women – the top 10*, *The Telegraph picture galleries*: 5406658.
18. [^]Caspari R., Lee S-H. (2004), *Older age becomes common late in human evolution*. *Proceedings of the National Academy of Sciences* 101(30): 10895-10900.
19. [^]Kaplan H., Hill K., Lancaster J., Hurtado A.M. (2000), *A theory of human life history evolution: Diet, intelligence, and longevity*. *Evol. Anthropol.*, 9: 156-185.
20. [^]Brown J.S., Cunningham J.J., Gatenby R.A. (2015), *The multiple facets of Peto's paradox: a life-history model for the evolution of cancer suppression*. *Phil. Trans. R. Soc. B* 370: 20140221.
21. [^]David A.R., Zimmerman M.R. (2010), *Cancer: an old disease, a new disease or something in between?*. *Nature Reviews Cancer*, 10(10): 728-733.
22. [^]Hummel S. (2003), *Ancient DNA Typing: Methods, Strategies and Applications*. Springer, ISBN 9783540430377.