

Open Peer Review on Qeios

Stage IIA Lung Cancer AJCC v7

National Cancer Institute

Source

National Cancer Institute. Stage IIA Lung Cancer AICC v7. NCI Thesaurus. Code C5644.

Stage IIA includes: (T2b, N0, M0); (T1a, N1, M0); (T1b, N1, M0); (T2a, N1, M0). T2b: Lung cancer with a tumor size more than 5 cm but 7 cm or less in greatest dimension. T1a: Lung cancer with a tumor size of 2 cm or less in greatest dimension, surrounded by lung or visceral pleura and without bronchoscopic evidence of invasion more proximal than the lobar bronchus (i.e., not in the main bronchus). The uncommon superficial tumor of any size with its invasive component limited to the bronchial wall, which may extend proximal to the main bronchus, is also classified as T1a. T1b: Lung cancer with a tumor size more than 2 cm but 3 cm or less in greatest dimension, surrounded by lung or visceral pleura and without bronchoscopic evidence of invasion more proximal than the lobar bronchus (i.e., not in the main bronchus). T2a: Lung cancer with a tumor size more than 3 cm but 5 cm or less in greatest dimension. N0: No regional lymph node metastasis. N1: Lung cancer with metastasis in ipsilateral peribronchial and/or ipsilateral hilar lymph nodes and intrapulmonary lymph nodes, including involvement by direct extension. M0: No distant metastasis. (AJCC 7th ed.)

Qeios ID: TPYH93 · https://doi.org/10.32388/TPYH93