

Wood decomposition (Diptera: Axymyiidae).

Carlos Henrique Marchiori¹

¹ Instituto Federal Goiano

Potential competing interests: No potential competing interests to declare.

Co-authors: Marco Vinícios de Oliveira Santana² and Klebert de Paula Malheiros³.

²⁻³Instituto Marco Santana, Goiânia, Goiás, Brazil.

Family Axymyiidae is part of Order: Diptera, Suborder Nematocera, and Infraorder: Axymyiomorpha. Are medium-sized mosquitoes, robustly built, discreetly dark to reddish-brown, with a body length of 9 to 12 millimeters, and with relatively short legs and antennae. The large head has very large and complex eyes. In males these collide at the top of the head, in females they are widely separated. There are also three ocelli. The short, pearl-like antennae with constrictions at the segment boundaries consist of 16 [1-4].

The trunk bears three short pairs of legs, the tibia without spurs, spines, or longer hairs. The wings are elongated and generally somewhat darkened. The wing vein is characteristic: the ray ends in four branches, where R2 branches from its trunk at almost right angles and reaches the edge vein near the confluence of R1. The MRI of the transverse artery is transparent and appears as a continuation of the radial artery (Figure 1) [2-5].



Figure 1. *Axymyia* McAtee, 1921. Source: Sample ID: CCDB-23503-B08. Attribution (2015).

The larvae family Axymyiidae live in dead wood that is free of bark and moss, mainly in forest swamps and on the banks of small forest streams. Each larva sits in a bottle-shaped auto-rotated cavity in the wood so that only the tip of the breathing tube protrudes out. It feeds on the wood mass eaten and probably on the microorganisms it contains. They develop in four larval stages. The mature larvae rotate in the cavity and widen the tunnel to the wooden surface, in whose mouth they transform into pupa. Adult mosquitoes hatch in North America in April or early May. They are rarely found and almost nothing is known about their way of life. Some species have been observed forming flocks, probably to mate (Figure 2) [2-7].



Figure 2. *Axymyia furcata* McAtee, 1921. Source: Photo#1215221, Danby, Tompkins County, New York, USA April 23, 2016.

The small family is spread across the Palearctic. It occurs in northern North America, Siberia, and northern Asia, from southern to southern China and Taiwan, and from western to eastern Europe [2-14].

The family Axymyiidae, of which only 6 extant species in 3 genera are known, and 3 fossil species. Genera and species: *Axymyia* McAtee 1921, *Axymyia furcata* McAtee 1921 and *Axymyia japonica* Ishida 1953. *Mesaxymyia* Mamaev, 1968. *Mkerteszi* (Doubt, 1930): *Mkerteszi stackelbergi* Mamaev 1968. *Protaxymyia* Mamaev & Krivosheina 1966: *Protaxymyia melanoptera* Mamaev & Krivosheina 1966 and *Protaxymyia sinica* Yang, 1993 [15-16].

The family Axymyiidae, of which only 6 extant species in 3 genera are known, and 3 fossil species. Genera and species: *Axymyia* McAtee 1921, *Axymyia furcata* McAtee 1921 and *Axymyia japonica* Ishida 1953. *Mesaxymyia* Mamaev, 1968. *Mkerteszi* (Doubt, 1930): *Mkerteszi stackelbergi* Mamaev 1968. *Protaxymyia* Mamaev & Krivosheina 1966: *Protaxymyiamelanoptera* Mamaev & Krivosheina 1966 and *Protaxymyia sinica* Yang, 1993 (Figure 3) [15-16].



Figure 3. Several long and intersecting larval galleries. Damage: Larvae rotate in the cavity and widen the tunnel to the wooden surface. <http://doi.org/10.5281/zenodo.231670>.

References

- [1] Amorim DS. A phylogenetic analysis of the basal groups of Bibionomorpha, with a critical reanalysis of the wing vein homology. *Revista Brasileira de Biologia*. 1993; 52(3): 379-399.
- [2] Schneeberg K, Krause K, Beutel R. The adult head of *Axymyia furcata* (Insecta: Diptera: Axymyiidae). *Arthropod Systematics & Phylogeny*. 2013; 71(2): 91-102.
- [3] Oosterbroek P. The European families of Diptera: Identification - diagnosis - biology. 1st ed. Leiden: KNNV Publishing Brill. 2006.
- [4] Marshal SAI. Flies. The natural history and diversity of Diptera. 1st ed. Ontario: Firefly Books Ltd. 2012.
- [5] Wood DM. Axymyiidae. In: McAlpine JF, Peterson BV, Shewel GE, Teskey HJ, Vockeroth JR, Wood DM, eds. *Manual of Nearctic Diptera*. 1st ed. Ottawa: Canadian Government Publishing Centre; 1981. p. 1-674.
- [6] Matthew W, Courtney GW. The distribution and life history of *Axymyia furcata* McAtee (Diptera: Axymyiidae), a semi-aquatic wood-dwelling fly. *Proceedings of the Entomological Society of Washington*. 2011; 113(3): 385-398.
- [7] Polevoi A, Ruokolainen A, Shorohova E. Eleven notable species of Diptera, emerged from fallen poplars in Kivach Nature Reserve Russian Karelia. *Biodiversity Data Journal*. 2018; 6: e22175.
- [8] Martynovský J, Roháček J. First records of *Synneuron annulipes* Lundström (Synneuridae) and *Mesaxymyia kerteszi* (Duda) (Axymyiidae) from Slovakia, with notes on their taxonomy and biology (Diptera). *Natural Sciences*. 1993; 42(1):

73-78.

- [9] Roskov Y, et al. Species 2000 & ITIS catalog of life, species 2000. 1st ed. Leiden: Naturalis. 2020.
- [10] Scott JF, Wood DM. A new species of Axymyiidae (Diptera) from western North America and a key to Nearctic species. Zootaxa. 2014; 3857(1): 101-113.
- [11] Blagoderov VA, Lukashevich ED. New Axymyiidae (Insecta: Diptera) from the Mesozoic of Eastern Siberia. Polish Journal of Entomology. 2013; 82(4): 257-271.
- [12] Bradley J, Sinclair BJ. Finally, rediscovered: an enigmatic new genus of Axymyiidae (Diptera) from western North America. Zootaxa. 2013; 3682(1): 143-150.
- [13] Borkent A, Sinclair BJ. The male genital track of Axymyiidae and Tanyderidae (Diptera). The Canadian Entomologist. 2012; 144: 266–272.
- [14] Ševčík J, et al. Molecular phylogeny of the infraorder megadiverse insect Bibionomorpha sensu lato (Diptera). PeerJ. 2016; 4: e2563.
- [15] Krivosheina MG. Family Axymyiidae. In: Papp L, Darvas B, eds. Contributions to a manual of Palaearctic Diptera. 1st ed. Budapest: Science Herald; 2000. p. 31–39.
- [16] Mamaev BM, Krivosheina NP. Family Axymyiidae. In: Soós Á, Papp L. Catalog of Palaearctic Diptera. Sciaridae-Anisopodidae. 4th ed. Budapest: Akadémiai Kiadó; 1986. p. 317–318.