v1: 5 August 2024

PEER-APPROVED

Consumption of Beverages Among Secondary and Intermediate Students in Riyadh Schools, Saudi Arabia: A Cross-Sectional Study

Peer-approved: 5 August 2024

© The Author(s) 2024. This is an Open Access article under the CC BY 4.0 license.

Qeios, Vol. 6 (2024) ISSN: 2632-3834

Ali Bani Mfarrij¹, Hassan Faqeehi¹, Issam AlFayyad²

1. Children's Specialised Hospital, King Fahad Medical City, Riyadh, Saudi Arabia; 2. King Fahad Medical City, Riyadh, Saudi Arabia

Background: Water is essential for body hydration to maintain human mental and physical function. However, hot and humid weather, in addition to physical activity, can further increase fluid loss. We aimed to evaluate beverage consumption among school children aged 12-18 years in Riyadh, Saudi Arabia. Methods: A cross-sectional study was conducted using an online self-administered questionnaire among intermediate and secondary school students. A random stratification technique was used to recruit schools, followed by a non-random sampling technique of children's parents to recruit the adolescents aged 12-18 years. Descriptive statistics were used to summarize the participants' characteristics and beverage consumption. Results: A total of 1548 children completed the questionnaire with a mean age of (15.69±1.68) years. The majority (65.2%) were females. The daily mean total beverage consumption was (1918.88±796.01) ml/day for females and (2238.53±879.96) ml/day for males, (p=<0.001). Children aged (14-18) years' beverage consumption was (2052.41±842.01) ml/day and (1881.33±811.38) ml/day for children aged (12-13) years, (p=0.007). The highest consumed beverage for females and males was water (1124.94±548.36) ml/day and (1209.59±534.31) ml/day, respectively (p=0.004). Children aged (14-18) years consumed higher amounts of coffee (206.84±234.29) ml/day compared to those aged (12-13) years (78.68±140.71) ml/day, (p<0.001). Girls were drinking coffee (208.71±236.60) ml/day more than boys (158.36±206.54) ml/day (p<0.001).

Conclusion: Water is the main source of beverages in all age groups; however, beverage consumption was lower than the World Health Organization recommendation. The largest proportion of adolescents is at risk of inadequate hydration.

Correspondence: $\underline{papers@team.qeios.com}$ — Qeios will forward to the authors

Introduction

Water is essential for life $\frac{[1][2]}{1}$. Without water, we cannot survive. As the water content in our bodies frequently fluctuates, it needs to be continuously replenished, and the best liquid for this purpose is water itself because it hydrates our bodies without providing extra energy (calories/kilojoules) $\frac{[3]}{2}$. Body hydration is essential for adequate mental and physical function. Slight dehydration may result in damaging outcomes both physically and intellectually, and excessive dehydration may accentuate serious potential ailments and can ultimately lead to death $\frac{[4]}{2}$. Fluid requirements vary between each individual according to many determinants, including age, weight, phase of perspiration, and nutritional lifestyle $\frac{[4][5]}{2}$. The percentage of fluids in the human body ranges from 55% to 75%, and water is required for the correct functioning of every single part of the human body. Water aids the kidneys in removing waste from the circulation and excreting it outside the body as urine $\frac{[6]}{2}$.

Dehydration occurs when the intake of water and fluids is less than the amount that leaves the body. When water loss exceeds 5%, this is considered to be a severe case of dehydration $\frac{[7]}{}$. Also, dehydration has significant therapeutic impacts on several body systems $\frac{[8]}{}$, especially the kidneys. It has been recognized for quite a long period that drinking water helps avoid the formation of kidney stones. Extensive observational studies have suggested a therapeutic role in chronic kidney disease prevention $\frac{[9]}{}$. A fluid loss of 1% can lead to problems in physical activity, thermoregulation, and appetite. At 4% fluid loss,

severe issues are noticed, such as inattentiveness, headaches, irritability, lassitude, an increase in body temperature, and labored breathing. A fluid loss of 8% or more results in death $\frac{[10]}{10}$. In the United States of America, a National Health and Nutrition Examination Survey was conducted, and a 54% prevalence of dehydration was reported between the ages of 6 and 19 $\frac{[11]}{10}$.

With the weather variations due to global warming, it is thought that unpredictable weather events will take place. It is projected that the situation in Asia will be dreadful in the 21st century as temperatures are projected to increase to unacceptable levels ^[12]. Temperature changes have already been detected using climate indices ^{[13][14][15]}. In the city of Riyadh, especially in the summer, the temperature reaches 40-45 degrees Celsius (°C) ^[16]. Humid and hot weather, along with physical exertion, can increase fluid loss; as a result, our bodies demand more fluid intake ^[17].

To deal with temperature fluctuations, the body's cooling system releases a large amount of fluid. Perspiration helps cool down the body temperature in warm weather and during physical activity. Its level depends on the ambient temperature, humidity, type of effort, and outfit $\frac{1181}{1181}$. Therefore, the daily fluid requirements on hot days (40 °C) could be triple those on colder days (20 °C) to compensate for water loss $\frac{1191}{1181}$.

Euhydration is a significant factor in the prevention of accidents at work or the progression of disease. This is because it improves performance in both mental and physical tasks and enhances the perception of wellbeing. No matter how smooth, dehydration is not a tempting situation; it denotes that there is an imbalance within the homeostatic exertion of the internal background. This might aggressively impact the rational volume and interfere with the acceptable implementation of work or learning-related movements that necessitate the application of precise rational abilities $\frac{[7]}{2}$. According to the World Health Organization (WHO) Guidelines for Drinking-Water Quality, the adequate intake of water for adolescents (9-13 years old) is 2,400 L/day for boys and 2,100 L/day for girls, and the intake for (14-18 years old) is 3,300 L/day for boys and 2,300 L/day for girls.

Dry tongue and lips, thirst, dry skin, lethargy, exhaustion, headache, muscle weakness, dizziness, a feeling of faintness, and difficulty focusing are all common signs of mild to moderate dehydration. Among males, acute dehydration (1.59% water loss) without producing an increase in body temperature has constrained but discernible impacts on particular aspects of cognitive performance and causes negative changes in mood, anxiety, and fatigue, which are related to antagonistic changes in cognitive performance in some circumstances. This idea is backed up by the findings in females at a comparative level of dehydration (1.39% water loss), which also caused small changes in cognitive performance but significant antagonistic changes in mood. Both genders experienced unfavorable but, to some degree, distinctive behavioral impacts of dehydration below 2% weight loss [21]. Such a small drop in body weight can occur during regular activities [22]. illustrating the significance of maintaining ideal hydration. However, people with medical conditions that can potentially cause dehydration, such as diabetics, along with the elderly and children, may be the most likely to encounter antagonistic behavioral impacts of slight dehydration [23]

In Saudi Arabia, the public's awareness of dehydration is low. A prior study of people in Riyadh, Saudi Arabia, to examine dehydration awareness and fluid consumption found that while participants knew the typical symptoms of dehydration, they didn't know the unusual symptoms or significant implications of dehydration ^[24]. Estimates of dehydration knowledge based on globally published studies may have limited applicability across fields, climates, and cultures. Reporting climate- and culturally valid data regarding hydration status and beverage consumption from intermediate and secondary school students might inform Saudi decision-makers, promote health policy, and establish programs that coach students on the importance of drinking healthy beverages. Due to Saudi Arabia's climate changes, and as the country's weather is becoming increasingly hot $\frac{[25]}{}$, we aimed in this study to evaluate the amounts and sources of fluids consumed by adolescents (12-18 years old) in Riyadh schools, Saudi Arabia, to determine whether these amounts and sources are sufficient and appropriate for adequate daily intake.

Materials and Methods

Study design and settings

A multicenter, cross-sectional study was conducted in the governmental schools in five regions of Riyadh City, namely, the north, south, east, west, and central regions. An online self-administered questionnaire was employed for data collection due to the COVID-19 pandemic.

Study population and sampling technique

The list of governmental schools in Riyadh City was obtained from the General Directorate of Education in Riyadh City. A stratified random sampling technique was used to select the schools (Boys' and girls' sections) according to the geographical area from each region. Two levels of stratification were used: the region strata (North, South, West, East, and Central) and the gender strata (Male and female). Accordingly, a total of 20 schools were selected from the five regions (4 from each region, 2 male schools, and 2 female schools). Then, a non-random sampling technique of children's parents was used to recruit the adolescents aged between 12–18 years (secondary and intermediate students). The schools' administrations were contacted and asked to

provide us with parents' WhatsApp groups to disseminate the study questionnaire. All students were invited to participate after the parents confirmed that their children meet the eligibility criteria described on the cover of the questionnaire.

The inclusion criteria were Saudi students aged between 12 and 18 in the designated schools. Any student who complained of significant medical diseases like renal failure, renal tubulopathy, diabetes insipidus, diabetes mellitus, chronic diarrhea, or cystic fibrosis was excluded from this study.

Data collection instrument

Due to the COVID-19 pandemic, an online questionnaire was employed for data collection and sent to the parents through parents' WhatsApp groups between March 2021 and June 2021. The parents were asked to assist their children in approximately evaluating their children's consumption of beverages. Incorporating the parents helped us to minimize the bias of children's comprehension of the study questionnaire.

This study survey was designed based on detailed literature reviews ^[25] ^{[26][27]}, study team professional experiences in nephrology care, and the recommendations of two pediatric nephrologists/experts. The first part of the survey consisted of the demographic and clinical data of the students. The second part assessed the amounts of beverages consumed, including water, juices, milk, coffee, tea, and soda. In this questionnaire, we used images labelled with their size for each type of beverage to facilitate children's comprehension of the survey questions and to reach as accurate an evaluation of beverage consumption as possible. The last part assessed the knowledge of the recommended amount of beverage intake per day.

Sample size calculation

Based on the literature, the prevalence of inadequate hydration in the USA is 54.4% (11), as well as an estimated 288,125 Saudi students in governmental schools at the secondary and intermediate stages [28]. The current study sample size is $n = [z\alpha/2p * q/(d * d)]$. p = estimated prevalence = 0.54, q = 1 - p = 0.46, d = estimating allowable error in the prevalence (margin of error) = 10% on either side of the stated prevalence of 54.4% = 0.054, otherwise power of study = 90%, α = probability of type I error = 0.05 (two-sided), z = 1.96; therefore, n = 1,497. With a nonresponse rate of 10%, the number of respondents needed for the study was 1,647 students.

Statistical analysis plan

Data were analyzed using the Statistical Package for the Social Sciences (SPSS) version 25.0 (Inc., Chicago, IL, USA). Descriptive statistics (mean, standard deviation, and percentages) were used to summarize the participants' characteristics and the distribution of total beverage consumption. T-tests were used to compare the means of beverage consumption. A p-value less than 0.05 was considered to indicate significance.

Ethics approval and consent to participate

Ethical approval was obtained from the Institutional Review Board of King Fahad Medical City (19-465). The questionnaire included a cover letter with informed consent indicating that completion of the questionnaire implies their informed consent to allow their children to participate in the study. Official approval was also obtained from the General Directorate of Education in Riyadh, Ministry of Education. Participants' identities were not captured.

Results

A total of 1647 online invitations were distributed, and 1548 participants responded and completed the questionnaire, with a response rate of

94%. The mean age of participants was 15.69±1.68, and the majority, 1345 (86.9%), were between the age group 14–18 years. Nearly two-thirds, 1010 (65.2%), were females. Participants were stratified by gender and age, and the majority of 907 (58.6%) were females aged 14–18 years, followed

by males, 438 (28.2%), aged 14-18 years. The majority of respondents, 1102 (71.2%), were from secondary schools, and 25 (1.6%) had significant medical conditions. Table 1

Table 1. Characteristics of 1548 students from Riyadh, Saudi Arabia	
Age (mean±SD)	15.69±1.68
Age (years)	
12-13	203 (13.1)
14-18	1345 (86.9)
Gender	
Female	1010 (65.2)
Male	538 (34.8)
Gender by Age	
Female (12-13)	103 (6.7)
Male (12-13)	100 (6.5)
Female (14-18)	907 (58.6)
Male (14-18)	438 (28.2)
School stage	
Intermediate	446 (28.8)
Secondary	1102 (71.2)
Height (mean±SD)	161.47±10.23
Weight (mean±SD)	59.64±14.82
Significant medical diseases	
Yes	25 (1.6)
No	1523 (98.4)

Table 2 shows that children aged 14–18 years for beverage consumption were higher than children aged 12–13 years, with a statistically significant difference (p=0.007). Males' beverage consumption (2238.53 \pm 879.96) was higher than females' (1918.88 \pm 796.01), with a statistically significant difference (p=<0.001). Furthermore, male

children in both age groups consumed beverages more than female children in the corresponding age group, with statistically significant differences (p=<0.001; for both). The stratification of beverage intake among secondary and intermediate shows no statistical difference (p=139). Table 2

	mean±SD	P-value
Age		
12-13	1881.33±811.38	0.007
14-18	2052.41±842.01	
Gender		
Female	1918.88±796.01	<0.001
Male	2238.53±879.96	
Gender by Age		
Female (12-13)	1755.93±712.07	<0.001
Male (12-13)	2010.50±887.50	
Female (14–18)	1937.39±803.26	
Male (14-18)	2290.59±870.89	
School stage		
Intermediate	1980.34±884.14	0.139
Secondary	2050.06±820.72	

Table 2. Comparison of total consumed amount of beverages by the participants with their characteristics

Beverage consumption

For water consumption, males consumed a higher amount (1209.59±534.31) than females (1124.94±548.36), and those aged 12–13 years (1051.77±529.92) reported less water intake than children aged 14–18 years (1169.84±545.58). Tabulation of water consumption showed statistically significant differences with age (p=0.004), gender (p=0.004), gender by age (p<0.001), and school stage (p=0.005). Table 3.

Boys (206.69±224.30) were significantly getting more juice than girls (123.76±149.83), younger adolescents aged 12-13 years (204.93±203.38) were drinking juice more than those aged 14-18 years (144.68±179.04), and the intermediate school age group (180.27±196.98) was drinking juice more than the secondary school age group (141.38±176.59). Tabulation of juice consumption showed statistically significant differences with age, gender, gender by age, and school stage, (p=<0.001; for all).

The results show less milk intake among children 14–18 years of age (138.88±198.01) compared to those aged 12–13 years of age (177.34±200.20). Boys (166.17±223.67) were drinking more milk than girls (132.08±182.97), and secondary school children (133.21±188.61) were drinking less milk than the children at the intermediate school (170.40±219.55). Tabulation of milk consumption showed statistically

significant differences with age, gender, gender by age, and school stage, (p=<0.010; for all). Table 3

The results show higher coffee intake among children aged 14–18 years (206.84±234.29) compared to those aged 12–13 years (78.68±140.71). Girls (208.71±236.60) were drinking more coffee than boys (158.36±206.54). Stratification of age and gender showed that girls in both age groups consumed more coffee than boys. Moreover, children in the secondary school (213.25±240.81) consumed a higher amount of coffee than those in the intermediate school (136.77±180.99). Tabulation of coffee consumption showed statistically significant differences with age, gender, gender by age, and school stage, (p=<0.001; for all). Table 3

Table 3 shows higher tea intake among children aged 14-18 years (170.11 \pm 227.10) compared to those aged 12-13 years (132.02 \pm 169.78). Boys reported higher tea consumption (202.364 \pm 256.02) than girls (147.72 \pm 202.36). Stratification of age and gender showed that boys in both age groups consumed more tea than girls. Tabulation of tea consumption showed statistically significant differences with age (p=0.024), gender (p=<0.001), and gender by age (p=<0.001). Coca consumption was significantly (p<0.001) higher among male children (299.94 \pm 351.29) than females (181.66 \pm 314.07). Stratification of age and gender showed that boys in both age groups consumed more coca than girls, (p<0.001). Table 3

	Water	p- value	Juice	p- value	Milk	p- value	Coffee	p- value	Tea	p- value	Coca	p- value
Age (year) 12-13 14-18	1051.77±529.92 1169.84±545.58	0.004	204.93±203.38 144.68±179.04	<0.001	177.34±200.20 138.88±198.01	0.010	78.68±140.71 206.84±234.29	<0.001	132.02±169.78 170.11±227.10	0.024	227.59±343.96 222.04±330.50	0.825
Gender Female Male	1124.94±548.36 1209.59±534.31	0.004	123.76±149.83 206.69±224.30	<0.001	132.08±182.97 166.17±223.67	0.010	208.71±236.60 158.36±206.54	<0.001	147.72±202.36 202.364±256.02	<0.001	181.66±314.07 299.94±351.29	<0.001
Gender by Age Female (12-13) Male (12-13) Female (14-18) Male (14-18)	1079.71±551.02 1023.00±508.43 1130.08±548.13 1252.19±531.48	<0.001	165.05±161.92 246.00±232.43 119.07±147.76 197.72±221.69	<0.001	143.69±188.74 212.00±206.60 130.76±182.37 155.71±226.31	0.010	104.85±145.10 70.001±34.47 220.51±242.09 178.54±214.78	<0.001	118.45±159.50 146.00±228.93 151.05±206.48 209.59±260.61	<0.001	144.17±220.02 313.50±420.51 185.92±322.84 296.85±333.97	<0.001
School stage Intermediate Secondary	1092.85±529.50 1179.26±549.21	0.005	180.27±196.98 141.38±176.59	<0.001	170.40±219.55 133.21±188.61	0.010	136.77±180.99 213.25±240.81	<0.001	152.91±217.09 170.05±226.20	0.172	247.13±346.24 212.91±325.97	0.066

Table 3. Comparison of the participants' consumed amounts (ml) of the beverages with their characteristics

Discussion

The current study aims to describe beverage consumption and its variation by age. The results show significant differences in the quantity and pattern of beverage consumption across the age and sex groups examined. The total mean fluid intake for males aged 12-13 years was 2011 mL/day, which is lower than the WHO recommendation for adequate intake for adolescents (9-13 years old) of 2,400 mL/day [20]. For females of the same age group, the mean fluid intake was 1756 mL/day, which is also lower than the WHO recommendation of 2,100 mL/day [20]. For other age groups 14-18 years, males' mean fluid consumption was 2291 mL/day, which is lower than the WHO recommendation of 3,300 mL/day, and for females aged 14-18 years, it was 1937 mL/day, which is also lower than the WHO recommendation of 2,300 mL/day. Sui et al. (2016) mentioned that the mean daily total water intake for children and adolescents (2-18 years old) is 1.7 L for boys and 1.5 L for girls [29]. Kant and Graubard (2010) mentioned that the adjusted mean intake of water in Americans aged 6-11 and 12-19 years is 1.6 L and 2.4 L, respectively. In particular, the average total water intake is insufficient; in general, the intake of boys was the least adequate [30]. Moreover, Kenney and Long (2009) mentioned that insufficient hydration is widespread among American children and adolescents, especially boys. Proper intake of water can reduce the risks of inadequate hydration [11].

According to the findings, water was the most significant contributor to fluid consumption (Males 1125 mL/day, Females 1210 mL/day), which is similar to the study published by Duffey et al. (2012) ^[26]. Moreover, Guelinckx et al. (2015), in a study on water and beverage intake among children and adolescents in 13 countries (from Latin America, Europe, and Asia), showed that across all the samples of these countries, the highest daily intake was observed for water, followed by milk, regular soft beverages, and juices ^[31]. One remarkable point throughout this analysis is that Coca-Cola drinks were the second contributor of fluids

(Males 300 mL/day, Females 182 mL/day), followed by coffee drinks (Males 158 mL/day, Females 209 mL/day), then juice (Males 207 mL/day, Females 124 mL/day), and the fifth contributor was tea (Males 202 mL/day, Females 148 mL/day), and the lowest was milk (Males 166 mL/day, Females 132 mL/day). Bello and Al-Hammad (2006) mentioned that potable water accounted for 37%, whereas fruit juices and carbonated soft drinks accounted for 25% and 26%, respectively ^[32]. Vieux et al. (2017) reported that the total intake of plain water is 1,338 mL/day, the intake of drinking water, coffee, soda, and tea increases with age, whereas the intake of milk decreases. Approximately 88.7% of children do not fulfill the European Food Safety Authority (EFSA) adequate intake criteria. The daily water shortfall ranged from 322 to 659 mL/day. The total water intake was at 74.8% of the EFSA reference standards ^[33].

The data were collected between February and June of this year, 2021, in Riyadh City, during the winter to summer seasons. In the summer, the temperature reaches 40-45°C (16). Children in hot regions may be more susceptible to heat sickness than adults because of their larger surface area to body mass ratio, lower rate of sweating, and a bit slower rate of heat acclimation [34][35]. Children may respond to hypohydration during activity with a larger relative increase in core temperature than adults do, and they may sweat less, reducing the advantages of evaporative cooling. According to Coyle (2004), fluid should be absorbed at rates that are most similar to sweating rates. Whenever that is not feasible, some sportsmen may be able to tolerate body water losses of up to 2% of body weight without affecting their physical well-being or performance when the weather is cold (5-10 °C) or temperate (21-22 °C). When exercising in a hot climate (>30 °C), however, dehydration by 2% of body weight impedes absolute power generation and predisposes individuals to heat injury [36]

This study has some limitations. The study is cross-sectional, based on self-reported beverage consumption. The self-reported drinks intake might have been overestimated or underestimated. Furthermore,

dietary data are mean values of a 24-hour recall, which is unlikely to be enough time to adequately reflect typical consumption, particularly in children and adolescents, where the ratio of within-subject variability is higher [37]. Selection bias might be present, as parents who have limited access to WhatsApp groups or are not interactive using this approach were likely to avoid participation. Measurement bias might also be present when comparing respondents who were assisted by their parents to respondents who were not assisted by their parents.

Conclusions

This study provides important information about beverage consumption in adolescents. Euhydration is important in maintaining mental and physical function, and dehydration is a risk factor for many health problems. A growing body of data demonstrates the link between fluid imbalances and diseases; however, the evidence is largely associative and inconsistent, with a limited number of published studies. Tap water is the main source of consumption for all ages, and consumption of other beverages varies by age. Coca-Cola drinks came second after water, coffee was third, followed by juice, then tea, and the lowest was milk. The consumption of soft drinks and mainly tap water should be increased.

Abbreviations

WHO: World Health Organization; COVID-19: Coronavirus Disease of 2019; EFSA: European Food Safety Authority; ℃: Degree Celsius; SPSS: Statistical Package for the Social Sciences.

Statements and Declarations

Ethics approval and consent to participate: Ethical approval was obtained from the Institutional Review Board of King Fahad Medical City (19-465). Official approval was also obtained from the General Directorate of Education in Riyadh, Ministry of Education. Participants' identities were not captured.

Consent for publication: Not Applicable.

Availability of data and materials: The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interests.

Funding: This work was supported by the Research Center at King Fahad Medical City [Grant #: 021-005].

Authors' contributions: AM and HF conceptualized the study. AM and IF collected the data and conducted the statistical analysis. AM and HF drafted the manuscript. AM and IF finalized the final manuscript. All authors approved the final version of this manuscript.

Acknowledgments: None.

Other References

 Ministry of Education. Statistics of governmental Education [Internet]. e-services. [cited 2019Jul31]. Available from: <u>https://departments.moe.gov.sa/Statistics/Educationstatistics/Pages/GESt</u>

References

- ^AIglesia I, Guelinckx I, De Miguel-Etayo PM, González-Gil EM, Salas-Sal vadó J, Kavouras SA, et al. Total fluid intake of children and adolescents: cross-sectional surveys in 13 countries worldwide. European journal of nutrition. 2015;54(2):57-67.
- A[¬]Özen A, Bibiloni MdM, Pons A, Tur J. Fluid intake from beverages acros s age groups: a systematic review. Journal of Human nutrition and Diete tics. 2015;28(5):417-42.

- 3. ^ADrewnowski A, Rehm CD, Constant F. Water and beverage consumptio n among children age 4-13y in the United States: analyses of 2005–2010 NHANES data. Nutrition journal. 2013;12(1):85.
- 4. ^{a.} ^bLieberman HR. Hydration and cognition: a critical review and recom mendations for future research. Journal of the American College of Nutr ition. 2007;26(sup5):555S-61S.
- 5. ^ABellisle F, Thornton S, Hebel P, Denizeau M, Tahiri M. A study of fluid in take from beverages in a sample of healthy French children, adolescents and adults. European journal of clinical nutrition. 2010;64(4):350.
- ^AArnaud MJ. Hydration Throughout Life: International Conference, Vitte l, France, June 9-12, 1998: John Libbey Eurotext; 1998.
- 7. ^{a. b}Adan A. Cognitive performance and dehydration. Journal of the Ame rican College of Nutrition. 2012;31(2):71-8.
- Ackleiner SM. Water: an essential but overlooked nutrient. Journal of the American Dietetic Association. 1999;99(2):200-6.
- ^ABouby N, Clark WF, Roussel R, Taveau C, Wang CJ. Hydration and kidne y health. Obesity facts. 2014;7(Suppl. 2):19-32.
- 10. [△]Grandjean AC, Reimers KJ, Buyckx ME. Hydration: issues for the 21st ce ntury. Nutrition reviews. 2003;61(8):261-71.
- 11. ^{a.} <u>b</u>Kenney EL, Long MW, Cradock AL, Gortmaker SL. Prevalence of inad equate hydration among US children and disparities by gender and rac e/ethnicity: National Health and Nutrition Examination Survey, 2009–2 012. American journal of public health. 2015;105(8):e113-e8.
- 12. [△]Pal JS, Eltahir EA. Future temperature in southwest Asia projected to e xceed a threshold for human adaptability. Nature Climate Change. 201 6;6(2):197.
- 13. [△]Alghamdi AS, Moore TW. Analysis and comparison of trends in extrem e temperature indices in Riyadh City, Kingdom of Saudi Arabia, 1985–2 010. Journal of Climatology. 2014;2014.
- <u>A</u>Athar H. Trends in observed extreme climate indices in Saudi Arabia d uring 1979–2008. International Journal of climatology. 2014;34(5):1561– 74.
- 15. ^AAl-Saleh MA. Variability and frequency of daily rainfall in Riyadh, Sau di Arabia. The Geographical Bulletin. 1997;39(1):48-57.
- 16. ^ARaggad B. Stationary and Non-stationary Extreme Value Approaches f or Modelling Extreme Temperature: the Case of Riyadh City, Saudi Arab ia. Environmental Modeling & Assessment. 2018;23(1):99–116.
- 17. [△]Medicine CoS, Fitness. Climatic heat stress and the exercising child and adolescent. Pediatrics. 2000;106(1):158-9.
- ^APopkin BM, D'Anci KE, Rosenberg IH. Water, hydration, and health. Nut rition reviews. 2010;68(8):439–58.
- ^ATani Y, Asakura K, Sasaki S, Hirota N, Notsu A, Todoriki H, et al. The inf luence of season and air temperature on water intake by food groups in a sample of free-living Japanese adults. European journal of clinical nut rition. 2015;69(8):907.
- 20. ^{a, b, c}Grandjean AC. Water requirements, impinging factors, and recomm ended intakes. Nutrients in drinking water. 2005:25.
- 21. ^AArmstrong LE, Ganio M, Lee E, McDermott B, Klau J, Yamamoto L, et a l. Mild dehydration degrades mood and symptoms, not cognitive perfor mance in females: a placebo-controlled study. Federation of American S ocieties for Experimental Biology; 2010.
- 22. [≜]Sawka MN, Cheuvront SN, Carter R. Human water needs. Nutrition rev tats akyys. 2005;63(supp[_]):S30–S9.
- 23. [△]Ganio MS, Armstrong LE, Casa DJ, McDermott BP, Lee EC, Yamamoto L M, et al. Mild dehydration impairs cognitive performance and mood of men. British Journal of Nutrition. 2011;106(10):1535-43.
- 24. ^AAlbasheer OB, Hakami A, Al Faqih AA, Akkam I, Soraihy SK, Mathary A, et al. Awareness of dehydration state and fluid intake practice among adults population in the Jazan Region of Saudi Arabia, 2019. Journal of Nutritional Science. 2021;10.
- 25. ^{a, b}Shaheen NA, Alqahtani AA, Assiri H, Alkhodair R, Hussein MA. Publi c knowledge of dehydration and fluid intake practices: variation by part icipants' characteristics. BMC public health. 2018;18(1):1-8.

- 26. ^{a.} ^bDuffey KJ, Huybrechts I, Mouratidou T, Libuda L, Kersting M, De Vrien dt T, et al. Beverage consumption among European adolescents in the H ELENA study. European journal of clinical nutrition. 2012;66(2):244-52.
- 27. [△]Laksmi PW, Morin C, Gandy J, Moreno L, Kavouras S, Martinez H, et al. Fluid intake of children, adolescents and adults in Indonesia: results of t he 2016 Liq. In 7 national cross-sectional survey. European journal of nu trition. 2018;57(3):89-100.
- ^AMinistry of Education. Statistics of governmental Education [Internet]. e-services. [cited 2019Jul31].Available from: https://departments.moe.go v.sa/Statistics/Educationstatistics/Pages/GEStats.aspx
- 29. [△]Sui Z, Zheng M, Zhang M, Rangan A. Water and beverage consumptio n: analysis of the Australian 2011–2012 National Nutrition and Physical Activity Survey. Nutrients. 2016;8(11):678.
- 30. [△]Kant AK, Graubard BI. Contributors of water intake in US children and adolescents: associations with dietary and meal characteristics—Nation al Health and Nutrition Examination Survey 2005–2006. The America n journal of clinical nutrition. 2010;92(4):887-96.

- 31. ^AGuelinckx I, Iglesia I, Bottin J, De Miguel-Etayo P, González-Gil E, Salas -Salvado J, et al. Intake of water and beverages of children and adolesce nts in 13 countries. European journal of nutrition. 2015;54(2):69-79.
- 32. [△]Bello L, Al-Hammad N. Pattern of fluid consumption in a sample of Sa udi Arabian adolescents aged 12–13 years. International Journal of Paed iatric Dentistry. 2006;16(3):168-73.
- 33. [△]Vieux F, Maillot M, Constant F, Drewnowski A. Water and beverage con sumption patterns among 4 to 13-year-old children in the United Kingd om. BMC Public Health. 2017;17(1):1-12.
- 34. [△]Falk B, Dotan R. Children's thermoregulation during exercise in the hea t—a revisit. Applied Physiology, Nutrition, and Metabolism. 2008;33(2):4 20-7.
- 35. [△]Bytomski JR, Squire DL. Heat illness in children. Current sports medicin e reports. 2003;2(6):320-4.
- 36. $^{\wedge}$ Coyle EF. Fluid and fuel intake during exercise: Routledge; 2004.
- 37. [△]Livingstone M, Robson P. Measurement of dietary intake in children. P roceedings of the Nutrition Society. 2000;59(2):279–93.

Declarations

Funding: Research Center at King Fahad Medical City [Grant #: 021–005]. Potential competing interests: No potential competing interests to declare.