

Short Communication

Timed Acupuncture-Mediated Memory Consolidation as a Strategy to Reset Pathological Homeostatic Set-Points: A Three-Step Hypothesis

Zicheng Cao¹

1. Longyan Hospital of Traditional Chinese Medicine, China

Chronic non-communicable diseases—including hypertension, type 2 diabetes, obesity, and chronic pain syndromes—are characterized by pathological shifts in homeostatic set-points within the hypothalamus and related neuroendocrine circuits. Current pharmacologic interventions transiently normalize physiological parameters but fail to reverse the underlying set-point deviation, resulting in relapse upon treatment withdrawal. We propose that homeostatic regulation shares fundamental mechanisms with memory consolidation: a transiently achieved healthy physiological state, if accompanied by appropriate neurochemical signals during a critical temporal window, can be consolidated into a new long-term set-point. Acupuncture is uniquely positioned to provide such signals through its capacity to release β -endorphin, endocannabinoids, oxytocin, and to modulate hypothalamic-pituitary-adrenal (HPA) axis activity in a temporally precise manner. We outline a three-step intervention strategy: (1) induce a comprehensive healthy physiological state via pharmacological or behavioral means; (2) administer acupuncture at specific acupoints during a circadian-defined memory consolidation window to release neuromodulatory factors that mark this state for long-term storage; (3) deliver periodic maintenance acupuncture to reinforce the new set-point until it becomes the system default. Specific testable predictions are provided. If validated, this hypothesis transforms acupuncture from a symptom-modulation technique into a precision tool for homeostatic reprogramming.

Correspondence: papers@team.qeios.com — Qeios will forward to the authors

1. Introduction

The pathophysiology of chronic metabolic and pain disorders involves more than peripheral organ dysfunction. Converging evidence indicates that these conditions represent stable attractors in a multi-level homeostatic control system centered on the hypothalamus. For example, obesity shows sustained upregulation of the hypothalamic NPY/AgRP orexigenic pathway and downregulation of the POMC/CART anorexigenic pathway despite caloric restriction [1]. Similarly, chronic pain involves central sensitization maintained by altered descending modulation from the periaqueductal gray and rostral ventromedial medulla [2]. The existence of such disease attractors explains the phenomenon of “set-point inertia”: even when external interventions temporarily normalize physiological parameters, the system spontaneously returns to its pathological baseline upon intervention withdrawal.

Current therapies are effective at symptom control but not at reprogramming the underlying set-point. Pharmacologic agents such as antihypertensives, hypoglycemics, and GLP-1 receptor agonists produce robust acute effects, yet discontinuation almost invariably leads to relapse. Bariatric surgery achieves lasting weight loss in some patients but is irreversible, costly, and carries surgical risk [3].

Recent advances in systems neuroscience suggest that homeostatic regulation shares computational principles with memory consolidation. During sleep, hippocampal–cortical replay strengthens newly encoded information [4]; during reconsolidation, retrieved memories become labile and can be updated [5]. This raises a critical therapeutic question: can a transiently achieved healthy physiological state be “remembered” by the homeostatic system as the new normal?

We propose that acupuncture can serve as the neuromodulatory “permission signal” to consolidate a transient healthy state into a long-term homeostatic set-point. Acupuncture stimulates the release of multiple neuromodulators—including β -endorphin, endocannabinoids, oxytocin, serotonin, and dopamine—while modulating HPA axis activity and vagal tone [6][7]. These neurochemicals are known to facilitate or gate synaptic plasticity and memory consolidation [8]. Importantly, acupuncture effects exhibit circadian sensitivity: stimulation at different times produces distinct hormonal responses [9]. This chronobiological property may allow targeting of specific consolidation windows.

2. The Hypothesis

We propose that a transiently achieved healthy physiological state can be consolidated into a long-term homeostatic set-point through a sequence of three interventions:

Step 1: Induction of a transient healthy state. A pharmacological, nutritional, or behavioral intervention (e.g., GLP-1 agonist for weight loss, intensive antihypertensive therapy for blood pressure normalization, or a controlled fasting protocol for metabolic reset) is applied to bring the organism into a full, healthy physiological state. The state must be comprehensive: metabolic parameters, autonomic tone, and neuroendocrine markers all within normal range. This state is inherently unstable; without further intervention, the system will revert to its prior pathological set-point.

Step 2: Acupuncture-mediated consolidation during the critical temporal window. During the period immediately following Step 1—corresponding to the consolidation window for homeostatic memory—acupuncture is administered at acupoints selected to activate specific neuroendocrine pathways. Based on the known circadian rhythm of cortisol secretion, glucocorticoid receptor sensitivity, and cholinergic tone, we hypothesize that this window optimally occurs in the late evening (pre-sleep period) when the HPA axis is at its nadir and sleep-dependent consolidation mechanisms can be engaged. The acupuncture stimulus triggers release of β -endorphin (from the arcuate nucleus), endocannabinoids (from peripheral and central sources), and oxytocin (from the paraventricular nucleus), which serve as “permission signals” for long-term potentiation in hypothalamic circuits encoding the new homeostatic set-point. Specifically, the coordinated release of these factors during a window of heightened synaptic plasticity enables the conversion of transient physiological data into stable engrams within the hypothalamus, amygdala, and autonomic brainstem nuclei.

Step 3: Maintenance reinforcement. Following initial consolidation, periodic acupuncture sessions are administered at increasing intervals (e.g., daily for 1 week, then every other day for 2 weeks, then weekly) to strengthen the newly encoded set-point. This period leverages the reconsolidation process: each retrieval of the new homeostatic state during maintenance sessions recruits molecular machinery that stabilizes the memory trace. After approximately 4–6 weeks, the new set-point becomes the system default and no longer requires periodic reinforcement.

3. Evidence Supporting The Hypothesis

3.1. *Acupuncture releases neuromodulators that gate memory consolidation*

Acupuncture at specific acupoints reliably increases plasma β -endorphin and enkephalin levels within 15–30 minutes [6]. Electroacupuncture (EA) at 2 Hz preferentially stimulates β -endorphin release via the arcuate nucleus–periaqueductal gray pathway, while 100 Hz EA targets dynorphin release via the spinal cord [10]. Acupuncture also elevates endocannabinoid levels—including anandamide and 2-arachidonoylglycerol—in both plasma and brain tissue [11]. Oxytocin release from the paraventricular nucleus following acupuncture has been demonstrated in rodent models of chronic pain [12].

These neuromodulators are known regulators of synaptic plasticity and memory consolidation. β -endorphin facilitates hippocampal long-term potentiation (LTP) via mu-opioid receptors [13]. Endocannabinoids mediate depolarization-induced suppression of inhibition (DSI) and can gate the induction of LTP in the hippocampus and amygdala [14]. Oxytocin modulates social memory consolidation and enhances LTP in the central amygdala [15]. The coordinated release of multiple consolidation-promoting factors during acupuncture creates a neurochemical milieu permissive for stable encoding of new homeostatic information.

3.2. *Acupuncture modulates HPA axis and autonomic function in a temporally specific manner*

Multiple studies demonstrate that acupuncture normalizes HPA axis hyperactivity in chronic stress models. A 2012 study demonstrated that acupuncture at HT7 (Shenmen) significantly decreased plasma corticosterone levels in rats subjected to chronic CORT administration, and this normalization was associated with restoration of spatial memory function and cholinergic neuron integrity in the hippocampus [16]. The study concluded that acupuncture “could influence secretion of CORT, thereby normalizing behavioral and neurochemical response,” suggesting that acupuncture’s modulation of the HPA axis is both temporally sensitive and functionally relevant to cognitive and homeostatic processes. This normalization is time-dependent: the magnitude of HPA modulation varies with the circadian phase of stimulation, with maximal effects observed during the light (rest) phase in rodents [9].

In obesity models, EA at ST36 (Zusanli) restores vagal tone and reduces sympathetic dominance, as measured by heart rate variability. A 2025 study published in *Chinese Medicine* demonstrated that the

therapeutic benefits of EA in appetite suppression and obesity mitigation are mediated by selective regulation of the “vagal–GLP-1–ARC” neural circuit, providing direct evidence for acupuncture-induced remodeling of central metabolic regulatory pathways [17]. Importantly, these autonomic effects persist beyond the stimulation period, suggesting that the intervention induces a lasting change in the set-point of autonomic regulation [18].

3.3. Acupuncture produces long-lasting effects consistent with set-point resetting

Clinical and animal studies report sustained therapeutic effects of acupuncture that outlast the treatment period. In a meta-analysis of acupuncture for hypertension, 10–20 sessions produced blood pressure reductions that persisted for 3–6 months after treatment discontinuation [19]. A rat model of diet-induced obesity showed that 4 weeks of EA was sufficient to produce weight loss and metabolic improvements that lasted at least 8 weeks post-treatment [20]. These observations are difficult to explain through acute modulation alone and suggest a mechanism involving lasting reorganization of homeostatic set-points.

3.4. Precedents: memory reconsolidation mediated by acupuncture

The hypothesis that acupuncture can engage memory reconsolidation mechanisms is supported by a recent proposal for treating PTSD. In a 2022 article published in *Translational Psychiatry*, the authors proposed that “memory-directed acupuncture” can serve as a neuromodulatory treatment for PTSD by inducing memory reconsolidation of traumatic memories, depotentiating maladaptive fear responses [21]. Their theoretical framework—which explicitly links acupuncture stimulation to the molecular machinery of memory reconsolidation—provides a direct precedent for our proposal: if acupuncture can destabilize and update pathological memory traces, it may also consolidate newly established adaptive ones. We extend this logic from emotional memory to homeostatic memory, proposing that the same reconsolidation mechanisms can be harnessed to stabilize a drug-induced healthy physiological state.

4. Testable Predictions

We propose the following specific, falsifiable predictions:

Prediction 1 (time window specificity): Acupuncture administered within a 2-hour window before sleep (the hypothesized consolidation period) will produce significantly greater set-point resetting than acupuncture administered at other times. Measured as: (a) change in homeostatic parameter (e.g., fasting

glucose, mean arterial pressure, body weight) from baseline at 8 weeks post-intervention; (b) percentage of drug-free days after medication withdrawal.

Prediction 2 (acupoint specificity): Acupuncture at neuroendocrine-regulating acupoints (e.g., HT7, ST36, CV4) will be superior to non-acupoint stimulation or sham acupuncture for set-point resetting, whereas acupoints with primarily local analgesic effects (e.g., LI4, GB34) will show no advantage over sham in this paradigm.

Prediction 3 (neuromodulator mediation): Pharmacologic blockade of the μ -opioid receptor (e.g., naloxone) immediately before Step 2 acupuncture will abolish the set-point resetting effect, as measured by the relapse rate.

Prediction 4 (biomarker signature): Successful set-point resetting will be associated with specific changes in hypothalamic resting-state functional connectivity (fMRI), increased hippocampal-prefrontal theta coherence (EEG during sleep), and elevated plasma oxytocin and β -endorphin during the consolidation window.

Prediction 5 (comparison to continuous intervention): A single optimized cycle of Step 1-3 (e.g., 2 weeks of drug + 4 weeks of consolidation acupuncture) will produce comparable or superior long-term outcomes compared to continuous drug therapy alone followed by abrupt cessation, in a randomized controlled trial.

5. Discussion and Limitations

We have presented a hypothesis for transforming acupuncture into a set-point reprogramming tool by co-opting the brain's memory consolidation mechanisms. The hypothesis reconciles several previously disconnected observations: the long-lasting effects of acupuncture, the circadian sensitivity of its efficacy, the role of neuropeptides in both acupuncture and memory, and the phenomenon of relapse upon drug withdrawal.

Potential limitations include: (1) The precise molecular mechanisms by which a physiological state is encoded as a homeostatic memory remain undefined. (2) Individual differences in circadian phase, glucocorticoid sensitivity, and opioid receptor polymorphisms may modify the consolidation window. (3) The optimal protocol (acupoint selection, stimulation parameters, duration, and number of sessions) requires empirical determination. (4) The hypothesis does not address whether the consolidated set-point is stable under challenge (e.g., high-fat diet after obesity treatment). (5) This hypothesis is based on

theoretical integration of existing findings; direct experimental evidence for the complete three-step sequence is currently lacking.

Nevertheless, the hypothesis generates clear, testable predictions amenable to investigation in both animal models and human clinical trials. If validated, this approach would represent a paradigm shift: rather than treating chronic disease as a lifelong pharmacotherapy problem, clinicians could aim for a time-limited intervention that resets the homeostatic system.

6. Conclusion

The convergence of acupuncture neuroscience, memory consolidation research, and homeostatic control theory suggests a novel therapeutic strategy: use acupuncture as a temporally precise neuromodulatory signal to consolidate a transiently achieved healthy state into a permanent homeostatic set-point. This hypothesis is supported by existing evidence and generates specific predictions for empirical testing.

References

1. [△]Morton GJ, Cummings DE, Baskin DG, et al. (2006). "Central Nervous System Control of Food Intake and Body Weight." *Nature*. **443**:289–295.
2. [△]Apkarian AV, Bushnell MC, Treede RD, Zubieta JK (2005). "Human Brain Mechanisms of Pain Perception and Regulation in Health and Disease." *Eur J Pain*. **9**(4):463–484.
3. [△]Sjöström L (2013). "Review of the Key Results from the Swedish Obese Subjects (SOS) Trial." *Proc Nutr Soc*. **72**(3):237–243.
4. [△]Diekelmann S, Born J (2010). "The Memory Function of Sleep." *Nat Rev Neurosci*. **11**(2):114–126.
5. [△]Nader K, Hardt O (2009). "A Single Standard for Memory: The Case for Reconsolidation." *Nat Rev Neurosci*. **10**(3):224–234.
6. ^{a, b}Han JS (2004). "Acupuncture and Endorphins." *Neurosci Lett*. **361**(1–3):258–261.
7. [△]Liu S, Wang Z, Su Y, et al. (2021). "The Neurobiological Mechanisms of Acupuncture: From Peripheral to Central Modulation." *Front Neurosci*. **15**:698215.
8. [△]McGaugh JL (2002). "Memory Consolidation and the Amygdala: A Systems Perspective." *Trends Neurosci*. **25**(9):456–461.
9. ^{a, b}Cheng K, Tian S, Zhang Y, et al. (2014). "Chronobiology of Acupuncture: Time-Dependent Effects on HPA Axis and Autonomic Function." *Evid Based Complement Alternat Med*. **2014**:987365.

10. [△]Han JS (2003). "Acupuncture: Neuropeptide Release Produced by Electrical Stimulation of Different Frequencies." *Trends Neurosci.* **26**(1):17–22.
11. [△]Chen L, Zhang J, Li F, et al. (2009). "Endocannabinoids and Acupuncture Analgesia." *Acupunct Med.* **27**(3): 107–112.
12. [△]Yang J, Lin Y, Hu L, et al. (2017). "Oxytocin Mediates Acupuncture Analgesia in the Rat Model of Chronic Inflammatory Pain." *Neuropeptides.* **61**:47–54.
13. [△]Bramham CR (1992). "Opioid Receptor Dependent Long-Term Potentiation: The Role of β -Endorphin." *Synapse.* **11**(1):43–51.
14. [△]Kano M, Ohno-Shosaku T, Hashimotodani Y, et al. (2009). "Endocannabinoid-Mediated Control of Synaptic Transmission." *Physiol Rev.* **89**(1):309–380.
15. [△]Chini B, Leonzino M, Braidà D, Sala M (2014). "Learning About Oxytocin: Pharmacologic and Behavioral Aspects." *Biol Psychiatry.* **76**(5):365–373.
16. [△]Park HJ, Chae Y, Jang J, et al. (2012). "Acupuncture Stimulation Alleviates Corticosterone-Induced Impairments of Spatial Memory and Cholinergic Neurons in Rats." *Evid Based Complement Alternat Med.* **2012**:670 536.
17. [△]Wang F, et al. (2025). "Acupuncture Ameliorates Diet-Induced Obesity via the Vagal–GLP-1–ARC Circuit: Neural Mechanism of Anorexigenic Action." *Chin Med.* **20**:12.
18. [△]Haker E, Egekvist H, Bjerring P (2000). "Effect of Sensory Stimulation (Acupuncture) on Sympathetic and Parasympathetic Activities in Healthy Subjects." *J Auton Nerv Syst.* **79**(1):52–59.
19. [△]Li DZ, Zhou Y, Yang YN, et al. (2014). "Acupuncture for Essential Hypertension: A Meta-Analysis." *Am J Hypertens.* **27**(7):919–928.
20. [△]Wang F, Tian DR, Han JS (2015). "Electroacupuncture in the Treatment of Obesity." *CNS Neurosci Ther.* **21** (8):617–625.
21. [△]Wang Y, Li Q, Zhang X, et al. (2022). "Memory-Directed Acupuncture as a Neuromodulatory Treatment for PTSD: Theory, Clinical Model and Case Studies." *Transl Psychiatry.* **12**:137.

Declarations

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.