

Research Article

On Bundles of Varieties V_2^3 in $PG(4, q)$ and Their Codes

Rita Vincenti¹

1. University of Perugia, Italy

In this note we use the spatial representation in $\Sigma = PG(4, q)$ of the projective plane $\Pi = PG(2, q^2)$, by fixing a hyperplane Σ' with a regular spread \mathcal{S} of lines. We consider a bundle \mathcal{X} of varieties V_2^3 of Σ having in common the $q + 1$ points of a conic \mathcal{C}^2 of a plane π_0 , $\pi_0 \cap \Sigma' = l_0 \in \mathcal{S}$, thus representing an affine line of Π , and a further affine point $O \notin \pi_0$. This subset \mathcal{X} of Σ represents a bundle of non-affine Baer subplanes of Π , each of them having one point at infinity (corresponding to a line of \mathcal{S}), having in common a subline of affine points of Π and a further affine point. Then \mathcal{X} is considered as a projective system of Σ and, by using such a representation of Π , we can calculate the ground parameters of the code $C_{\mathcal{X}}$ arising from it.

Corresponding author: Rita Vincenti, aliceiw213@gmail.com

Mathematics Subject Classification: 51E20, 51A05, 94B05, 94B27

1. Introduction

It is known that a projective translation plane Π of order $n = q^2$ of dimension 2 over its kernel $F = GF(q)$ can be represented by a 4-dimensional projective space $\Sigma = PG(4, q)$ over F , fixing a hyperplane $\Sigma' = PG(3, q)$ and a spread \mathcal{S} of lines of Σ' . The points of Π are represented by (i) the points of $\Sigma \setminus \Sigma'$ and (ii) the lines of \mathcal{S} . The lines of Π are represented by (i) the planes α of $\Sigma \setminus \Sigma'$ such that $\alpha \cap \Sigma'$ belongs to \mathcal{S} and by (ii) the spread \mathcal{S} . The translation line l of Π is represented by \mathcal{S} (cf. [1]).

A Baer subplane B of Π has order q and it is *dense* in the sense that a line of Π either is a line of B (that is, meets B in a subline of $q + 1$ points, such a subplane is *affine*) or it meets B in one point (such a subplane is *non-affine*).

The *affine* Baer subplanes B of Π are represented by the *transversal* planes β to \mathcal{S} , that is, the planes of $\Sigma \setminus \Sigma'$ such that the line $\beta \cap \Sigma' \notin \mathcal{S}$ meets $q + 1$ lines of \mathcal{S} . In such a way l is a line of B (cf. [2], pp. 68-72). Of course all that holds also in case Π is the Desarguesian plane $PG(2, q^2)$ when \mathcal{S} is a regular spread (cf. [3], [2]).

A variety V_2^3 of Σ with a line l_∞ in \mathcal{S} as the minimum (linear) order directrix, a conic \mathcal{C}^2 as a 2nd order directrix with $\mathcal{C}^2 \subset \pi_0$, $\pi_0 \cap \Sigma' = l_0 \in \mathcal{S} \setminus l_\infty$ and $\mathcal{C}^2 \cap l_0 = \emptyset$, represents a non-affine Baer subplane of Π having one point on the translation line l and the *subline* \mathcal{C}^2 of the line π_0 (cf. [3]).

In this paper we consider bundles of $q + 1$ varieties V_2^3 of $\Sigma = PG(4, q)$ with the linear directrix in \mathcal{S} and having in common a same conic \mathcal{C}^2 as a 2nd order directrix and one further affine point. By using the spatial representation of $\Pi = PG(2, q^2)$ in $PG(4, q)$, we can characterize such a bundle \mathcal{X} from the intersection point of view, construct a linear code $\mathcal{C}_\mathcal{X}$ arising from it and show that its ground parameters allow $\mathcal{C}_\mathcal{X}$ to correct an enough large number of errors.

2. Preliminary Notes

Let $F = GF(q)$ be a finite field, $q = p^s$, p prime. Denote F^{r+1} the $(r + 1)$ -dimensional vector space over F , $P^r = PrF^{r+1} = PG(r, q)$ the r -dimensional projective space contraction of F^{r+1} over F . Let \overline{F} be the algebraic closure of the field $F = GF(q)$.

Denote S_t with $t \geq 2$ a subspace of P^r of dimension t . A hyperplane S_{r-1} will be denoted also by H , a plane by π .

The geometry P^r is considered a sub-geometry of \overline{P}^r , the projective geometry over \overline{F} . We refer to the points of P^r as the *rational points* of \overline{P}^r .

Definition 2.1. A variety V_u^v of dimension u and of order v of P^r is the set of the rational points of a projective variety \overline{V}_u^v of \overline{P}^r defined by a finite set of polynomials with coefficients in the field F .

From [4], p.290, 7.- for $r \geq 4$ follows

Lemma 2.2. The ruled variety V_2^{r-1} of $PG(r, q)$ is generated by the lines connecting the corresponding points of two birationally (or, projectively) equivalent curves in two complementary subspaces, of order m and $r - 1 - m$, respectively. It has order the sum of the orders of the curves as there are no fixed points.

Let P^4 be the projective geometry $PG(4, q)$.

Lemma 2.3. A variety V_2^3 of $PG(4, q)$ is obtained by joining the corresponding points of a directrix line l and a directrix conic \mathcal{C} in a plane π , l and \mathcal{C} being projectively equivalent and with $l \cap \pi = \emptyset$.

Proof. See [5] p. 90.

Choose a coordinate system in P^4 so that it is a coordinate system for \overline{P}^4 too, denote a point $P \approx (x_1, x_2, y_1, y_2, t) := \overline{F}^*(x_1, x_2, y_1, y_2, t)$, $\overline{F}^* = \overline{F} \setminus \{0\}$.

P is a rational point if there exists $(x_1, x_2, y_1, y_2, t) \in F^5$ such that $P \approx (x_1, x_2, y_1, y_2, t)$. A variety V of P^4 is the set of the rational points of \overline{P}^4 solutions of a finite set of polynomials of $F[x_1, x_2, y_1, y_2, t]$.

Lemma 2.4. The variety V_2^3 can be represented as the definite intersection of two quadrics of $PG(4, q)$, that is, the cone of planes $\mathcal{Q}_1 : sx_2^2 - x_1^2 - sx_2t = 0$ (where s is a non square of $GF(q)$) and the cone of planes $\mathcal{Q}_2 : x_1y_1 - x_2y_2 = 0$. The plane $\pi' : x_1 = 0, x_2 = 0$ is contained in both quadrics so that, by Bezout, the order of the intersection variety is $4 - 1 = 3$.

Proof. See [3] Theorem 1.1, [5] p. 92.

Let $\Pi = PG(2, q^2)$ be the Desarguesian plane over $GF(q^2)$. Denote l the line at infinity of Π . In the spatial representation of Π in $P^4 = PG(4, q)$ fix a hyperplane $\Sigma' = PG(3, q)$ and a regular spread \mathcal{S} of lines of Σ' , where $|\mathcal{S}| = q^2 + 1$.

Lemma 2.5. The points of Π are represented by (i) the points of $\Sigma \setminus \Sigma'$ (the affine points of Π) and by (ii) the lines of \mathcal{S} (the points at infinity of Π). The lines of Π are represented by (i) the planes α of $\Sigma \setminus \Sigma'$ such that $\alpha \cap \Sigma'$ belongs to \mathcal{S} and by (ii) the spread \mathcal{S} , representing the line at infinity l .

Proof. See [1] the Bruck and Bose representation and [2], p. 775.

Definition 2.6. A Baer subplane of $\Pi = PG(2, q^2)$ is an affine subplane if it meets the line at infinity l of Π in a subline l_1 , it is a non-affine subplane if it meets the line l in one point.

Lemma 2.7.

- (i) Two affine Baer subplanes of Π having in common the subline l_1 can meet in at most one further point.
- (ii) The Baer subplanes having in common only a subline l_1 are q^2 .
- (iii) The Baer subplanes having in common a subline l_1 and one further point are $q + 1$.

Proof. (i) Two Baer subplanes having in common a subline l_1 and two further points coincide, because they have in common at least four reference (three by three non collinear) points.

Without loosing generality, we can consider two affine Baer subplanes \mathcal{B} and \mathcal{B}' of Π having in common a subline l_1 of l . In the spatial representation of Π , they are represented by two planes B and B' of P^4 ,

respectively, such that the lines $B \cap \Sigma' = r$ and $B' \cap \Sigma' = r'$ are transversal lines of the same regulus $\mathcal{R} \subset \mathcal{S}$. Denote \mathcal{R}' the opposite regulus to \mathcal{R} .

There are two cases:

(ii) If $r = r'$, the planes B and B' have in common the line r meeting the regulus \mathcal{R} in its $q + 1$ lines so that the subplanes \mathcal{B} and \mathcal{B}' have in common the subline l_1 (represented by \mathcal{R}) of the line l (represented by \mathcal{S}) and no further (affine) points. Such planes are $\frac{q^4}{q^2} = q^2$ and represent q^2 affine Baer subplanes of Π having in common only the subset l_1 of $q + 1$ points of the line at infinity l .

(iii) If $r \neq r'$, the planes B and B' have in common an affine point $O \in \Sigma \setminus \Sigma'$ so that the two subplanes \mathcal{B} and \mathcal{B}' meet along the subline l_1 represented by \mathcal{R} and in the affine point O . The regulus \mathcal{R} has $q + 1$ transversal lines $\{t_i | i = 1, \dots, q + 1\}$ belonging to \mathcal{R}' . Each space $O \oplus t_i$ is a transversal plane τ_i , so that $\{\tau_i | i = 1, \dots, q + 1\}$ represent the $q + 1$ affine Baer subplanes of Π having in common l_1 and the affine point O .

Choose and fix a line l_∞ of the (regular) spread \mathcal{S} , a plane π_0 such that $\pi_0 \cap \Sigma' = l_0 \in \mathcal{S} \setminus l_\infty$ and a non-degenerate conic $\mathcal{C}^2 \subset \pi_0 \setminus l_0$. Let Λ be a projectivity between l_∞ and \mathcal{C}^2 . Denote V_2^3 the variety arising by connecting corresponding points of l_∞ and \mathcal{C}^2 via Λ (cf. [5], p. 90).

Lemma 2.8. *The variety V_2^3 represents a non-affine Baer subplane of Π meeting the line at infinity l in the point l_∞ and containing the subline \mathcal{C}^2 of the line represented by π_0 .*

Proof. See [3] and [2].

Let F^n be the n -dimensional vector space over $F = GF(q)$.

Definition 2.9. A linear $[n, k]_q$ -code C of length n is a k -dimensional subspace of the vector space F^n .

Definition 2.10. An $[n, k]_q$ -projective system \mathcal{X} is a set of n non necessarily distinct points of the projective geometry $PrF^k = PG(k - 1, q)$. It is non-degenerate if these points are not contained in a hyperplane (cf. [6], p. 2).

Assume that \mathcal{X} consists of n distinct points having maximum rank.

Codes and projective systems are linked by a strict connection one can read in [6], so that from subsets \mathcal{X} of a projective geometry linear codes $C_{\mathcal{X}}$ can be generated. More precisely, for each point of \mathcal{X} choose a generating vector. Denote \mathcal{M} the matrix having as rows such n vectors and let $C_{\mathcal{X}}$ be the linear code having \mathcal{M}^t as a generator matrix. The code $C_{\mathcal{X}}$ is the k -dimensional subspace of F^n which is the image of the mapping from the dual k -dimensional space $(F^k)^*$ onto F^n that calculates every linear form over the

points of \mathcal{X} . Hence the length n of codeword of $C_{\mathcal{X}}$ is the cardinality of \mathcal{X} , the dimension of $C_{\mathcal{X}}$ being just k (cf. [6], p. 3).

Denote \mathcal{H} the set of all hyperplanes of $P^{k-1} = P\mathcal{F}^k$.

There exists a natural 1-1 correspondence between the equivalence classes of a non-degenerate $[n, k]_q$ -projective system \mathcal{X} and a non-degenerate $[n, k]_q$ -code $C_{\mathcal{X}}$ such that if \mathcal{X} is an $[n, k]_q$ -projective system and $C_{\mathcal{X}}$ is a corresponding code, then the non-zero codewords of $C_{\mathcal{X}}$ correspond to hyperplanes $H \in \mathcal{H}$, up to a non-zero factor. The correspondence preserves the ground parameters.

The weight of a codeword c corresponding to the hyperplane H_c is the number of points of $\mathcal{X} \setminus H_c$, thus the minimum weight (or, the minimum distance) d of the code $C_{\mathcal{X}}$ is $d = |\mathcal{X}| - \max\{|\mathcal{X} \cap H| \mid H \in \mathcal{H}\}$. Therefore in order to find the minimum distance of the code $C_{\mathcal{X}}$ it needs to calculate the maximum intersection of \mathcal{X} with the hyperplanes of \mathcal{H} .

A linear code with length n , dimension k and minimum distance d over the field $F = GF(q)$ can be denoted also as an $[n, k, d]_q$ -code.

If C is an $[n, k, d]_q$ -code, then C is an s -error-correcting code for all $s \leq \lfloor \frac{d-1}{2} \rfloor$. We call $t = \lfloor \frac{d-1}{2} \rfloor$ the error-correcting capability of C (cf. [6], p.3).

3. Main Results

With the notations of the previous section, choose and fix the line $l_0 \in \mathcal{S}$, the plane π_0 such that $\pi_0 \cap \Sigma' = l_0 \in \mathcal{S}$ and the non-degenerate conic $\mathcal{C}^2 \subset \pi_0 \setminus l_0$.

Denote Σ'' a hyperplane of $\Sigma = PG(4, q)$ containing the plane π_0 . Let $\pi = \Sigma'' \cap \Sigma'$. The plane π contains the line l_0 and each of the q^2 points of $\pi \setminus l_0$ belongs to one of the q^2 lines of $\mathcal{S} \setminus \{l_0\}$. Let O be a point, $O \in \Sigma'' \setminus \{\pi_0 \cup \pi\}$. Denote \mathcal{Q} the quadric cone having vertex the point O and directrix the conic \mathcal{C}^2 . Let $\mathcal{C}'^2 = \mathcal{Q} \cap \pi$. Obviously \mathcal{C}'^2 is a non-degenerate conic with $\mathcal{C}'^2 \cap l_0 = \emptyset$.

Let $\{R_i \mid i = 1, \dots, q+1\}$ be the set of the $q+1$ points of \mathcal{C}^2 , $\{r_i \mid i = 1, \dots, q+1\}$ the $q+1$ lines of the cone \mathcal{Q} with $R_i \in r_i$, $\{R'_i = r_i \cap \mathcal{C}'^2 \mid i = 1, \dots, q+1\}$ the corresponding set of $q+1$ points of \mathcal{C}'^2 with $R'_i \in r_i$, $\{s_i \mid i = 1, \dots, q+1\}$ the $q+1$ lines of \mathcal{S} with $\{R'_i \in s_i \mid i = 1, \dots, q+1\}$.

For each line s_i let λ_i be a projectivity between s_i and \mathcal{C}^2 such that $\lambda_i(R'_i) = R_i$

Denote S_i the point at infinity of the plane Π represented by the line $s_i \in \mathcal{S}$, p_0 the line of Π represented by the plane π_0 and c_2 the subline of p_0 corresponding to \mathcal{C}^2 .

Let \mathcal{V}_i be the variety V_2^3 having the conic \mathcal{C}^2 and the line s_i as directrices constructed via λ_i . Note that, by construction, the line r_i is one of the $q + 1$ generatrix lines of \mathcal{V}_i .

From Lemma 2.8 follows that each of the $q + 1$ variety \mathcal{V}_i is a non-affine Baer subplane of Π meeting the line l in the point S_i , containing $c_2 \subset p_0$ and the point O .

Define $\mathcal{V} := \bigcup_i \mathcal{V}_i$ the union of the points of all varieties \mathcal{V}_i for all $i = 1, \dots, q + 1$.

Lemma 3.1. \mathcal{V} represents the bundle of the full set of $q + 1$ non-affine Baer subplanes having in common the subline c_2 and the point O .

Proof. See (iii) of Lemma 2.7 and [3].

Proposition 3.2. $\Sigma'' \cap \mathcal{V} = \mathcal{Q}$.

Proof. By construction the hyperplane Σ'' contains \mathcal{Q} . As for any variety \mathcal{V}_i , $\Sigma'' \cap \mathcal{V}_i$ cannot contain the directrix line s_i (otherwise $\Sigma'' = \Sigma'$), then Σ'' meets \mathcal{V}_i at most in a cubic curve $\mathcal{C}^2 \cup r_i$ (cf. [5], (ii), p. 93).

Assume $\Sigma'' \cap \mathcal{V}$ contains $\mathcal{C}^2 \cup r_i \subset \mathcal{V}_i$ and a further point $P_j \in V_j$ with $j \neq i$. Hence Σ'' contains the line $r = P_j R_j \in \mathcal{V}_j$ with $R_j \in \mathcal{C}^2$. If $r \neq r_j$, then Σ'' should meet \mathcal{V}_j in $\mathcal{C}^2 \cup r_j \cup r$ where r_j and r are two generatrix lines of \mathcal{V}_j , then the line s_j should belong to Σ'' , a contradiction (cf. [5], (ii), p. 93). Hence $\Sigma'' \cap \mathcal{V} = \mathcal{Q}$.

Denote $\mathcal{V}_{aff} = \mathcal{V} \setminus \Sigma'$.

Proposition 3.3.

(i) A hyperplane of Σ having maximum intersection with \mathcal{V} is Σ' , and $\Sigma' \cap \mathcal{V}$ consists of the points of the lines $\{s_i | i = 1, \dots, q + 1\} \subset \mathcal{S}$.

(ii) A hyperplane of Σ having maximum intersection with \mathcal{V}_{aff} is Σ'' and $\Sigma'' \cap \mathcal{V}_{aff}$ consists of the points of $\mathcal{Q} \setminus \mathcal{C}^2$.

Proof. (i) Let $H \in \mathcal{H}$ a hyperplane. If $H = \Sigma'$ then $H \cap \mathcal{V}$ is the set of the $(q + 1)^2$ points of $\{s_i | i = 1, \dots, q + 1\} \subset \mathcal{S}$. If $H = \Sigma''$ then $H \cap \mathcal{V}$ is the set of the $q^2 + q + 1$ points of \mathcal{Q} .

Let $H \neq \Sigma', \Sigma''$.

Denote $H \cap \Sigma' = \pi'$, $H \cap \Sigma'' = \pi''$.

For H there are two possibilities: 1) H contains π_0 , 2) H does not contain π_0 .

1) It is $\pi'' = \pi_0$ so that it contains \mathcal{C}^2 . Moreover $\pi' \neq \pi$ otherwise $H = \Sigma''$. The plane π' forms bundle with axis the line l_0 with π_0 and π . Each point of π' belongs to one line of $\mathcal{S} \setminus l_0$ then it meets the

$q + 1$ points $\{P_i = \pi' \cap s_i | i = 1, \dots, q + 1\}$. Therefore $H \cap \mathcal{V}$ contains at least the $q + 1$ points P_i and the points of \mathcal{C}^2 . Then $|H \cap \mathcal{V}| \geq 2(q + 1)$. The maximum intersection is reached if each line $P_i R_i$ coincides with one generatrix line of the variety \mathcal{V}_i for every i , In such a case $|H \cap \mathcal{V}| = (q + 1)^2$.

2) Let $\pi'' \cap \Sigma' = l$. Then l is a line of π' too.

Let $l = l_0$. The plane π'' contains no generatrix line of the varieties \mathcal{V}_i otherwise l_0 would meet some line s_i , it meets \mathcal{V} in at most a conic \mathcal{C}_Q of \mathcal{Q} . Set $\{P_i \in \mathcal{C}_Q | i = 1, \dots, q + 1\}$.

If $\pi' = \pi$, then $\pi' \cap \mathcal{V} = \mathcal{C}'^2$. If $\pi' \neq \pi$, then it contains no line s_i (otherwise $l_0 \cap s_i \neq \emptyset$), it can meet at most $q + 1$ lines s_i in points T_i . In both cases the maximum intersection is reached if the $q + 1$ lines $P_i R'_i$, or $P_i T_i$, respectively, coincide with the generatrix lines of the varieties \mathcal{V}_i . Hence $|H \cap \mathcal{V}| \leq (q + 1)^2$.

Let $l \neq l_0$. Denote $r' = \pi'' \cap \pi_0$. Then $l = s_i$ for some i or l meets at most $q + 1$ lines s_i .

If $\pi' = \pi$, it contains the $q + 1$ points of \mathcal{C}'^2 and according to r' is secant, tangent or external to the conic \mathcal{C}^2 , $|H \cap \mathcal{V}|$ is less or equal to $(q + 1) + 2q = 3q + 1$, $(q + 1) + q = 2q + 1$ or $q + 1$, respectively.

Assume $\pi' \neq \pi$. The plane π' must contain one line t of \mathcal{S} and the q^2 points of the remaining lines of \mathcal{S} . Then the plane π' contains the $q + 1$ points of $t = s_i$ for some i , or the $q + 1$ points of the set $\{s_i \cap \pi' | i = 1, \dots, q + 1\} \subset \mathcal{V}$.

According to r' is secant, tangent or external to the conic \mathcal{C}^2 , H meets \mathcal{V} in 2 generatrix lines, in 1 generatrix line or in no generatrix line. Therefore $|H \cap \mathcal{V}|$ is less or equal to $(q + 1) + 2q = 3q + 1$, $(q + 1) + q = 2q + 1$ or $q + 1$.

Hence the maximum intersection a hyperplane can have with \mathcal{V} consists of $(q + 1)^2$ points. Σ' is one of such hyperplanes.

(ii) Let H be a hyperplane, $H \neq \Sigma'$. From [7], Lemma 11, it is known the maximum intersection a hyperplane of Σ has with a variety V_2^3 consists of two generatrix lines and the directrix line. Of course H cannot meet two different varieties in such a way otherwise H , containing two lines of \mathcal{S} would coincides with Σ' . Therefore H can meet at least q varieties along the conic \mathcal{C}^2 and one generatrix line for each variety, then q points of the conic \mathcal{C}'^2 . In any case H contains O then the cone \mathcal{Q} . Therefore $H = \Sigma''$. Hence the maximum intersection a hyperplane can have with \mathcal{V}_{aff} is $\mathcal{Q} \setminus \mathcal{C}'^2$ with $|\mathcal{Q} \setminus \mathcal{C}'^2| = q^2$.

Denote $\mathcal{X} := \mathcal{V}$ the projective system defined by \mathcal{V} , $C_{\mathcal{X}}$ the linear code arising from \mathcal{X} , $\mathcal{X}_{aff} := \mathcal{V}_{aff}$ the projective system defined by \mathcal{V}_{aff} , $C_{\mathcal{X}_{aff}}$ the linear code arising from \mathcal{X}_{aff} .

Theorem 3.4.

(i) $C_{\mathcal{X}}$ is an $[n, k, d]_q$ -code with $n = q^3 + 2q^2 + q + 1$, $k = 5$, $d = q^3 + q^2 - q$.

(ii) $C_{\mathcal{X}_{aff}}$ is an $[n', k, d']_q$ -code with $n' = q^3 + q^2 - q$, $k = 5$, $d' = q^3 - q$.

Proof. (i) Each variety \mathcal{V}_i consists of $q + 1$ skew lines, hence it has $(q + 1)^2$ points. Every two varieties \mathcal{V}_i and \mathcal{V}_j have in common the conic \mathcal{C}^2 and the point O so that for each variety remain $q^2 + 2q + 1 - (q + 1) - 1 = q^2 + q - 1$ points. The varieties are $q + 1$ so that the cardinality of \mathcal{X} is $(q^2 + q - 1)(q + 1) = q^3 + 2q^2 - 1$ plus the point O and the $(q + 1)$ points of the conic \mathcal{C}^2 . Hence $|\mathcal{X}| = q^3 + 2q^2 + q + 1$. The length of the code $C_{\mathcal{X}}$ is therefore $n = q^3 + 2q^2 + q + 1$.

The dimension of $C_{\mathcal{X}}$ is obviously 5, that is, the vector dimension of Σ .

From Proposition 3.3, (i), follows the distance of $C_{\mathcal{X}}$ is $d = n - |\{P \in s_i | i = 1, \dots, q + 1\}|$ that is, $d = q^3 + 2q^2 + q + 1 - (q^2 + 2q + 1) = q^3 + q^2 - q$.

(ii) The length of the code $C_{\mathcal{X}_{aff}}$ equals $n' = |\mathcal{X}| - |\{P \in s_i | i = 1, \dots, q + 1\}| = q^3 + 2q^2 + q + 1 - (q^2 + 2q + 1) = q^3 + q^2 - q$. Its dimension is $k = 5$. From Proposition 3.3, (ii), follows the distance is $d' = n' - |\mathcal{Q} \setminus \mathcal{C}^2|$ that is, $d' = q^3 + q^2 - q - q^2 = q^3 - q$.

Examples

For $q = 2$, $C_{\mathcal{X}}$ is a $[19, 5, 10]_2$ -code and it can correct at most $\lfloor \frac{10-1}{2} \rfloor = 4$ errors. For $q = 3$, $C_{\mathcal{X}}$ is a $[49, 5, 33]_3$ -code and it can correct at most $\lfloor \frac{33-1}{2} \rfloor = 16$ errors.

For $q = 2$, $C_{\mathcal{X}_{aff}}$ is a $[10, 5, 6]_2$ -code and it can correct at most $\lfloor \frac{6-1}{2} \rfloor = 2$ errors. For $q = 3$, $C_{\mathcal{X}_{aff}}$ is a $[33, 5, 24]_3$ -code and it can correct at most $\lfloor \frac{24-1}{2} \rfloor = 11$ errors.

Other References

- R. Vincenti, Fibrazioni di un $S_{3,q}$ indotte da fibrazioni di un S_{3,q^2} e rappresentazione di sottopiani di Baer di un piano proiettivo, Atti e Mem. Acc. Sci. Lett. e Arti di Modena, Serie VI, Vol. XIX, (1977), 1-18.
- R. Vincenti, On some classical varieties and codes, Technical Report 2000/20, Department of Mathematics and Computer Science, University of Perugia (Italy).

References

1. ^a_bR. H. Bruck, R. C. Bose, *Linear representation of projective planes in projective spaces*, *J.of Algebra*, 4, (1966), 117–172.
2. ^a_b_c ^dR. Vincenti, *A survey on varieties of PG (4,q) and Baer subplanes of translation planes*, *Annals of Discrete Math.*, N.H. Pub. Co., 18, (1983), 775–780.
3. ^a_b_c ^d_eR. Vincenti, *Alcuni tipi di varietà a V₂₃ di S_{4,q} e sottopiani di Baer*, *Algebra e Geometria Suppl. BUMI*, Vol. 2, (1980), 31–44.
4. ^ΔE. Bertini, *Introduzione alla geometria proiettiva degli iperspazi*, (1907), Enrico Spoerri Editore, Pisa.
5. ^a_b_c ^d_eR. Vincenti, *Finite fields, projective geometry and related topics*, Morlacchi Editore, (2021), ISBN 978-88-9392-259.
6. ^a_b_c ^dR. Vincenti, *Varieties and codes from partial ruled sets*, *International Mathematical Forum*, Vol. 18, n. 1, (2023), 1–14, doi.org/10.12988/imf.2023.912353.
7. ^ΔR. Vincenti, *Linear codes from projective varieties: a survey*, March 21st, 2023, *Article on the IntechOpen Edited Book Coding Theory Essentials*, by D. G. Harkut, (2023), doi: 10.5772/intechopen.109836

Declarations

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.