

Pathologic Stage IIIC Cutaneous Melanoma AJCC v8

National Cancer Institute

Source

National Cancer Institute. *Pathologic Stage IIIC Cutaneous Melanoma AJCC v8*. NCI Thesaurus. Code C137673.

Stage IIIC includes: (T0, N2b, N2c, N3b or N3c); (T1a-T3a, N2c or N3a/b/c, M0); (T3b/T4a, Any N greater than or equal to N1, M0); (T4b, N1a-N2c, M0). T0: No evidence of primary tumor (e.g., unknown primary or completely regressed melanoma). T thickness: Not applicable. Ulceration status: Not applicable. T1a: Tumor measuring less than 0.8 mm in thickness. Ulceration status: Without ulceration. T1b: Tumor measuring less than 0.8 mm in thickness with ulceration, or 0.8-1.0 mm with or without ulceration. T2a: Tumor measuring more than 1.0 and equal to or less than 2.0 mm in thickness. Ulceration status: Without ulceration. T2b: Tumor measuring more than 1.0 and equal to or less than 2.0 mm in thickness. Ulceration status: With ulceration. T3a: Tumor measuring more than 2.0 and equal to or less than 4.0 mm in thickness. Ulceration status: Without ulceration. T3b: Tumor measuring more than 2.0 and equal to or less than 4.0 mm in thickness. Ulceration status: With ulceration. T4a: Tumor measuring more than 4.0 mm in thickness. Ulceration status: Without ulceration. T4b: Tumor measuring more than 4.0 mm in thickness. Ulceration status: With ulceration. N2b: Two or three nodal metastases at least one of which was clinically detected. Presence of in-transit, satellite, and/or microsatellite metastases: No. N2c: One clinically occult nodal metastasis. Presence of in-transit, satellite, and/or microsatellite metastases: Yes. N3b: Four or more nodal metastases, at least one of which was clinically detected, or presence of any number of matted nodes. Presence of in-transit, satellite, and/or microsatellite metastases: No. N3c: Two or more clinically occult nodal metastases. Presence of in-transit, satellite, and/or microsatellite metastases: Yes. N3a: Four or more clinically occult nodal metastasis (i.e., detected by sentinel lymph node biopsy). Presence of in-transit, satellite, and/or microsatellite metastases: No. N1a: One clinically occult nodal metastasis (i.e., detected by sentinel lymph node biopsy). Presence of in-transit, satellite, and/or microsatellite metastases: No. N1b: One clinically detected nodal metastasis.

Presence of in-transit, satellite, and/or microsatellite metastases: No. N1c: No regional lymph node disease. Presence of in-transit, satellite, and/or microsatellite metastases: Yes. N2a: Two or three clinically occult nodal metastases (i.e., detected by sentinel lymph node biopsy). Presence of in-transit, satellite, and/or microsatellite metastases: No. M0: No evidence of distant metastasis. LDH level is not applicable. (AJCC 8th ed.)